

**Action principles, restoration of BRS symmetry and the
renormalization group equation for chiral non-Abelian gauge theories
in dimensional renormalization with a non-anticommuting γ_5**

C. P. Martín* and D. Sánchez-Ruiz†

Departamento de Física Teórica I, Universidad Complutense, 28040 Madrid, Spain

The one-loop renormalization of a general chiral gauge theory without scalar and Majorana fields is fully worked out within Breitenlohner and Maison dimensional renormalization scheme. The coefficients of the anomalous terms introduced in the Slavnov-Taylor equations by the minimal subtraction algorithm are calculated and the asymmetric counterterms needed to restore the BRS symmetry, if the anomaly cancellation conditions are met, are computed. The renormalization group equation and its coefficients are worked out in the anomaly free case. The computations draw heavily from the existence of action principles and BRS cohomology theory.

1. Introduction

Dimensional Regularization [1,2] is the standard regularization method in four dimensional perturbative quantum field theory as applied to particle physics. Both its axiomatics and properties were rigorously established long ago [3,4,5,6,7,8,9,10] and quite a number of computational techniques based on the method have been developed over the years [11,12]. Involved multiloop computations of the parameters of the Standard Model, a must due to the availability of high precision tests of the model in particle accelerators, have been carried out in the simplest possible setting thanks to dimensional regularization [13]. The success of Dimensional Regularization as a practical regularization method stems from the fact that it preserves the BRS symmetry of vector-like non-supersymmetric gauge theories without distorting neither the shape of the integrand of a regularized Feynman diagram nor the properties of the algebraic objects involved (no chiral objects being present). The minimal subtraction scheme, the famous $\overline{\text{MS}}$ scheme [11], then leads [2] to a renormalized BRS-invariant theory [3,6]. Counterterms are generated by multiplicative renormalization of the tree-level Lagrangian, so that β functions and anomalous dimensions can be computed easily [10].

It is an empirical fact that electroweak interactions are chiral and hence vector-like gauge theories does fail to account for them. It turn thus out that chiral gauge theories are of key importance to understanding Nature. Unfortunately, Dimensional Regularization loses its smartness when applied to chiral gauge theories. The algebraic properties

* email: carmelo@elbereth.fis.ucm.es

† email: domingos@eucmos.sim.ucm.es

that the matrix γ_5 has in four dimensions cannot be maintained without introducing algebraic inconsistencies as we move away from four dimensions [14]. Hence, within the framework of Dimensional Regularization, the definition of the object γ_5 in “ d complex dimensions” demands a new set of algebraic identities. There exists such a set of identities, they were introduced in ref. [3], following refs. [2,15], and they entail that the object γ_5 anticommutes no longer with the object γ_μ . The axiomatics of dimensional regularization so established is the only one which has been shown to be thoroughly consistent at any order in perturbation theory if cyclicity of the trace is not given up (see ref. [16] for the non-cyclic trace alternative). Minimal subtraction [11] of the singular part of the dimensionally regularized Feynman diagrams, a subtraction procedure known as minimal dimensional renormalization [3,4,5,6], leads to a renormalized quantum field theory that satisfies Hepp axioms [20] of renormalization theory. Field equations, the action principles and Zimmermann-Bonneau identities hold in dimensional renormalization. This has been rigorously shown in refs. [3,4,5,6,7,17] and it constitutes one of the key ingredients of the modern approach to the quantization of gauge theories by using BRS methods [18,19].

That Dimensional Regularization can be used along with algebraic BRS techniques is of the utmost importance, for this regularization method and the minimal subtraction algorithm that comes with it break, generally speaking, chiral gauge symmetries. This breaking gives rise to both physical and non-physical anomalies [7,21]. Physical anomalies in the currents of the Lagrangian impose anomaly cancellation conditions [23], lest the theory ceases to make sense as a quantum theory. These anomalies belong to the cohomology of the Slavnov-Taylor operator that governs the chiral gauge symmetry at the quantum level and they are not artifacts of the regularization method nor due to the renormalization scheme employed. Non-physical anomalies correspond to trivial objects in the cohomology of the BRS operator and, therefore, they can be set to zero by an appropriate choice of finite counterterms [21]. Non-physical anomalies are artifacts either of the regularization method, the renormalization method or both.

Notwithstanding the unique status that as a thoroughly consistent framework the Breitenlohner and Maison scheme enjoys among the dimensional renormalization prescriptions, there is, up to the best of our knowledge, no complete one-loop study of a general non-Abelian chiral gauge theory, let alone the Standard Model. Most of the implementations of dimensional regularization in quantum field theories which involve the matrix γ_5 and are anomaly-free, *e.g.* the Standard Model, take a fully anticommuting γ_5 in “ d complex dimensions”. This is the so-called “naive” prescription for the object γ_5 [22]. The purpose of this paper is to remedy this situation by carrying out such one-loop study for a general chiral gauge theory with neither scalar nor Majorana fields. The inclusion of scalar fields will be discussed elsewhere.

The layout of this paper is as follow. In Section 2 we give a quick account of the dimensional renormalization algorithm *à la Breitenlohner and Maison*, the Regularized and Quantum Action Principles and Bonneau identities. Section 3 is devoted to a thorough study of the one-loop dimensional renormalization of a general chiral gauge theory for a simple gauge group. In this section the BRS anomaly is computed for the first time in the Breitenlohner and Maison framework. Of course, the standard anomaly cancellation conditions are thus obtained. The beta function and the anomalous dimensions of the

theory are computed in this section as well. Again, the one-loop renormalization group equation for the model at hand had never been work out as yet in Breitenlohner and Maison scheme. In section 4 we adapt the results obtained in the previous section to the case of a non-simple gauge group. We also include three appendices. In Appendices A and B we show how to use the action principles to obtain the breaking of the Slavnov-Taylor identities. In Appendix C we compare, for a simple gauge group, the one-loop renormalized chiral gauge theory obtained by means of the Breitenlohner and Maison scheme with the corresponding one-loop renormalized theory obtained with the help of an fully anticommuting γ_5 in “ d complex dimensions”.

2. Dimensional renormalization: notation and general results

As explained in the Introduction, we will use the formulation of dimensional renormalization given by Breitenlohner and Maison [3,4,5], because it is a systematic and consistent procedure (valid in the sense of Hepp [20] to all orders in perturbation theory) in which tools like field equations, action principles and Ward identities can be rigorously implemented. This applies specially to the treatment of γ_5 .

We will also compare Breitenlohner and Maison formulation with the “naive” prescription which sets the object γ_5 to anticommute with the object γ_μ [22].

2.1. The Breitenlohner and Maison “ d -dimensional covariants”

Breitenlohner and Maison define the usual d -dimensional “Lorentz Covariants” ($g_{\mu\nu}$, p_μ , γ_μ , *etc.*) to be formal objects obeying the standard algebraic identities that they would satisfy in spaces of integral dimension [3] (d is a *complex* number, indices do not take any value and questions like of “Lorentz invariance” are meaningless!).

Besides the “ d -dimensional” metrics $g_{\mu\nu}$, they introduce a new one, $\hat{g}_{\mu\nu}$, which can be considered as a “ $(d-4)$ -dimensional covariant”. Moreover the ϵ tensor is considered to be a “4 dimensional covariant” object (because of axiom in eq. (2) below).

The symbols are required to obey (apart from obvious rules concerning contractions of indices, addition, multiplication by numbers, commutation of some symbols, *etc.*):

$$\begin{aligned} g^{\mu\nu} p_\nu &= p^\mu, & \hat{g}^{\mu\nu} p_\nu &= \hat{p}^\mu, & g^{\mu\nu} \gamma_\nu &= \gamma^\mu, & \hat{g}^{\mu\nu} \gamma_\nu &= \hat{\gamma}^\mu, \\ g_{\mu\nu} g^\nu{}_\rho &= g_{\mu\rho}, & \hat{g}_{\mu\nu} g^\nu{}_\rho &= \hat{g}_{\mu\nu} \hat{g}^\nu{}_\rho = \hat{g}_{\mu\rho}, \\ g^\mu{}_\mu &= d, & \text{Tr } \mathbb{I} &= 4, & \{\gamma^\mu, \gamma^\nu\} &= 2g^{\mu\nu} \mathbb{I}; \end{aligned} \tag{1}$$

$$\epsilon_{\mu_1 \dots \mu_4} \epsilon_{\nu_1 \dots \nu_4} = - \sum_{\pi \in S_4} \text{sign } \pi \prod_{i=1}^4 (g_{\mu_i \nu_{\pi(i)}} - \hat{g}_{\mu_i \nu_{\pi(i)}}), \tag{2}$$

where S_4 denotes the permutation group of 4 objects, and $\mathbb{I}$ is the unit of the symbolic algebra of gammas.

With the definitions

$$\bar{g}^{\mu\nu} = g^{\mu\nu} - \hat{g}^{\mu\nu}, \quad \bar{g}^{\mu\nu} p_\nu = \bar{p}^\mu, \quad \bar{g}^{\mu\nu} \gamma_\nu = \bar{\gamma}^\mu,$$

$$\gamma_5 = \frac{i}{4!} \epsilon_{\mu_1 \dots \mu_4} \gamma^{\mu_1} \dots \gamma^{\mu_4} \quad (3)$$

and the *assumption of cyclicity* of the symbol Tr , the following properties can be proved algebraically [3]

$$\begin{aligned} \hat{g}_{\mu\nu} \bar{g}^\nu{}_\rho &= 0, \quad \hat{p} \cdot \bar{q} = 0, \quad g_{\mu\nu} \bar{g}^\nu{}_\rho = \bar{g}_{\mu\rho}, \quad \hat{g}^\mu{}_\mu = d - 4, \quad \bar{g}^\mu{}_\mu = 4, \\ \{\bar{\gamma}^\mu, \bar{\gamma}^\nu\} &= 2\bar{g}^{\mu\nu} \mathbb{I}, \quad \{\hat{\gamma}^\mu, \hat{\gamma}^\nu\} = 2\hat{g}^{\mu\nu} \mathbb{I}, \\ \epsilon_{\mu_1 \dots \mu_4} &= \text{sign } \pi \epsilon_{\mu_{\pi(1)} \dots \mu_{\pi(4)}}, \\ \epsilon_{\mu_1 \dots \mu_4} \hat{g}^{\mu_i \nu_i} &= 0, \quad \epsilon_{\mu_1 \dots \mu_4} g^{\mu_i \nu_i} = \epsilon_{\mu_1 \dots \mu_4} \bar{g}^{\mu_i \nu_i}; \\ \text{Tr } \gamma^\mu &= \text{Tr } \gamma_5 = 0, \quad \gamma_5^2 = \mathbb{I}, \\ \text{Tr } [\gamma^\mu \gamma^\nu \gamma_5] &= \text{Tr } [\bar{\gamma}^\mu \bar{\gamma}^\nu \gamma_5] = \text{Tr } [\hat{\gamma}^\mu \hat{\gamma}^\nu \gamma_5] = \text{Tr } [\bar{\gamma}^\mu \hat{\gamma}^\nu \gamma_5] = 0, \\ \text{Tr } [\gamma^{\mu_1} \dots \gamma^{\mu_4} \gamma_5] &= \text{Tr } [\bar{\gamma}^{\mu_1} \dots \bar{\gamma}^{\mu_4} \gamma_5] = i \text{Tr } \mathbb{I} \epsilon_{\mu_1 \dots \mu_4}; \end{aligned} \quad (4)$$

$$\begin{aligned} \{\gamma_5, \gamma^\mu\} &= \{\gamma_5, \hat{\gamma}^\mu\} = 2\gamma_5 \hat{\gamma}^\mu = 2\hat{\gamma}^\mu \gamma_5, \\ [\gamma_5, \gamma^\mu] &= [\gamma_5, \bar{\gamma}^\mu] = 2\gamma_5 \bar{\gamma}^\mu = -2\bar{\gamma}^\mu \gamma_5, \\ \{\gamma_5, \bar{\gamma}^\mu\} &= [\gamma_5, \hat{\gamma}^\mu] = 0. \end{aligned} \quad (5)$$

$\bar{g}^{\mu\nu}$ can be thought of as a projector over the “4-dimensional space” and $\hat{g}^{\mu\nu}$ as a projector over the “(d−4)-dimensional” one.

All usual formulae (not involving γ_5) used for computing traces of strings of gammas in terms of combinations of the metric remain valid, even when the gammas are hatted or barred: in these cases one has only to put hats or bars over the corresponding metrics. Also the trace of an odd number of (normal, hatted or barred) γ s vanish.

Strings of gammas with contracted indices are simplified with the aid of formulae like:

$$\begin{aligned} \gamma^\mu \gamma^\nu \gamma_\mu &= (2 - d) \gamma^\nu, \\ \bar{\gamma}^\mu \bar{\gamma}^\nu \bar{\gamma}_\mu &= -2 \bar{\gamma}^\nu, \quad \bar{\gamma}^\mu \hat{\gamma}^\nu \bar{\gamma}_\mu = -4 \hat{\gamma}^\nu, \\ \hat{\gamma}^\mu \hat{\gamma}^\nu \hat{\gamma}_\mu &= (6 - d) \hat{\gamma}^\nu, \quad \hat{\gamma}^\mu \bar{\gamma}^\nu \hat{\gamma}_\mu = (4 - d) \bar{\gamma}^\nu. \end{aligned}$$

Of course, when γ_5 or the ϵ tensor appear the situation is very different. In the “naive” prescription, one assumes the *anticommutativity* of γ_5 : $\{\gamma_\mu, \gamma_5\} = 0$, and the *cyclicity* of the trace. From these assumptions one obtains [3,10,14,21]

$$\begin{aligned} 2d \text{Tr}[\gamma_5] &= 0, \\ 2(d - 2) \text{Tr}[\gamma_{\mu_1} \gamma_{\mu_2} \gamma_5] &= 0 \quad \text{if } d \neq 0, \\ 2(d - 4) \text{Tr}[\gamma_{\mu_1} \dots \gamma_{\mu_4} \gamma_5] &= 0 \quad \text{if } d \neq 0, 2; \end{aligned} \quad (6)$$

which has the consequence that $\text{Tr}[\gamma_{\mu_1} \dots \gamma_{\mu_4} \gamma_5]$ is identically 0 in the dimensionally regularized theory ($d \neq 4$). So, it also vanishes in the dimensionally renormalized theory, which

is incompatible with the true property (see eq. (4)) in four dimensions. We could introduce the symbol ϵ with the requirement that it satisfies eq. (4), but then the symbol ϵ would be identically 0. Despite this well known fact, people use the “naive” prescription in *e.g.* the Standard Model, but they do not set ϵ equal to 0 and at the end of calculations the symbol ϵ is supposed to have its usual meaning as the Levi-Civita tensor. This is clearly a mathematical inconsistency (the axioms are not compatible) and when mathematical inconsistencies are present, results obtained from the axioms are ambiguous and cannot be trusted.

Eq. (6) is obtained by computing $\text{Tr}[\gamma_{\mu_1} \dots \gamma_{\mu_m} \gamma^\alpha \gamma_\alpha \gamma_5]$ in two different ways: first, one contracts the index α without moving around the γ_5 and, second, one anticommutates the γ_α which is next to the γ_5 with this very γ_5 and puts it on the left thanks to the cyclicity of the trace; then, one anticommutates it to the right through the rest of γ s until the other γ^α is met, contraction of the index α is now carried out. So, at least the trace of six γ s, having two indices contracted, and a γ_5 are needed to obtain this inconsistency.

Here, the ambiguity in the results is a consequence of the choice of the position of the γ_5 . For example, if the γ_5 is not moved in any case,

$$\begin{aligned} \text{Tr}[\gamma^{\mu_1} \dots \gamma^{\mu_4} \gamma^\alpha \gamma_\alpha \gamma_5] &= d \text{Tr}[\gamma^{\mu_1} \dots \gamma^{\mu_4} \gamma_5], \\ &\neq -\text{Tr}[\gamma_\alpha \gamma^{\mu_1} \dots \gamma^{\mu_4} \gamma^\alpha \gamma_5] = (8 - d) \text{Tr}[\gamma^{\mu_1} \dots \gamma^{\mu_4} \gamma_5], \end{aligned} \quad (7)$$

unless the trace of four γ s and one γ_5 (or the ϵ tensor) is 0.

Notice that the ambiguity will be always proportional to $(d - 4)$ or a $(d - 4)$ object like $\hat{\gamma}_\mu$, but there are poles in $(d - 4)$ in the divergent diagrams that will make finite this ambiguity in the dimensionally renormalized theory, even in one loop calculations.

Then, why do people go on using the “naive” prescription in calculations of, *e.g.* Standard Model? The key is that the ambiguities are claimed to be always proportional to the coefficient of the (chiral gauge) anomaly. So, it would appear that one could freely use the “naive” prescription in theories with cancellation of anomalies (like the Standard Model). Calculations to low orders in perturbation theory in some models support this idea but there is not a rigorous proof of it valid to all orders in perturbation theory. (Also, it seems difficult to build a consistent theory with inconsistent elements).

We shall close this subsection with a few words regarding the construction of Lorentz covariants due to Wilson and Collins [24,10]. It is worth mentioning it since it furnishes explicit expressions for the Lorentz covariants satisfying eqs. (1)-(2), thus showing that no inconsistencies arise. In this construction gamma “matrices” are represented as infinite dimensional objects. The “matrix” γ_5 being defined by eq. (3), with $\epsilon_{\mu_1 \dots \mu_4}$ as defined by eq. (2). However, we will not use this construction but the algebraic approach of abstract symbols by Breitenlohner and Maison.

2.2. Minimal dimensional renormalization and the renormalized theory

Dimensional renormalization *à la* Breitenlohner and Maison is carried out by applying the pole subtraction algorithm [3,4,5,6] as given by the forest formula to each dimensionally regularized 1PI Feynman diagram. This algorithm can also be implemented by adding to the “ d -dimensional” Lagrangian (see below) counterterms whose coefficients have poles at

$d = 4$. Both the set of dimensionally regularized Feynman diagrams, which conforms what is referred to as the regularized theory, and the renormalized 1PI functional obtained from them are best established in the following way:

- i) Write a classical, *i.e.* order zero in $\hbar$, action, S_0 , in d space-time dimensions (d being a generic positive integer) which yield the four dimensional action of the theory when the formal limit $d \rightarrow 4$ is taken. The “ d -dimensional” covariants in S_0 are defined as in the previous subsection. We shall assume as in ref. [3] that each free term, $\int d^d x \frac{1}{2} \phi D \phi$, in S_0 is such that D^{-1} is the same algebraic expression as in four dimensions, although it is expressed in terms of “ d -dimensional” covariants. ϕ denotes a collection of bosonic or fermionic fields. $i\hbar D^{-1}$ gives the free propagator. S_0 will be referred to as the Dimensional Regularization classical action. We shall call the “ d -dimensional” space-time used above to set the Dimensional Regularization classical action, S_0 , the “ d -dimensional” space-time of Dimensional Regularization.
- ii) Use the Dimensional Regularization classical action, S_0 , along with standard path integrals textbook techniques, formally applied, to obtain the set of “ d -dimensional” Feynman rules stemming from it. These Feynman rules will lead to the collection of Feynman diagrams in the d -dimensional space-time of Dimensional Regularization, d being a generic positive integer, which is to be turned into the dimensionally regularized Feynman diagrams by using the algorithms in refs. [3,4,5,10]. This set of dimensionally regularized Feynman diagrams defines the regularized theory.
- iii) Introduce the Dimensional Regularization generating functional $Z_{\text{DReg}}[J; K; \lambda]$:

$$Z_{\text{DReg}}[J; K; \lambda] = \int \mathcal{D}\phi \exp\left\{\frac{i}{\hbar} (S_0[\phi; K; \lambda] + \int d^d x J_i(x) \phi_i(x))\right\}, \quad (8)$$

where the right hand side of eq. (8) is defined as the formal power expansion in $\hbar$, K and J given by the Feynman diagrams obtained by using the “ d -dimensional” Feynman rules previously established. The symbols K and λ denote, respectively, any external field, usually coupled linearly to composite operators, and any parameter occurring in the action. Let us stress that eq. (8) is merely a symbol which denotes the whole set of Feynman diagrams in the “ d -dimensional” space-time of Dimensional Regularization, d being a generic positive integer, which are converted into dimensionally regularized Feynman diagrams by analytic continuation in d . However, this symbol is very useful due to the Regularized Action Principle [3,4,5] discussed below. Indeed, the standard formal manipulations of the path integral, *e.g.* variations of fields and differentiation with respect to parameters, that in four dimensions lead to equations of motion, Ward identities and so on, can be formally performed in the path integral in eq. (8); thus leading to formal functional equations involving Z_{DReg} . These formal manipulations are mathematically well defined if expressed in terms of the dimensionally regularized Feynman diagrams arising from the right hand side of eq. (8), and the formal functional equations they lead to are also mathematically sound if they are considered as symbols denoting in a condensed manner the equations verified by the corresponding dimensionally regularized Feynman diagrams. These equations are the dimensionally regularized counterparts of the four dimensional equations of motion, Ward identities, *etc.* It is also useful to introduce the Dimensional Regularization 1PI functional,

$\Gamma_{\text{DReg}}[\phi; K; \lambda]$, which can be obtained from eq. (8) through formal Legendre transform; a procedure well defined in terms of regularized Feynman diagrams. Again, the Regularized Action Principle guarantees that the formal functional equations verified by $\Gamma_{\text{DReg}}[\phi; K; \lambda]$ make sense mathematically when expressed in terms of dimensionally regularized Feynman diagrams. We shall refer to the formal equations satisfied by Z_{DReg} and Γ_{DReg} as dimensionally regularized equations, keeping always in mind the previous discussion.

- iv) If there are classical symmetries in the four dimensional classical theory that should hold in the quantum theory, one generalizes next the corresponding field transformations to the d -dimensional space-time of Dimensional Regularization. Objects in this d dimensional space-time should be defined according to the algebra of “ d -dimensional” covariants given in the previous subsection. The field transformations in the d -dimensional space-time thus obtained should yield, in the limit $d \rightarrow 4$, the classical four dimensional transformations. Generally speaking, the transformations in the d dimensional space-time of Dimensional Regularization do not leave invariant the Dimensional Regularization classical action S_0 . This lack of invariance will make the Dimensional Regularization generating functionals, Z_{DReg} and Γ_{DReg} , satisfy anomalous Ward identities, which can be derived (see Appendix A) by performing formal manipulations of the path integral in eq. (8). Again, the Regularized Action Principle guarantees that both these formal manipulations and the identities they lead to have a well defined mathematical meaning when expressed in terms of dimensionally regularized Feynman diagrams. The dimensionally regularized identities so obtained are of enormous help in the computation of the anomalous breaking terms of the dimensionally renormalized four dimensional theory (see next subsection), which otherwise would have to be computed by evaluating the complete dimensionally renormalized 1PI functional.
- v) The minimal subtraction algorithm of references [3,6] is applied next to every dimensionally regularized Feynman diagram coming from the 1PI functional $\Gamma_{\text{DReg}}[\phi; K; \lambda]$. The minimally renormalized 1PI functional $\Gamma_{\text{minren}}[\phi; K; \lambda; \mu]$ (μ stands for the Dimensional Regularization scale) is obtained by taking the limit $d \rightarrow 4$, first, and then setting every hatted object to zero in every subtracted 1PI diagram. The minimal subtraction algorithm amounts to subtracting the pole at $d = 4$ from every diagram once subdivergences have been taken care of, and it can be formulated in terms of singular, at $d = 4$, counterterms [3] added to the Dimensional Regularization classical action S_0 . These singular counterterms, which can be read from the forest formula [3,10], are local polynomials of the fields and their derivatives in the d -dimensional, with $d \neq 4$, space-time of Dimensional Regularization. The coefficients of these polynomials are the principal part at $d = 4$ of a certain meromorphic function of complex d . The singular counterterms in question give rise to new “ d -dimensional” vertices, which in turn yield new Feynman diagrams that cancel, after dimensionally regularized, the singular behaviour of the dimensionally regularized diagrams provided by S_0 .

According to ref. [6] the dimensional regularization scale, μ , is introduced by replacing every loop momentum measure $\frac{d^d p}{(2\pi)^d}$ with $\mu^{4-d} \frac{d^d p}{(2\pi)^d}$ before applying the subtraction

algorithm. If one follows the procedure of singular counterterms, the previous replacement should be made regardless the diagram involves singular counterterms. The introduction of the scale μ will render dimensionally homogeneous the Laurent expansion around $d = 4$ of a given dimensionally regularized diagram.

Unfortunately, if there is a symmetry we wanted preserved at the quantum level, the renormalized functional $\Gamma_{\text{minren}}[\phi; K; \lambda; \mu]$ would not do, since it would not define, in general, a quantum theory having such a symmetry: the regularization and also the subtraction process may break the symmetry. However, the situation is not hopeless. Algebraic BRS renormalization along with the Quantum Action Principle [18] comes in our aid: if the anomaly cancellation conditions [23] are met, the anomalous breaking terms, called spurious, can be canceled at each order in $\hbar$ by adding appropriate counterterms to the four dimensional classical action of the theory.

Suppose that n is the order in the $\hbar$ expansion at which a non-anomalous symmetry is broken for the first time in the dimensionally renormalized theory. Then,

- vi)* Compute the breaking, which will be a local four dimensional functional, with the help of the action principles and the Bonneau identities [6,7,8]. This can be easily done (see next subsection) by taking as action, not the classical one, S_0 , but a new action $S^{(n)} = S_0 + S_{\text{sct}}^{(n)}$, where $S_{\text{sct}}^{(n)}$ denotes the set of singular counterterms needed to minimally renormalize the theory at order $\hbar^n$, and then redo steps *i)* to *v)*. Indeed, if $d \neq 4$, the Regularized Action Principle still holds for the Dimensional Regularization generating functionals, $Z_{\text{DReg}}[J; K; \lambda]$ and $\Gamma_{\text{DReg}}[\phi; K; \lambda]$, defined for $S^{(n)}$, and, hence, we can take advantage of the dimensionally regularized equations, in the sense explained in *ii)*, to obtain the renormalized equations verified by $\Gamma_{\text{minren}}[\phi; K; \lambda; \mu]$.

Next, extract from the breaking the four dimensional finite counterterms needed to restore the symmetry, generalize them to d space-time dimensions with the help of the algebra of d -dimensional covariants (we shall denote this generalization by $S_{\text{fct}}^{(n)}$), and add them to the action $S^{(n)}$ to obtain yet another new action $S_{\text{DReg}}^{(n)} = S_0 + S_{\text{sct}}^{(n)} + S_{\text{fct}}^{(n)}$. $S_{\text{DReg}}^{(n)}$ will be called the Dimensional Regularization action. Furnished with this new action $S_{\text{DReg}}^{(n)}$, establish then new perturbation theory: redo steps *i)* to *v)* upon replacing S_0 with $S_{\text{DReg}}^{(n)}$. The new renormalized 1PI functional, $\Gamma_{\text{ren}}[\phi; K; \lambda; \mu]$, will satisfy up to order $\hbar^n$ the Ward identities that govern the given symmetry at the quantum level. Move on to *vi)* and compute the breaking at order $\hbar^{n+1}$, and so on and so forth.

- vii)* Use the Bonneau [6,7] identities to obtain the renormalization group equation at every order in the perturbative expansion for the theory being analyzed. Thus the beta functions and anomalous dimensions of the theory will be evaluated.

Let us close this subsection by making some remarks on the generalization to d space-time dimensions, *i.e.* to the d -dimensional space-time of Dimensional Regularization, of the four dimensional classical action, the four dimensional counterterms needed to restore non-anomalous symmetries and the four dimensional symmetry transformations. If the classical action of the theory involves objects whose properties depend on the dimension of space-time (*e.g.* γ_5 , the Levi-Civita symbol,...) there is no canonical Dimensional Regularization classical action, S_0 , in d space-time dimensions. Any local functional which formally

go to the four dimensional classical action as $d \rightarrow 4$ would do, provided the free terms of the former lead to the propagators described in *i*). For instance, let κ_1 and κ_2 be a couple of positive real numbers, then, the four dimensional metric and gamma matrices in the interaction part of the four dimensional classical action can be replaced with $\bar{g}_{\mu\nu} + \kappa_1 \hat{g}_{\mu\nu}$ and $\bar{\gamma}_\mu + \kappa_2 \hat{\gamma}_\mu$, respectively, to obtain an admissible biparametric family of Dimensional Regularization classical actions. Each member of this family yields a particular regularization of the theory, this situation is somewhat reminiscent of the lattice regularization method. Generally speaking the difference between two admissible “ d -dimensional” lagrangians will always be a local “evanescent” operator[†] vanishing as the coupling constants go to zero. Analogously, in the type of theories under scrutiny, namely, chiral gauge theories, there is no canonical generalization to d -dimensional space-time of Dimensional Regularization of the four dimensional finite counterterms needed to restore a non-anomalous broken symmetry. Again, two such generalizations of a given four dimensional finite counterterm which agree up to order $\hbar^{n-1}$ will differ in a “ d -dimensional” integrated “evanescent” operator of order $\hbar^n$. However, the dependence on the choice of generalization will show in the renormalized theory at order $\hbar^{n+1}$, never at order $\hbar^n$. Last, but not least, from the point of view of renormalization, there is no canonical generalization to d space-time dimensions of the symmetry transformations of the fields for theories which are not vector-like. Again, two such generalizations differ in an “evanescent” operator. However, this arbitrariness in the choice of Dimensional Regularization classical action, “ d -dimensional” finite counterterms and symmetry transformations is useful since one can play around with it so as to simplify the form of the symmetry breaking contributions. Finally, for vector-like theories such as QCD, there is, of course, a canonical choice of Dimensional Regularization classical action and vector-like transformations: the usual one [10].

2.3. Regularized and Quantum Action Principles

As we said in the previous subsection, one of the main characteristics of the dimensional renormalization procedure is that action principles are precisely stated and proved [3-5]. These principles are most efficiently expressed in terms of the Dimensional Regularization generating functional, $Z_{\text{DReg}}[J; K; \lambda]$, for the Greens function given by the Gell-Mann-Low series and the dimensional renormalization 1PI functional, $\Gamma_{\text{ren}}[\phi; K; \lambda; \mu]$, obtained from it (see previous subsection).

Let us take as Dimensional Regularization action (see *vi*) in the previous subsection)

$$S_{\text{DReg}}^{(n)} = S_{\text{free}}[\phi; \lambda] + S_{\text{int}}[\phi; K; \lambda], \quad (9)$$

where ϕ denotes a collection of commuting or anticommuting quantum fields, K set of commuting or anti-commuting external fields, λ is a generic parameter, and

$$\begin{aligned} S_{\text{free}}[\phi; \lambda] &= S_{0\text{free}}[\phi; \lambda], \\ S_{\text{int}}[\phi; K; \lambda] &= S_{0\text{int}}[\phi; K; \lambda] + S_{\text{sct}}^{(n)}[\phi; K; \lambda] + S_{\text{fct}}^{(n)}[\phi; K; \lambda]. \end{aligned}$$

[†] An “evanescent” operator is an operator whose tree level contribution vanishes as $d \rightarrow 4$.

We next introduce

$$S_{\text{INT}}[\phi, J, K, \lambda] = S_{\text{int}}[\phi, K, \lambda] + \int d^d x J_i(x) \phi_i(x)$$

and define, following eq. (8), the functional Z_{DReg}

$$Z_{\text{DReg}}[J; K; \lambda] = \int \mathcal{D}\phi \exp\left\{\frac{i}{\hbar}(S_{\text{DReg}}^{(n)} + \int J_i \phi_i)\right\} \equiv \langle \exp\left\{\frac{i}{\hbar} S_{\text{INT}}[\phi; J; K; \lambda]\right\} \rangle_0. \quad (10)$$

The symbol $\exp$ in the previous equation stands for its formal power series and the symbol $\langle \cdots \rangle_0$ denotes the usual vacuum expectation value

$$\langle \cdots \rangle_0 = \int \mathcal{D}\phi \cdots \exp\left\{\frac{i}{\hbar} S_{\text{free}}[\phi; \lambda]\right\},$$

defined by gaussian integration, which gives a formal power series in $\hbar$ and the external fields J and K . Each coefficient of this series is a sum of “ d -dimensional” Feynman diagrams. Every “ d -dimensional” Feynman diagram can be converted into a meromorphic function of d by promoting d to a complex variable and understanding the “ d -dimensional” covariants as in subsection 2.1.

Now, the *Regularized Action Principle* states that the following three functional equations hold in the dimensionally regularized theory:

- 1) Arbitrary polynomial variations of the quantized fields ϕ , $\delta\phi(x) = \delta\theta(x) P(\phi(x))$ leave Z_{DReg} invariant and

$$\delta Z_{\text{DReg}}[J, K] \equiv \langle \delta(S_{\text{free}} + S_{\text{INT}}) \exp\left\{\frac{i}{\hbar} S_{\text{INT}}[\phi, J, K, \lambda]\right\} \rangle_0 = 0, \quad (11)$$

where the variations are the linear parts in $\delta\theta$.

- 2) variations of external fields $E(x) \equiv (K(x), J(x))$ give

$$\left\langle \frac{\delta S_{\text{INT}}}{\delta E(x)} \exp\left\{\frac{i}{\hbar} S_{\text{INT}}[\phi, J, K, \lambda]\right\} \right\rangle_0 = -i\hbar \frac{\delta Z_{\text{DReg}}[J, K, \lambda]}{\delta E(x)}. \quad (12)$$

- 3) variations of parameters give

$$\left\langle \frac{\partial(S_{\text{free}} + S_{\text{INT}})}{\partial \lambda} \exp\left\{\frac{i}{\hbar} S_{\text{INT}}[\phi, J, K, \lambda]\right\} \right\rangle_0 = -i\hbar \frac{\partial Z_{\text{DReg}}[J, K, \lambda]}{\partial \lambda}. \quad (13)$$

Let us explain in what sense are the functional equations (11), (12) and (13) mathematically meaningful. First, these equations are to be understood diagrammatically only. The formal expansion in powers of $\hbar$, K and J of both sides of each equation leads to an infinite set of equations involving only “ d -dimensional” Feynman diagrams. It is this infinite set of diagrammatic equations in the d -dimensional space-time of Dimensional Regularization what is taken as the definition of the corresponding functional equation. The *Regularized Action Principle* [3,6] states that this very set of diagrammatic equations still

holds if every formal “ d -dimensional” Feynman diagram in them is replaced with its dimensionally regularized counterpart, with the scale μ introduced as explained in 2.2.v).[†] A mathematically sound set of equations is thus obtained. Of course, it is easier to handle the functional equations above than their dimensionally regularized diagrammatic definition. Furthermore, this functional equations can be deduced by formal path integral manipulations, which in turn can be given a diagrammatic definition valid within the Dimensional Regularization method *à la* Breitenlohner and Maison. Let us finally note that the limit $d \rightarrow 4$ of dimensionally regularized the Green functions coming from the right hand side of eqs. (12) and (13) does exist at each order in the $\hbar$ expansion up to order n . Notice that we have included in the action $S_{\text{DReg}}^{(n)}$, defined in eq. (9), the appropriate singular counterterms up to order $\hbar^n$. Hence, eqs. (12) and (13) have a well-defined $d \rightarrow 4$ limit up to order $\hbar^n$ and can be used to compute the variations with $E(x)$ and λ of the renormalized functional $Z_{\text{ren}}[J; K; \lambda]$ up to order $\hbar^n$.

The *Regularized Action Principle* given by eqs. (11), (12) and (13) leads [3,6,7,10] to the *Quantum Action Principle* (see ref. [25] and references therein) which states that the following functional equations hold for the dimensionally renormalized theory:

QAP1)

$$\frac{\partial \Gamma_{\text{ren}}}{\partial \lambda} = N\left[\frac{\partial(S_0 + S_{\text{fct}}^{(n)})}{\partial \lambda}\right] \cdot \Gamma_{\text{ren}}. \quad (14)$$

QAP2)

$$\frac{\delta \Gamma_{\text{ren}}}{\delta E(x)} = N\left[\frac{\delta(S_0 + S_{\text{fct}}^{(n)})}{\delta E(x)}\right] \cdot \Gamma_{\text{ren}}. \quad (15)$$

QAP3) Let $\phi'(x) = P(\phi(x))$ denote a linear transformation of $\phi(x)$ whose coefficients do not depend explicitly on d , then

$$\phi'(x) \frac{\delta \Gamma_{\text{ren}}}{\delta \phi(x)} = N\left[\phi'(x) \frac{\delta(S_0 + S_{\text{fct}}^{(n)})}{\delta \phi(x)}\right] \cdot \Gamma_{\text{ren}}. \quad (16)$$

QAP4) If $\delta\phi(x)$ is a non-linear field transformation in the four dimensional space-time and $K(x)$ is the external field coupled to it, then

$$\frac{\delta \Gamma_{\text{ren}}}{\delta K(x)} \frac{\delta \Gamma_{\text{ren}}}{\delta \phi(x)} = N[\mathcal{O}(x)] \cdot \Gamma_{\text{ren}}, \quad (17)$$

where the symbols $N[\dots]$ and $N[\dots] \cdot \Gamma_{\text{ren}}$ denote normal product as defined in refs. [6,7,10] and its insertion in the renormalized 1PI functional, respectively. $N[\mathcal{O}(x)]$ is a local normal product of ultraviolet dimension $4 - \dim(\phi) + \dim(\delta\phi)$. S_0 and $S_{\text{fct}}^{(n)}$ have been defined in *i)* and *v)* of subsection 2.2.

Eqs. (14)-(17) have been shown to hold by using the forest formula [3,4,6,7] and the singular counterterms algorithm [10], respectively. Of course, both proofs are equivalent.

[†] The scale μ is introduced so as to render dimensionally homogeneous the Laurent expansions around $d = 4$.

The *Quantum Action Principle* is of the utmost importance to the renormalization of BRS symmetries [18], for it guarantees (see QAP1 and QAP2) that the breaking of such symmetry is given by the insertion into the dimensionally renormalized 1PI Γ_{ren} of a certain integrated normal operator, $N[\mathcal{O}] = \int d^4x N[\mathcal{O}(x)]$, where $N[\mathcal{O}(x)]$ is a local normal operator of ultraviolet dimension $4 - \dim(\Phi) + \dim(s\Phi)$ and ghost number 1. Indeed, the following equation holds for $\Gamma_{\text{ren}}[\varphi, \Phi; K_\Phi]$

$$\mathcal{S}(\Gamma_{\text{ren}}) \equiv \int d^4x (s\varphi) \frac{\delta\Gamma_{\text{ren}}}{\delta\varphi} + \frac{\delta\Gamma_{\text{ren}}}{\delta K_\Phi} \frac{\delta\Gamma_{\text{ren}}}{\delta\Phi} = \Delta_{\text{breaking}}. \quad (18)$$

with $\Delta_{\text{breaking}} = N[\mathcal{O}] \cdot \Gamma_{\text{ren}}$. In the previous equation φ and Φ stand for fields which undergo linear and non-linear BRS transformations, respectively. As it is customary $s\varphi$ denotes a BRS transformation. K_Φ stands for the external field coupled to the non-linear BRS transformation $s\Phi$ in the four-dimensional classical action [18].

Now, if finite counterterms have been added to the four-dimensional classical action so that the BRS symmetry is preserved up to order $\hbar^{n-1}$ (*i.e.* the first non-vanishing contribution to the right hand side of eq. (18) is order $\hbar^n$), eq. (18) reads

$$\mathcal{S}(\Gamma_{\text{ren}}) = N[\mathcal{O}]^{(n)} + O(\hbar^{n+1}).$$

Where $N[\mathcal{O}]^{(n)}$ is the contribution to $N[\mathcal{O}]$ of order $\hbar^n$. We have taken into account that $N[\mathcal{O}] \cdot \Gamma_{\text{ren}} = N[\mathcal{O}]^{(n)} + O(\hbar N[\mathcal{O}]^{(n)})$, since by assumption $N[\mathcal{O}] \cdot \Gamma_{\text{ren}} = O(\hbar^n)$.

Next, if $N[\mathcal{O}]^{(n)} = -\hbar^n b S_{\text{fct},n}$ (*i.e.* $N[\mathcal{O}]^{(n)}$ is b -exact) for some four-dimensional integrated local functional of the fields, the BRS symmetry can be restored, up to order $\hbar^n$, by adding to the four-dimensional classical action the finite counterterm $S_{\text{fct},n}$ [18]. Here b denotes the linearized BRS operator:

$$b = s + \int d^4x \left\{ \frac{\delta\Gamma_{\text{ren}}^{(0)}}{\delta\Phi(x)} \frac{\delta}{\delta K_\Phi(x)} \right\},$$

and $\Gamma_{\text{ren}}^{(0)}$ is the order $\hbar^0$ contribution to Γ_{ren} , *i.e.* the BRS invariant classical four dimensional action. Then, we set $S_{\text{fct}}^{(n)} = \sum_{m=1}^n \hbar^m S_{\text{fct},m}$. Of course, if $N[\mathcal{O}]^{(n)}$ is not b -exact the theory is anomalous at order $\hbar^n$.

The computation of the right hand side of eq. (18), Δ_{breaking} , is the issue we will discuss now. One can always compute the complete 1PI functional Γ_{ren} at each order in $\hbar$, insert it in the left hand side of equation (18), work out the functional derivatives and thus obtain the right hand side. This method is, of course, very impractical. The *Regularized Action principle*, as given in eqs. (11) and (12), provides us with a more efficient way to compute the symmetry breaking term of the BRS symmetry. Indeed, in Appendix A we show that the following functional equation hold, in the same sense as eqs. (11) and (12), for the 1PI formal functional Γ_{DReg}

$$\begin{aligned} \mathcal{S}_d(\Gamma_{\text{DReg}}) &\equiv \int d^d x (s_d \varphi) \frac{\delta\Gamma_{\text{DReg}}}{\delta\varphi} + \frac{\delta\Gamma_{\text{DReg}}}{\delta K_\Phi} \frac{\delta\Gamma_{\text{DReg}}}{\delta\Phi} = \\ &= \Delta \cdot \Gamma_{\text{DReg}} + \Delta_{\text{ct}} \cdot \Gamma_{\text{DReg}} + \int d^d x \left[\frac{\delta S_{\text{ct}}^{(n)}}{\delta K_\Phi(x)} \cdot \Gamma_{\text{DReg}} \right] \frac{\delta\Gamma_{\text{DReg}}}{\delta\Phi(x)}, \end{aligned} \quad (19)$$

were Γ_{DReg} is the 1PI functional computed with the Dimensional Regularization action, $S_{\text{DReg}}^{(n)}$, defined at the beginning of the current subsection. The symbol s_d denotes the generalization to d dimensions of the four-dimensional BRS transformations (see 2.2.iv)). The operators Δ and Δ_{ct} are given by $\Delta = s_d S_0$ and $\Delta_{\text{ct}} = s_d S_{\text{ct}}^{(n)}$, respectively. Here, $S_0 = S_{0\text{free}} + S_{0\text{int}}$ and $S_{\text{ct}}^{(n)} = S_{\text{fct}}^{(n)} + S_{\text{sct}}^{(n)}$ (see eq. (9) and the paragraph below it for definitions).

Let us recall that Γ_{DReg} is defined as a formal series expansion in powers of $\hbar$ and the fields φ , Φ and K_ϕ . The coefficients of this power series are 1PI Green functions of the theory at a given order in $\hbar$. Taking into account that Γ_{DReg} contains the contributions coming from the singular counterterms up to order $\hbar^n$, denoted by $S_{\text{sct}}^{(n)}$, one concludes that the dimensionally regularized 1PI Green functions obtained from Γ_{DReg} are finite in the limit $d \rightarrow 4$ up to order $\hbar^n$. These 1PI Green functions are the dimensionally renormalized 1PI Green functions upon introduction of the Dimensional Regularization scale μ (see subsection 2.2.). Hence, up to order $\hbar^n$, one may formally write

$$\text{LIM}_{d \rightarrow 4} \Gamma_{\text{DReg}}[\varphi, \Phi; K_\Phi; \mu] = \Gamma_{\text{ren}}[\varphi, \Phi; K_\Phi; \mu] \quad (20)$$

In the previous equation, $\text{LIM}_{d \rightarrow 4}$ is defined by the following process: one takes first the limit $d \rightarrow 4$ of the dimensionally regularized 1PI Green functions and then sets to zero every hatted object (*i.e.* $(d-4)$ -dimensional covariant). Recall that the Dimensional Regularization scale, μ , is introduced by hand to render dimensionally homogeneous the regularized Green functions upon Laurent expansion around $d = 4$ [6].

Now, by taking $\text{LIM}_{d \rightarrow 4}$ of the left hand side of eq. (19) after discarding any contribution of order greater than $\hbar^n$, and bearing in mind eq. (20), one obtains the left hand side of eq. (18) up to order $\hbar^n$. Hence, Δ_{breaking} in the latter equation is given by

$$\Delta_{\text{breaking}} = \text{LIM}_{d \rightarrow 4} \left\{ \Delta \cdot \Gamma_{\text{DReg}} + \Delta_{\text{ct}} \cdot \Gamma_{\text{DReg}} + \int d^d x \left[\frac{\delta S_{\text{ct}}^{(n)}}{\delta K_\Phi(x)} \cdot \Gamma_{\text{DReg}} \right] \frac{\delta \Gamma_{\text{DReg}}}{\delta \Phi(x)} \right\}. \quad (21)$$

Of course, the previous equation only makes sense up to order $\hbar^n$. The order $\hbar$ contribution to eq. (21) reads particularly simple since $s_d S_0$ is an evanescent operator (there is no $O(\hbar^0)$ contribution to $N[\Delta]$ thereby) and the lowest order contribution to S_{ct} is order $\hbar$:

$$\Delta_{\text{breaking}}^{(1)} = \text{LIM}_{d \rightarrow 4} \left\{ \left[\Delta \cdot \Gamma_{\text{DReg}} \right]_{\text{singular}}^{(1)} + b_d S_{\text{sct}}^{(1)} \right\} + \left[N[\Delta] \cdot \Gamma_{\text{ren}} \right]^{(1)} + b S_{\text{fct}}^{(1)}, \quad (22)$$

where the superscript (1) denotes contributions of order $\hbar$ and Δ is equal to the evanescent operator $s_d S_0$. In eq. (22), b_d stands for the linearized BRS operator in d dimensions:

$$b_d = s_d + \int d^d x \left\{ \frac{\delta S_0}{\delta \Phi(x)} \frac{\delta}{\delta K_\Phi(x)} \right\}.$$

The symbol $\left[\Delta \cdot \Gamma_{\text{DReg}} \right]_{\text{singular}}^{(1)}$ denotes the expression, singular at $d = 4$, that it is subtracted from $\Delta \cdot \Gamma_{\text{DReg}}$ to obtain $\left[N[\Delta] \cdot \Gamma_{\text{ren}} \right]^{(1)}$ according to the forest formula [3,6]. Hence,

$$\left[\Delta \cdot \Gamma_{\text{DReg}} \right]^{(1)} = \left[\Delta \cdot \Gamma_{\text{DReg}} \right]_{\text{singular}}^{(1)} + \left[N[\Delta] \cdot \Gamma_{\text{ren}} \right]^{(1)} + \text{vanishing terms}. \quad (23)$$

By “vanishing terms” we mean contributions that vanish as d is sent to 4 first and then every hatted object is set to zero.

There are some comments regarding eq. (22) which we would like to make. First, if the theory is not anomalous, the finite counterterms in $S_{\text{fct}}^{(1)}$ can be chosen so that $\Delta_{\text{breaking}}^{(1)} = 0$ [18]. Second, there are two possible sources of symmetry breaking in eq. (22). One is the regularization method; in particular, the Dimensional Regularization classical action, S_0 , one began with. The other is the minimal subtraction algorithm (*i.e.* the renormalization algorithm) we are employing. That the regularization method breaks the symmetry shows itself in the contributions $[\Delta \cdot \Gamma_{\text{DReg}}]_{\text{singular}}^{(1)}$ and $[N[\Delta] \cdot \Gamma_{\text{ren}}]^{(1)}$. The subtraction algorithm gives rise to the contribution $b_d S_{\text{sct}}^{(1)}$, which need not be zero as we shall show later on. Third, the “limit”

$$\text{LIM}_{d \rightarrow 4} \left\{ [\Delta \cdot \Gamma_{\text{DReg}}]_{\text{singular}}^{(1)} + b_d S_{\text{sct}}^{(1)} \right\}$$

need not be zero either, as we shall discuss in section 3. Of course, this “limit” yields a finite local four-dimensional functional of the fields and their derivatives. Fourth, the operator $\Delta = s_d S_0$ being evanescent ushers in the techniques introduced by Bonneau [6,7] to express normal products of evanescent operators, also called *anomalous* normal products, in terms of a basis of standard (non-evanescent operators) normal products. We shall address this subject in the next subsection. Suffice it to say that expansion of the normal product of an evanescent operator in terms of “standard” normal products has coefficients that are series expansions in $\hbar$ with no $O(\hbar)$ term. Hence, there is no breaking of the BRS symmetry at order $\hbar^0$ and $[N[\Delta] \cdot \Gamma_{\text{ren}}]^{(1)}$ is, in agreement with the *Quantum Action Principle*, a sum of standard local operators.

We shall finally recall that the equation of motion holds in dimensional renormalization [3,6,10]:

$$\frac{\delta \Gamma_{\text{ren}}}{\delta \phi(x)} = N \left[\frac{\delta (S_0 + S_{\text{fct}}^{(n)})}{\delta \phi(x)} \right] \cdot \Gamma_{\text{ren}} = \text{LIM}_{d \rightarrow 4} \left\{ \frac{\delta S_{\text{DReg}}^{(\infty, n)}}{\delta \phi(x)} \cdot \Gamma_{\text{DReg}} \right\}, \quad (24)$$

where $S_{\text{fct}}^{(n)}$ and $S_{\text{DReg}}^{(\infty, n)}$ are defined in subsection 2.2. and eq. (9). The superscript ∞ tell us that the singular counterterms needed to minimally renormalize the theory at any order in $\hbar$ have been included; however, one has added finite counterterms only up to order $\hbar^n$. Eq. (24) can be obtained from QAP3 (eq. (16)) by setting $\phi'(x)$ to 1. Alternatively, eq. (11), for $P(\phi(x)) = 1$, leads to eq. (24).

The use of both the equation of motion of the ghost field and the equation of motion of the auxiliary field coupled to the gauge fixing condition simplifies the renormalization of theories with BRS symmetries.

2.4. Expression of anomalous terms with the aid of Bonneau identities

The computation of the insertions of normal products of evanescent operators, also called anomalous normal products, are, in dimensional renormalization, of key importance to the calculation of the symmetry breaking terms in Ward identities [6,10]. We shall see in the next section that $[N[\Delta] \cdot \Gamma_{\text{ren}}]^{(1)}$ (see eq. (22)) yields both the essential and the

spurious BRS anomalies for the regularized theory we will consider. The purpose of the current subsection is to summarily recall that the normal product of an evanescent operator can be expressed as linear combinations of normal products of standard (non-evanescent) operators. Further details can be found in refs. [6,7,8].

In renormalized perturbation theory, any set of insertions of quantum operators such that their classical approximations form a basis in the linear space of classical operators of dimensions bounded by D is also a basis in the linear space of insertions, with dimension bounded by the same D , of quantum operators [18]. So, in dimensional renormalization, if $\{\mathcal{O}^i\}$ form such a basis in the space of classical operators, then $\{N[\mathcal{O}^i] \cdot \Gamma_{\text{ren}}\}$, where $\mathcal{O}^i$ is any of the possible generalizations to d dimensions of the corresponding classical four-dimensional operator, is a basis in the space of quantum insertions. We will call standard operators the operators in the “ d -dimensional” space-time of Dimensional Regularization which are generalizations of the classical operators in four dimensions. These standard operators are obtained with the help of the algebra “ d -dimensional” covariants and they are non-evanescent. Hence, $N[\mathcal{O}^i] \cdot \Gamma_{\text{ren}}$ is an insertion of an standard normal product.

It is thus expected that anomalous normal products, *i.e.* normal products of non-evanescent operators minimally subtracted should be decomposable in terms of some basis of standard normal products, minimally subtracted again. This was shown by Bonneau [8] who, remarkably, proved a linear system of Zimmermann-like identities which expresses a decomposition of any anomalous normal product on all possible normal products of monomials of operators, including standard and anomalous ones.

If there is only a scalar field, the Bonneau identities have the form

$$\begin{aligned}
N[\hat{g}_{\mu\rho}\mathcal{O}^{\mu\rho}](x) \cdot \Gamma_{\text{ren}} = & - \sum_n^4 \sum_r^{4-n} \sum_{\substack{\{i_1 \dots i_r\} \\ 1 \leq i_j \leq n}} \\
& \left\{ \text{r.s.p.} \frac{(-i)^r}{r!} \frac{\partial^r}{\partial p_{i_1}^{\mu_1} \dots \partial p_{i_r}^{\mu_r}} \overline{\langle \tilde{\phi}(p_1) \dots \tilde{\phi}(p_n) N[\check{g}_{\mu\rho}\mathcal{O}^{\mu\rho}] \rangle^{\text{1PI}}} \Big|_{p_i \equiv \check{g} \equiv 0} \right\} \\
& \times N\left[1/n! \prod_{k=1}^n \left\{ \left(\prod_{\{\alpha/i_\alpha=k\}} \partial_{\mu_\alpha} \right) \phi \right\} \right](x) \cdot \Gamma_{\text{ren}}, \tag{25}
\end{aligned}$$

where the tilde indicates Fourier transformed fields, the bar means that only the minimal subtraction of the subdivergences has been done, r.s.p. stands for “residue of the simple pole in $\varepsilon = 4 - d$ ” (this is the reason why the global sign -1 occurs: $\hat{g}^\mu{}_\mu = -\varepsilon$) and the $\times$ symbol means that the tensorial structure $\{\dots\}$ really appears *inside* the normal product. Of course, any colour or pure number factor can be taken out of the normal product, but note that $g_{\mu\nu}N[\mathcal{O}^{\mu\nu}] = N[\bar{g}_{\mu\nu}\mathcal{O}^{\mu\nu}] \neq N[g_{\mu\nu}\mathcal{O}^{\mu\nu}]$, the difference being the anomalous normal product $N[\hat{g}_{\mu\nu}\mathcal{O}^{\mu\nu}]$, which need not vanish.

Notice that the sum in n is a sum in the number of fields in the monomial and the sum in r , a sum in the number of derivatives in the monomial.

The tensor $\check{g}$ is a new one, which has been introduced in order to simplify the calculations. Its properties are:

$$\check{g}_{\mu\nu} = \check{g}_{\nu\mu}, \quad \check{g}_\mu{}^\mu = 1,$$

$$\begin{aligned}\check{g}_{\mu\rho}g^\rho{}_\nu &= \check{g}_{\mu\rho}\hat{g}^\rho{}_\nu = \check{g}_{\mu\nu} \\ \check{g}_{\mu\rho}\bar{g}^{\rho\mu} &= 0, \quad \text{N}[\check{g}_{\mu\nu}\mathcal{O}^{\mu\nu}] = \check{g}_{\mu\nu}\text{N}[\mathcal{O}^{\mu\nu}].\end{aligned}$$

Eq. (25) is easily generalized when there are several fields, not necessarily scalar, involved. One just sums over all kind of fields, taking special care of the symmetry factors and the fermionic signs. We shall give below the generalization of eq. (25) that is relevant to our computations in subsection 3.5.

Due to the fact that all subdivergences have been previously subtracted, the barred 1PI function of eq. (25) has a polynomial singular part. We are interested in the coefficients of the simple pole. These coefficients will be in general combinations of metrics, constants and colour factors. In vector-like gauge theories with the usual regularizations, its tensorial structure will involve only usual metrics, which combined with the tensor indices of the normal product will get the expansions of the anomalous normal product in terms of a basis of standard monomial normal products. However, if in the regularized theory or in the calculations some hatted objects appear, then the Bonneau identities (eq. (25)) would express any anomalous normal product in terms of a collection of standard ($\mathcal{M}^i$) and also evanescent ($\hat{\mathcal{M}}^j$) monomial normal products. That is

$$\text{N}[\hat{g}_{\mu\rho}\mathcal{O}^{\mu\rho}](x) \cdot \Gamma_{\text{ren}} = \sum_i \alpha_i \text{N}[\mathcal{M}^i](x) \cdot \Gamma_{\text{ren}} + \sum_j \hat{\alpha}_j \text{N}[\hat{\mathcal{M}}^j](x) \cdot \Gamma_{\text{ren}}.$$

Therefore, the rest of independent anomalous monomials should be also expanded by a similar formula, then getting a system of identities.

But notice that the left hand side of the Bonneau identities and also the coefficients given by the r.s.p. of the right hand side are both of order $\hbar^1$, in the least. Therefore, the Bonneau identities are not a trivial system but a linear system whose unique solution give the desired expansion of any anomalous operator in terms of a quantum basis of standard insertions:

$$\text{N}[\hat{g}_{\mu\rho}\mathcal{O}^{\mu\rho}](x) \cdot \Gamma_{\text{ren}} = \sum_i q_i \text{N}[\mathcal{M}^i](x) \cdot \Gamma_{\text{ren}}.$$

Of course, the coefficients q_i are formal series in $\hbar$.

Clearly, at lowest order the linear system is decoupled and has an easy interpretation: the loops with the anomalous insertion are replaced with sums of tree diagrams corresponding to insertion of standard operators, but with coefficients of order $\hbar^1$. This will be most relevant to our computations below. In general, for the calculation of q_i at order $\hbar^m$, it is needed the coefficients α_i up to order $\hbar^m$ and the coefficients $\hat{\alpha}_i$ up to order $\hbar^{(m-1)}$.

3. Chiral non-Abelian Yang-Mills theories: simple gauge groups

This section is devoted to the study of the one-loop dimensional renormalization of a general Chiral non-Abelian gauge theory with neither Majorana nor scalar fields. We shall assume for the time being that the gauge group is a compact simple Lie group and leave for section 4. the case of a general compact group.

Let us give first some definitions and display some properties

$$P_L = \frac{1 - \gamma_5}{2}, \quad P_R = \frac{1 + \gamma_5}{2}, \quad P_L^2 = P_L, \quad P_R^2 = P_R, \quad P_L P_R = P_R P_L = 0$$

$$\psi_L = P_L \psi, \quad \psi_R = P_R \psi, \quad \bar{\psi}_L = \bar{\psi} P_R, \quad \bar{\psi}_R = \bar{\psi} P_L,$$

and also the special properties of the symbols in the algebra of covariants:

$$\begin{aligned} P_R \gamma^\mu P_R &= P_R \hat{\gamma}^\mu P_R = \hat{\gamma}^\mu P_R \neq 0, \\ P_L \gamma^\mu P_L &= P_L \hat{\gamma}^\mu P_L = \hat{\gamma}^\mu P_L \neq 0, \\ P_L \gamma^\mu P_R &= P_L \bar{\gamma}^\mu P_R = \bar{\gamma}^\mu P_R = P_L \bar{\gamma}^\mu, \\ P_R \gamma^\mu P_L &= P_R \bar{\gamma}^\mu P_L = \bar{\gamma}^\mu P_L = P_R \bar{\gamma}^\mu, \\ P_L \bar{\gamma}^\mu P_L &= P_R \bar{\gamma}^\mu P_R = 0 = P_L \hat{\gamma}^\mu P_R = P_R \hat{\gamma}^\mu P_L. \end{aligned} \tag{26}$$

3.1. The Classical Action

The BRS invariant classical four dimensional action is

$$S_{\text{cl}} = S_{\text{inv}} + S_{\text{gf}} + S_{\text{ext}}, \tag{27}$$

with

$$\begin{aligned} S_{\text{inv}} &= \int d^4x \left[-\frac{1}{4g^2} \text{Tr} F_{\mu\nu} F^{\mu\nu} + \frac{i}{2} \bar{\psi} \overleftrightarrow{D}_L \psi + \frac{i}{2} \bar{\psi}' \overleftrightarrow{D}_R \psi' \right], \\ S_{\text{gf}} &= \int d^4x \left[\text{Tr} \frac{\alpha}{2} B^2 + \text{Tr} B(\partial_\mu A^\mu) - \text{Tr} \bar{\omega} \partial^\mu \nabla_\mu \omega \right], \\ S_{\text{ext}} &= \int d^4x \left[\text{Tr} \rho^\mu s A_\mu + \text{Tr} \zeta s \omega + \bar{L} s \psi + L s \bar{\psi} + \bar{R} s \psi' + R s \bar{\psi}' \right], \end{aligned}$$

where $A_\mu = A_\mu^a \tau^a$, $\omega, \bar{\omega}, B, \rho$ and ζ take values on the Lie algebra of a compact simple Lie group G . τ^a are the generators of G in a given finite dimensional representation, normalized so that $\text{Tr} [\tau^a \tau^b] = \delta^{ab}$. We thus have $[\tau^a, \tau^b] = i c^{abc} \tau^c$, c^{abc} being completely antisymmetric, which defines the adjoint representation $(T_A^a)_{ij} = -i c^{aij}$ with a certain normalization $\text{Tr} [T_A^a T_A^b] = T_A \delta^{ab}$, $(T_A^e T_A^e)_{ij} = C_A \delta_{ij}$, $T_A = C_A$. The following definitions will be used in the sequel

$$\begin{aligned} F_{\mu\nu} &= F_{\mu\nu}^a \tau^a = \partial_\mu A_\nu - \partial_\nu A_\mu - i[A_\mu, A_\nu], \\ \nabla_\mu \phi^a &= \partial_\mu \phi^a + c^{abc} A_\mu^b \phi^c, \quad \phi \text{ being a Lie algebra valued object} \\ D_{L\mu} \psi &= (\partial_\mu - i A_\mu^a T_L^a P_L) \psi, \\ D_{R\mu} \psi' &= (\partial_\mu - i A_\mu^a T_R^a P_R) \psi'. \end{aligned}$$

ψ (ψ') represents a collection of left-handed (right-handed) fermionic multiplets carrying finite representations, T_L^a (T_R^a) of the group generators. The following equations hold

$$\begin{aligned} [T_L^a, T_L^b] &= ic^{abc} T_L^c, & [T_R^a, T_R^b] &= ic^{abc} T_R^c, \\ \text{Tr} [T_L^a T_L^b] &= T_L \delta^{ab}, & \text{Tr} [T_R^a T_R^b] &= T_R \delta^{ab}, \\ T_L^e T_L^e &= C_L, & T_R^e T_R^e &= C_R. \end{aligned}$$

We also introduce the shorthand notation:

$$\begin{aligned} \text{Tr} [T_L^{a_1} \dots T_L^{a_n}] &\equiv T_L^{a_1 \dots a_n}, & \text{Tr} [T_R^{a_1} \dots T_R^{a_n}] &\equiv T_R^{a_1 \dots a_n}, \\ T_L^{a_1 \dots a_n} + T_R^{a_1 \dots a_n} &\equiv T_{L+R}^{a_1 \dots a_n}, & T_L^{a_1 \dots a_n} - T_R^{a_1 \dots a_n} &\equiv T_{L-R}^{a_1 \dots a_n}, \\ \text{Tr}_L [\phi_1 \dots \phi_n] &\equiv \phi_1^{a_1} \dots \phi_n^{a_n} \text{Tr} [T_L^{a_1} \dots T_L^{a_n}], & \text{Tr}_R [\phi_1 \dots \phi_n] &\equiv \phi_1^{a_1} \dots \phi_n^{a_n} \text{Tr} [T_R^{a_1} \dots T_R^{a_n}]. \end{aligned}$$

The appropriate index labeling different fermions will be understood throughout this paper: every left handed multiplet can yield a different, say, T_L , C_L ...

For the action we have chosen, the free boson propagator is (in momentum space):

$$g^2 \frac{-i}{k^2 + i\epsilon} \delta^{ab} \left[g^{\mu\nu} - \left(1 - \frac{\alpha}{g^2}\right) \frac{k^\mu k^\nu}{k^2} \right] = g^2 \frac{-i}{k^2 + i\epsilon} \delta^{ab} \left[\left(g^{\mu\nu} - \frac{k^\mu k^\nu}{k^2}\right) + \frac{\alpha}{g^2} \frac{k^\mu k^\nu}{k^2} \right].$$

Because in perturbative calculations the combination α/g^2 will often appear, we denote it with α' . Therefore, the Feynman gauge here is defined by $\alpha = g^2$ or $\alpha' = 1$ †.

S_{cl} is left invariant in four dimensions by the BRS transformations:

$$\begin{aligned} s\psi &= i\omega^a T_L^a P_L \psi, & s\bar{\psi} &= i\bar{\psi} T_L^a P_R \omega^a, \\ s\psi' &= i\omega^a T_R^a P_R \psi', & s\bar{\psi}' &= i\bar{\psi}' T_R^a P_L \omega^a, \\ sA_\mu &= \nabla_\mu \omega, & s\omega &= i\omega^2, & s\bar{\omega} &= B, & sB &= 0, \\ s\rho^\mu &= 0, & s\zeta &= 0, & sL &= s\bar{L} = sR = s\bar{R} = 0. \end{aligned} \tag{28}$$

This is due to the anticommutativity of γ_5 in four dimensions, which allow us to write S_{inv} in the gauge invariant form

$$S_{\text{inv}} = \int d^4x \left[-\frac{1}{4g^2} F_{\mu\nu} F^{\mu\nu} + \frac{i}{2} \bar{\psi}_L \overleftrightarrow{D}_L \psi_L + \frac{i}{2} \bar{\psi}'_R \overleftrightarrow{D}_R \psi'_R + \frac{i}{2} \bar{\psi}_R \overleftrightarrow{\partial} \psi_R + \frac{i}{2} \bar{\psi}'_L \overleftrightarrow{\partial} \psi'_L \right].$$

The action preserves the ghost number. Its value together with the dimensions and the commutativity for the different fields are shown in Table 1.

The four-dimensional BRS linear operator b is:

† If the fields are rescaled by g : $A \rightarrow gA'$, $\omega \rightarrow g\omega'$, $\bar{\omega} \rightarrow g^{-1}\bar{\omega}'$; and $\alpha \rightarrow g^2\alpha'$, then the action adopt the other usual form with g being interpreted as the “coupling” constant. But let us remember that the loop expansion is an expansion in $\hbar$ rather than in g .

	s	A_μ	ψ, ψ'	ω	$\bar{\omega}$	B	ρ_μ	ζ	L, R
Ghost numb.	1	0	0	1	-1	0	-1	-2	-1
Dimension	0	1	3/2	0	2	2	3	4	5/2
Commutat.	-1	+1	-1	-1	-1	+1	-1	+1	+1

Table 1: Ghost number and dimension of the fields. In the last row, +1 (-1) means that the symbol commutes (anticommutes)

$$b \equiv s + \int d^4x \left\{ \text{Tr} \frac{\delta S_{\text{cl}}}{\delta A_\mu} \frac{\delta}{\delta \rho^\mu} + \text{Tr} \frac{\delta S_{\text{cl}}}{\delta \omega} \frac{\delta}{\delta \zeta} + \frac{\delta S_{\text{cl}}}{\delta \psi} \frac{\delta}{\delta \bar{L}} + \frac{\delta S_{\text{cl}}}{\delta \psi'} \frac{\delta}{\delta \bar{R}} + \frac{\delta S_{\text{cl}}}{\delta \bar{\psi}} \frac{\delta}{\delta L} + \frac{\delta S_{\text{cl}}}{\delta \bar{\psi}'} \frac{\delta}{\delta R} \right\}, \quad (29)$$

which satisfies $b^2 = 0$ because is the linearization of the BRS operator and the classical action satisfy the BRS identities [18]. The symbol s has been defined in eq. (28).

The classical b -invariant combinations are:

$$\begin{aligned} L_g &= g \frac{\partial S_{\text{cl}}}{\partial g} = \frac{1}{2g^2} \int d^4x \text{Tr} F_{\mu\nu} F^{\mu\nu}, \\ L_A &= b \cdot \int d^4x \tilde{\rho}_\mu^a A^{a\mu}, \quad L_\omega = -b \cdot \int d^4x \zeta^a \omega^a, \\ L_\psi^L &= -b \cdot \int d^4x \bar{L} P_L \psi + \bar{\psi} P_R L = 2 \int d^4x \frac{i}{2} \bar{\psi} \overleftrightarrow{\not{\partial}} P_L \psi + \bar{\psi} \gamma^\mu P_L T_L^a \psi A_\mu^a, \\ L_{\psi'}^R &= -b \cdot \int d^4x \bar{R} P_R \psi' + \bar{\psi}' P_L R = 2 \int d^4x \frac{i}{2} \bar{\psi}' \overleftrightarrow{\not{\partial}} P_R \psi' + \bar{\psi}' \gamma^\mu P_R T_R^a \psi' A_\mu^a, \\ L_\psi^R &= -b \cdot \int d^4x \bar{L} P_R \psi + \bar{\psi} P_L L = 2 \int d^4x \frac{i}{2} \bar{\psi} \overleftrightarrow{\not{\partial}} P_R \psi, \\ L_{\psi'}^L &= -b \cdot \int d^4x \bar{R} P_L \psi' + \bar{\psi}' P_R R = 2 \int d^4x \frac{i}{2} \bar{\psi}' \overleftrightarrow{\not{\partial}} P_L \psi'; \end{aligned} \quad (30)$$

where $\tilde{\rho}^\mu = \rho^\mu + \partial^\mu \bar{\omega}$. L_ψ^R and $L_{\psi'}^L$ will prove not to be useful due to our choice of the regularized chiral vertex.

It will be shown later that the non-trivial L_g is associated with the finite renormalizations of the coupling constant g , whereas the b -exact terms, L_ψ , $L_{\psi'}$, L_A and L_ω give rise to finite renormalizations of the corresponding wave functions in the space of field functionals in four dimensional space-time.

3.2. The Dimensional Regularization Action

We shall now follow *i)* of subsection 2.2. and set the Dimensional Regularization classical action S_0 . The kinetic terms are uniquely defined, not so the interaction terms. Notice, for instance, that the Dirac matrix part of the left-handed-fermion-gauge-boson vertex has the following equivalent forms in 4 dimensions: $\gamma^\mu P_L = P_R \gamma^\mu = P_R \gamma^\mu P_L$. All these forms are different in the d -dimensional space-time of Dimensional Regularization because of the non-anticommutativity of γ_5 . Of course, the generalization of the interaction to the Dimensional Regularization space is not unique, and any choice is *equally correct*, although some choices will be more convenient than others.

We choose the following generalization of S_{inv} :

$$\begin{aligned} \int d^d x & - \frac{1}{4g^2} \text{Tr} F_{\mu\nu} F^{\mu\nu} + \frac{i}{2} \bar{\psi} \overleftrightarrow{\partial} \psi + \frac{i}{2} \bar{\psi}' \overleftrightarrow{\partial} \psi' + (\bar{\psi} P_R \gamma^\mu P_L T_L^a \psi + \bar{\psi}' P_L \gamma^\mu P_R T_R^a \psi') A_\mu^a \\ & = \int d^d x - \frac{1}{4g^2} \text{Tr} F_{\mu\nu} F^{\mu\nu} + \frac{i}{2} \bar{\psi} \overleftrightarrow{\partial} \psi + \frac{i}{2} \bar{\psi}' \overleftrightarrow{\partial} \psi' + (\bar{\psi} \bar{\gamma}^\mu P_L T_L^a \psi + \bar{\psi}' \bar{\gamma}^\mu P_R T_R^a \psi') A_\mu^a. \end{aligned}$$

We next generalize, in the obvious way, to the d -dimensional space-time of Dimensional Regularization the BRS variations (which we will denote by s_d and call Dimensional Regularization BRS variations) and the gauge-fixing terms. The Dimensional Regularization BRS transformations read

$$\begin{aligned} s_d \psi &= i\omega^a T_L^a P_L \psi, & s_d \bar{\psi} &= i\bar{\psi} T_L^a P_R \omega^a, \\ s_d \psi' &= i\omega^a T_R^a P_R \psi', & s_d \bar{\psi}' &= i\bar{\psi}' T_R^a P_L \omega^a, \\ s_d A_\mu &= \nabla_\mu \omega, & s_d \omega &= i\omega^2, & s_d \bar{\omega} &= B, & s_d B &= 0, \\ s_d \rho^\mu &= 0, & s_d \zeta &= 0, & s_d L &= s_d \bar{L} = s_d R = s_d \bar{R} = 0. \end{aligned} \tag{31}$$

Notice that we have chosen the Dimensional Regularization BRS transformations so that no explicit evanescent operator occurs in them. This will certainly simplify the computation of the anomalous breaking term in BRS identities for the minimally renormalized theory. We refer the reader to Appendix B for the discussion of the computation of the BRS anomalous breaking term when the Dimensional Regularization BRS transformations involve coefficients that vanish as $d \rightarrow 4$.

In summary, our Dimensional Regularization classical action, S_0 , has the following expression:

$$\begin{aligned} S_0 &= \int d^d x - \frac{1}{4g^2} \text{Tr} F_{\mu\nu} F^{\mu\nu} + \frac{i}{2} \bar{\psi} \overleftrightarrow{\partial} \psi + \frac{i}{2} \bar{\psi}' \overleftrightarrow{\partial} \psi' + (\bar{\psi} \bar{\gamma}^\mu P_L T_L^a \psi + \bar{\psi}' \bar{\gamma}^\mu P_R T_R^a \psi') A_\mu^a + \\ & \int d^d x \text{Tr} \frac{\alpha}{2} B^2 + \text{Tr} B(\partial_\mu A^\mu) - \text{Tr} \bar{\omega} \partial^\mu \nabla_\mu \omega + \\ & \int d^d x \text{Tr} \rho^\mu s_d A_\mu + \text{Tr} \zeta s_d \omega + \bar{L} s_d \psi + L s_d \bar{\psi} + \bar{R} s_d \psi' + R s_d \bar{\psi}'. \end{aligned} \tag{32}$$

The fermionic part of the Dimensional Regularization classical Lagrangian has the following non-gauge invariant form:


$$\frac{i}{2} \bar{\psi}_L \overleftrightarrow{\partial} \psi_L + \frac{i}{2} \bar{\psi}'_R \overleftrightarrow{\partial} \psi'_R + \frac{i}{2} \bar{\psi}_R \overleftrightarrow{\partial} \psi_R + \frac{i}{2} \bar{\psi}'_L \overleftrightarrow{\partial} \psi'_L$$

$$+ \frac{i}{2} \bar{\psi}_R \overleftrightarrow{\hat{\partial}} \psi_R + \frac{i}{2} \bar{\psi}_L \overleftrightarrow{\hat{\partial}} \psi_L + \frac{i}{2} \bar{\psi}'_L \overleftrightarrow{\hat{\partial}} \psi'_L + \frac{i}{2} \bar{\psi}'_R \overleftrightarrow{\hat{\partial}} \psi'_R. \quad (33)$$

Hence, S_0 is not BRS invariant, the breaking, $s_d S_0$, coming from the the last four terms of eq. (33):

$$\begin{aligned} s_d S_0 &= s_d \int d^d x \frac{i}{2} \bar{\psi} \hat{\gamma}^\mu P_L \overleftrightarrow{\partial}_\mu \psi + \frac{i}{2} \bar{\psi} P_R \hat{\gamma}^\mu \overleftrightarrow{\partial}_\mu \psi + \frac{i}{2} \bar{\psi}' \hat{\gamma}^\mu P_R \overleftrightarrow{\partial}_\mu \psi' + \frac{i}{2} \bar{\psi}' P_L \hat{\gamma}^\mu \overleftrightarrow{\partial}_\mu \psi' \\ &= \int d^d x \frac{1}{2} \omega^a \left\{ \bar{\psi} \hat{\gamma}^\mu \gamma_5 T_L^a \overleftrightarrow{\partial}_\mu \psi + \partial_\mu (\bar{\psi} \hat{\gamma}^\mu T_L^a \psi) - \bar{\psi}' \hat{\gamma}^\mu \gamma_5 T_R^a \overleftrightarrow{\partial}_\mu \psi' + \partial_\mu (\bar{\psi}' \hat{\gamma}^\mu T_R^a \psi') \right\} \\ &\equiv \hat{\Delta} \equiv \int d^d x \hat{\Delta}(x). \end{aligned} \quad (34)$$

The Feynman rule of the insertion of this anomalous breaking is given by fig. 1. We denote hereafter the line of the fermion which interact left(right)-handedly as L(R).



$$\begin{aligned} &\equiv \tilde{\Gamma}_{\psi^{(\prime)} \bar{\psi}^{(\prime)} \omega; \hat{\Delta}}^{\beta \alpha, a}(p_1, p_2; q) = \\ &= \frac{i}{2} [T_{L(R)}^a]_{ij} [+(-)(\hat{p}_2 - \hat{p}_1) \gamma_5 + (\hat{p}_1 + \hat{p}_2)] \Big|_{\alpha \beta} \end{aligned}$$

Figure 1. Feynman rule of the insertion of the non-integrated breaking eq. (34)

Notice that if the fermion representation(s) were compatible with a CP invariance of the classical action, then the Dimensional Regularization action with this choice of chiral vertex would also be formally CP invariant, and the breaking eq. (34) would have a definite CP value (+1 if we assign $(\omega_{L(R)})' = -(\omega_{L(R)})^t$, -1 if we assign $(\omega_{L(R)})' = (\omega_{L(R)})^t$, where $\omega_{L(R)} \equiv \omega^a T_{L(R)}^a$ and t stands for transposition of the colour matrices).

The breaking is an (implicit) “(d-4)-object”, *i.e.* an evanescent operator, and, clearly, this would be also true for any other choice Dimensional Regularization classical action. For example, if we had chosen as the regularized interaction

$$\bar{\psi} \gamma^\mu P_L T_L^a \psi A_\mu^a + \bar{\psi}' \gamma^\mu P_R T_R^a \psi' A_\mu^a,$$

the breaking would have been

$$\begin{aligned} &\int d^d x \frac{1}{2} \omega^a \left\{ \bar{\psi} \hat{\gamma}^\mu \gamma_5 T_L^a \overleftrightarrow{\partial}_\mu \psi + \partial_\mu (\bar{\psi} \hat{\gamma}^\mu \gamma_5 T_L^a \psi) - \bar{\psi}' \hat{\gamma}^\mu \gamma_5 T_R^a \overleftrightarrow{\partial}_\mu \psi' - \partial_\mu (\bar{\psi}' \hat{\gamma}^\mu \gamma_5 T_R^a \psi') \right\} \\ &\quad - c^{abc} \omega^b (\bar{\psi} \hat{\gamma}^\mu P_L T_L^a \psi + \bar{\psi}' \hat{\gamma}^\mu P_R T_R^a \psi') A_\mu^c \\ &\quad + i \omega^b (\bar{\psi} \hat{\gamma}^\mu P_L T_L^c T_L^b \psi + \bar{\psi}' \hat{\gamma}^\mu P_R T_R^c T_R^b \psi') A_\mu^c, \end{aligned}$$

which is certainly more involved. (And with this vertex, the formal CP invariance of the Dimensional Regularization classical action for CP invariant classical actions would be lost).

Remember that in the Breitenlohner and Maison prescription $\hat{\gamma}^\mu$ anticommutes whereas $\bar{\gamma}^\mu$ commutes with γ_5 ; so, in general the algebraic manipulations of the strings of γ s would be a bit more tedious with the second kind of vertex because $\gamma^\mu = \bar{\gamma}^\mu + \hat{\gamma}^\mu$.

Obviously, due to the fact that these two vertices are different, the results obtained by using minimal subtraction would be different—they would be *two different renormalization schemes*—and they should be related by finite terms. Moreover, in general, it should be expected that both results are different from the one obtained with the usual “naive” prescription of an anticommuting γ_5 every time a fermionic loop occurs. But the “hermitian vertex” of ref. [26] is not the only correct one as claimed there, but also any other one differing from it by an evanescent operator of dimension 4 as correctly remarked in [27].

The fact that in general the results obtained by minimal subtraction *à la* Breitenlohner and Maison in theories with cancellation of anomalies *do not satisfy* the Ward identities is a drawback from the practical point of view but not a reason to cast doubt on Breitenlohner and Maison schemes [28]. Indeed, we should always remember that we have the freedom to add *any* finite counterterm to restore the identities. And *in theories with anomalies the mathematical rigour of the regularization scheme we are considering has not been surpassed by any other dimensional regularization prescription*.

Furnished with the action S_0 given in eq. (32) one next develops a dimensionally regularized perturbation theory by following steps *i*) to *iv*) in subsection 2.2. This regularized theory is not invariant under the Dimensional Regularization BRS transformations in eq. (31). Indeed, the Dimensional Regularization 1PI functional Γ_0 obtained from S_0 satisfies anomalous BRS identities, the symmetry breaking terms being given by the insertion in Γ_0 of the operator $\hat{\Delta}$ in eq. (34).

Since our ultimate goal is to obtain a BRS invariant theory (if the anomaly cancellation conditions [23] are met) we shall further proceed and develop a dimensionally regularized perturbation theory by using the following action

$$S_{\text{DReg}}^{(n)} = S_0 + S_{\text{ct}}^{(n)}, \quad S_{\text{ct}}^{(n)} = S_{\text{sct}}^{(n)} + S_{\text{fct}}^{(n)}, \quad (35)$$

instead of the Dimensional Regularization classical action S_0 . We have named, in subsection 2.2., $S_{\text{DReg}}^{(n)}$ the *Dimensional Regularization action* (at order $\hbar^n$). It is computed order by order in the perturbative expansion in $\hbar$ by proceeding as in steps *v*) to *vii*) of subsection 2.2. Let us recall that $S_{\text{sct}}^{(n)}$ denotes the singular (at $d = 4$) counterterms needed to minimally renormalize the theory dimensionally up to order $\hbar^n$, whereas $S_{\text{fct}}^{(n)}$ stands for the finite (at $d = 4$) counterterms needed to turn the minimally renormalized theory into a BRS invariant theory up to order $\hbar^n$. In this paper we shall be concerned only with the renormalized theory at order $\hbar$. Hence, we will just compute $S_{\text{sct}}^{(1)}$ and $S_{\text{fct}}^{(1)}$.

Note that we have denoted by Γ_0 , instead of Γ_{DReg} , the Dimensional Regularization 1PI functional for S_0 to avoid confusion with the Dimensional Regularization 1PI functional for $S_{\text{DReg}}^{(n)}$, which shall denote by Γ_{DReg} .

3.3. The one-loop singular counterterms: $S_{\text{sct}}^{(1)}$

Let $\Gamma_{0\text{sing}}^{(1)}$ be the one-loop singular contribution at $d = 4$ to Γ_0 , the latter being the Dimensional Regularization 1PI functional obtained from S_0 . S_0 is given in eq. (32). Then, by definition

$$S_{\text{sct}}^{(1)} = -\Gamma_{0\text{sing}}^{(1)}. \quad (36)$$

We show the 1PI functions contributing to $S_{\text{sct}}^{(1)}$ in figs. 2 and 3.


Let $\varepsilon = 4 - d$. Then, the functional $\Gamma_{0\text{sing}}^{(1)}$ is given by

$$\begin{aligned} \Gamma_{0\text{sing}}^{(1)} = & \frac{\hbar}{\varepsilon} \frac{1}{(4\pi)^2} g^2 \left\{ -C_A \left[\frac{10}{3} + (1 - \alpha') \right] S_{0AA} + \frac{8}{3} \frac{T_{L+R}}{2} \bar{S}_{0AA} \right. \\ & - C_A \left[\frac{4}{3} + \frac{3}{2}(1 - \alpha') \right] S_{0AAA} + \frac{8}{3} \frac{T_{L+R}}{2} \bar{S}_{0AAA} \\ & - C_A \left[-\frac{2}{3} + 2(1 - \alpha') \right] S_{0AAAA} + \frac{8}{3} \frac{T_{L+R}}{2} \bar{S}_{0AAAA} \\ & + C_L 2\alpha' \bar{S}_{0\bar{\psi}\psi} + C_R 2\alpha' \bar{S}_{0\bar{\psi}'\psi'} \\ & + \left[\left(2 - \frac{(1 - \alpha')}{2} \right) C_A + 2\alpha' C_L \right] \bar{S}_{0\bar{\psi}\psi A} \\ & + \left[\left(2 - \frac{(1 - \alpha')}{2} \right) C_A + 2\alpha' C_R \right] \bar{S}_{0\bar{\psi}'\psi' A} \\ & - \left(1 + \frac{(1 - \alpha')}{2} \right) C_A S_{0\bar{\omega}\omega} + \alpha' C_A S_{0\bar{\omega}\omega A} \\ & + \alpha' C_A [S_{0\bar{L}s\psi} + S_{0\bar{R}s\psi'} + S_{0Ls\bar{\psi}} + S_{0Rs\bar{\psi}'}] \\ & - \left(1 + \frac{(1 - \alpha')}{2} \right) C_A S_{0\rho\omega} + \alpha' C_A S_{0\rho\omega A} \\ & \left. + \alpha' C_A S_{0\zeta\omega\omega} \right\} \\ & + \frac{\hbar}{\varepsilon} \frac{1}{(4\pi)^2} \frac{T_{L+R}}{2} \frac{4}{3} \int d^d x \frac{1}{2} \bar{A}_\mu \hat{\square} \bar{A}^\mu, \end{aligned} \quad (37)$$


where the S_{0X} terms are the corresponding (α -independent) terms of the Dimensional Regularization classical action S_0 . The bar above some of them means that all the index in the term are barred.

Let us define the Dimensional Regularization linearized BRS operator b_d as follows


$$\begin{aligned} b_d \equiv s_d + \int d^d x \left\{ \text{Tr} \frac{\delta S_0}{\delta A_\mu} \frac{\delta}{\delta \rho^\mu} + \text{Tr} \frac{\delta S_0}{\delta \omega} \frac{\delta}{\delta \zeta} + \right. \\ \left. + \frac{\delta S_0}{\delta \psi} \frac{\delta}{\delta \bar{L}} + \frac{\delta S_0}{\delta \psi'} \frac{\delta}{\delta \bar{R}} + \frac{\delta S_0}{\delta \bar{\psi}} \frac{\delta}{\delta L} + \frac{\delta S_0}{\delta \bar{\psi}'} \frac{\delta}{\delta R} \right\}, \end{aligned}$$




$$\begin{aligned}
& -\tilde{\Gamma}_{AA \ 0 \text{ sing}}^{(1) \ \mu_1 \mu_2, a_1 a_2}(k_1) = \\
& -\frac{i}{(4\pi)^2} C_A \delta^{a_1 a_2} \left[\frac{5}{3} + \frac{(1-\alpha')}{2} \right] \frac{2}{\varepsilon} (k_1^2 g^{\mu_1 \mu_2} - k_1^{\mu_1} k_1^{\mu_2}) \\
& -\frac{i}{(4\pi)^2} T_{L+R} \delta^{a_1 a_2} \frac{2}{3} \frac{2}{\varepsilon} \left[(\bar{k}_1^{\mu_1} \bar{k}_1^{\mu_2} - \bar{k}_1^2 \bar{g}^{\mu_1 \mu_2}) - \frac{1}{2} \hat{k}_1^2 \bar{g}^{\mu_1 \mu_2} \right]
\end{aligned}$$




$$\begin{aligned}
& -\tilde{\Gamma}_{AAA \ 0 \text{ sing}}^{(1) \ \mu_1 \mu_2 \mu_3, a_1 a_2 a_3}(k_1, k_2, k_3 = -k_1 - k_2) = \\
& -\frac{1}{(4\pi)^2} C_A c^{a_1 a_2 a_3} \left[\frac{2}{3} + \frac{3}{4}(1-\alpha') \right] \frac{2}{\varepsilon} \begin{bmatrix} g^{\mu_1 \mu_2}(k_1 - k_2)^{\mu_3} + \\ g^{\mu_1 \mu_3}(k_3 - k_1)^{\mu_2} + \\ g^{\mu_2 \mu_3}(k_2 - k_3)^{\mu_1} \end{bmatrix} \\
& + \frac{1}{(4\pi)^2} \frac{T_{L+R}}{2} c^{a_1 a_2 a_3} \frac{4}{3} \frac{2}{\varepsilon} \begin{bmatrix} \bar{g}^{\mu_1 \mu_2}(\bar{k}_1 - \bar{k}_2)^{\mu_3} + \\ \bar{g}^{\mu_1 \mu_3}(\bar{k}_3 - \bar{k}_1)^{\mu_2} + \\ \bar{g}^{\mu_2 \mu_3}(\bar{k}_2 - \bar{k}_3)^{\mu_1} \end{bmatrix}
\end{aligned}$$




$$\begin{aligned}
& -\tilde{\Gamma}_{AAAA \ 0 \text{ sing}}^{(1) \ \mu_1 \mu_2 \mu_3 \mu_4, a_1 a_2 a_3 a_4}(k_1, k_2, k_3, k_4 = -k_1 - k_2 - k_3) = \\
& + \frac{i}{(4\pi)^2} C_A \left[-\frac{1}{3} + (1-\alpha') \right] \frac{2}{\varepsilon} \begin{bmatrix} c^{e a_1 a_2} c^{a_3 a_4 e} (g^{\mu_2 \mu_3} g^{\mu_1 \mu_4} - g^{\mu_2 \mu_4} g^{\mu_1 \mu_3}) + \\ c^{e a_1 a_3} c^{a_4 a_2 e} (g^{\mu_3 \mu_4} g^{\mu_1 \mu_2} - g^{\mu_3 \mu_2} g^{\mu_1 \mu_4}) + \\ c^{e a_1 a_4} c^{a_2 a_3 e} (g^{\mu_1 \mu_3} g^{\mu_2 \mu_4} - g^{\mu_1 \mu_2} g^{\mu_4 \mu_3}) \end{bmatrix} \\
& - \frac{i}{(4\pi)^2} \frac{T_{L+R}}{2} \frac{4}{3} \frac{2}{\varepsilon} \begin{bmatrix} c^{e a_1 a_2} c^{a_3 a_4 e} (\bar{g}^{\mu_2 \mu_3} \bar{g}^{\mu_1 \mu_4} - \bar{g}^{\mu_2 \mu_4} \bar{g}^{\mu_1 \mu_3}) + \\ c^{e a_1 a_3} c^{a_4 a_2 e} (\bar{g}^{\mu_3 \mu_4} \bar{g}^{\mu_1 \mu_2} - \bar{g}^{\mu_3 \mu_2} \bar{g}^{\mu_1 \mu_4}) + \\ c^{e a_1 a_4} c^{a_2 a_3 e} (\bar{g}^{\mu_1 \mu_3} \bar{g}^{\mu_2 \mu_4} - \bar{g}^{\mu_1 \mu_2} \bar{g}^{\mu_4 \mu_3}) \end{bmatrix}
\end{aligned}$$




$$\begin{aligned}
& -\tilde{\Gamma}_{\omega \bar{\omega} \ 0 \text{ sing}}^{(1) \ ab}(p) = \\
& + \frac{i}{(4\pi)^2} C_A \delta^{ab} g^2 \frac{1}{2} \left[1 + \frac{(1-\alpha')}{2} \right] \frac{2}{\varepsilon} p^2
\end{aligned}$$



$$\begin{aligned}
& -\tilde{\Gamma}_{\omega \bar{\omega} A \ 0 \text{ sing}}^{(1) \ abc \ \mu}(p, q) = \\
& + \frac{1}{(4\pi)^2} C_A g^2 c^{abc} \frac{\alpha'}{2} \frac{2}{\varepsilon} p^\mu
\end{aligned}$$




$$\begin{aligned}
& -\tilde{\Gamma}_{\psi^{(\prime)} \bar{\psi}^{(\prime)} A \ 0 \text{ sing}}^{(1) \ \beta \alpha, ji}(p) = \\
& -\frac{i}{(4\pi)^2} g^2 \delta_{ji} C_{L(R)} \alpha' \frac{2}{\varepsilon} \bar{\not{p}} [P_{L(R)}]^{\alpha \beta}
\end{aligned}$$




$$\begin{aligned}
& -\tilde{\Gamma}_{\psi^{(\prime)} \bar{\psi}^{(\prime)} A \ 0 \text{ sing}}^{(1) \ \beta \alpha \ a \mu, ji}(p, q) = \\
& -\frac{i}{(4\pi)^2} g^2 \left[\left(1 - \frac{1-\alpha'}{4} \right) C_A + \alpha' C_{L(R)} \right] \frac{2}{\varepsilon} [T_{L(R)}^a]_{ij} \bar{\gamma}^\mu [P_{L(R)}]^{\alpha \beta}
\end{aligned}$$


Figure 2: Feynman rules for the order $\hbar$ singular counterterms. Here, only functions with no external fields are shown




$$-\tilde{\Gamma}_{\psi^{(\prime)}\omega;s\psi\ 0\ \text{sing}}^{(1)\ \beta a;\alpha i}(p;q) = -\frac{i}{(4\pi)^2}g^2C_A\frac{\alpha'}{2}\frac{2}{\varepsilon}[T_{L(R)}^a]^{ij}[P_{L(R)}]^{\alpha\beta}$$




$$-\tilde{\Gamma}_{\bar{\psi}^{(\prime)}\omega;s\bar{\psi}\ 0\ \text{sing}}^{(1)\ \beta a;\alpha i}(p;q) = \frac{i}{(4\pi)^2}g^2C_A\frac{\alpha'}{2}\frac{2}{\varepsilon}[T_{L(R)}^a]^{ij}[P_{R(L)}]^{\alpha\beta}$$



$$-\tilde{\Gamma}_{\omega A;sA\ 0\ \text{sing}}^{(1)\ e;a\ \mu}(;q) = \frac{i}{(4\pi)^2}g^2C_A\left(1+\frac{1-\alpha'}{2}\right)\frac{1}{\varepsilon}\delta^{ae}q^\mu$$



$$-\tilde{\Gamma}_{\omega A;sA\ 0\ \text{sing}}^{(1)\ e b\nu;a\ \mu}(p;q) = -\frac{1}{(4\pi)^2}g^2C_A\alpha'\frac{1}{\varepsilon}c^{abe}$$



$$-\tilde{\Gamma}_{\omega\omega;s\omega\ 0\ \text{sing}}^{(1)\ ab\ c}(p;q) = \frac{1}{(4\pi)^2}g^2C_A\alpha'\frac{1}{\varepsilon}c^{abc}$$

Figure 3: The same as for fig. 2 but with functions involving external fields

Notice that $b_d^2 \neq 0$ since $s_d S_0 \neq 0$. The action of s_d on the fields has been defined in eq. (31). If we define in the d -dimensional space-time of Dimensional Regularization the following integrated field polynomials

$$\begin{aligned} L_g &\equiv \frac{1}{2g^2} \int d^d x \text{Tr } F_{\mu\nu} F^{\mu\nu} = g \frac{\partial}{\partial g} S_0, & \bar{L}_g &\equiv \frac{1}{2g^2} \int d^d x \text{Tr } \bar{F}_{\mu\nu} \bar{F}^{\mu\nu}, \\ L_{\psi^{(\prime)}}^{L(R)} &\equiv -b_d \cdot \int d^d x \{ \bar{L}(\bar{R}) P_{L(R)} \psi^{(\prime)} + \bar{\psi}^{(\prime)} P_{R(L)} L(R) \} = \left(N_{\psi}^{L(R)} - N_{L(R)} \right) S_0 = \\ &= 2 \int d^d x \left\{ \frac{i}{2} \bar{\psi}^{(\prime)} \overleftrightarrow{\not{D}} P_{L(R)} \psi^{(\prime)} + \bar{\psi}^{(\prime)} \bar{\gamma}^\mu P_{L(R)} T_{L(R)}^a \psi^{(\prime)} A_\mu^a \right\}, \\ \bar{L}_{\psi^{(\prime)}}^{L(R)} &\equiv L_{\psi^{(\prime)}} - \int d^d x i \bar{\psi}^{(\prime)} \overleftrightarrow{\not{D}} P_{L(R)} \psi^{(\prime)} = 2 \int d^d x \left\{ \frac{i}{2} \bar{\psi}^{(\prime)} \overleftrightarrow{\not{D}} P_{L(R)} \psi^{(\prime)} + \bar{\psi}^{(\prime)} \bar{\gamma}^\mu P_{L(R)} T_{L(R)}^a \psi^{(\prime)} A_\mu^a \right\}, \end{aligned}$$




Figure 4: One-loop Feynman graphs contributing to the 1PI functions involving external fields

$$L_A \equiv b_d \cdot \int d^d x \tilde{\rho}_\mu^a A^{a\mu} = \left(\text{Tr} [N_A - N_\rho - N_B - N_{\bar{\omega}}] + 2\alpha \frac{\partial}{\partial \alpha} \right) S_0,$$

$$L_\omega \equiv -b_d \cdot \int d^d x \zeta^a \omega^a = \text{Tr} [N_\omega - N_\zeta] S_0,$$

where

$$\begin{aligned} N_\phi &\equiv \int d^d x \phi(x) \frac{\delta}{\delta \phi(x)}, & \phi &= A_\mu, \rho_\mu, B, \bar{\omega}, \omega \text{ and } \zeta, \\ N_\psi^{L(R)} &\equiv \int d^d x (P_{L(R)})_\beta \psi_\beta \frac{\delta}{\delta \psi_\beta}, & N_{\psi'}^{R(L)} &\equiv \int d^d x (P_{R(L)})_\beta \psi'_\beta \frac{\delta}{\delta \psi'_\beta}, \\ N_{\bar{\psi}}^{R(L)} &\equiv \int d^d x (\bar{\psi} P_{R(L)})_\beta \frac{\delta}{\delta \bar{\psi}_\beta}, & N_{\bar{\psi}'}^{L(R)} &\equiv \int d^d x (\bar{\psi}' P_{L(R)})_\beta \frac{\delta}{\delta \bar{\psi}'_\beta}; \end{aligned} \quad (38)$$

then, the functional $\Gamma_{0\text{sing}}^{(1)}$ in eq. (37) can be cast into the form:

$$\Gamma_{0\text{sing}}^{(1)} = \frac{1}{(4\pi)^2} g^2 \frac{\hbar}{\varepsilon} \left[\frac{11}{3} C_A L_g + \left(2 - \frac{1 - \alpha'}{2} \right) C_A L_A + \alpha' C_A L_\omega \right]$$

$$\begin{aligned}
& + \frac{1}{(4\pi)^2} g^2 \frac{\hbar}{\varepsilon} \left[-\frac{4}{3} \frac{T_{L+R}}{2} \bar{L}_g + \alpha' C_L \bar{L}_\psi + \alpha' C_R \bar{L}_{\psi'} \right] \\
& + \frac{1}{(4\pi)^2} \frac{\hbar}{\varepsilon} \frac{T_{L+R}}{2} \frac{4}{3} \int d^d x \frac{1}{2} \bar{A}_\mu \hat{\Delta} \bar{A}^\mu;
\end{aligned} \tag{39}$$

i.e. $b_d \Gamma_{0\text{sing}}^{(1)} = O(\hat{g}_{\mu\nu})$.

The second and third lines of the previous equation show explicitly that the standard “infinite” multiplicative renormalization of the coupling constant g and of the fields of S_0 is already lost at the one-loop level.

3.4. The BRS identity and the anomalous symmetry breaking term

The *Regularized Action Principle*, explained in subsection 2.3. and Appendix A, leads to the following BRS identity

$$\begin{aligned}
\mathcal{S}_d(\Gamma_{\text{DReg}}) & \equiv \int d^d x \left\{ \text{Tr} \frac{\delta \Gamma_{\text{DReg}}}{\delta \rho^\mu} \frac{\delta \Gamma_{\text{DReg}}}{\delta A_\mu} + \text{Tr} \frac{\delta \Gamma_{\text{DReg}}}{\delta \zeta} \frac{\delta \Gamma_{\text{DReg}}}{\delta \omega} + \text{Tr} B \frac{\delta \Gamma_{\text{DReg}}}{\delta \bar{\omega}} \right. \\
& \quad \left. + \frac{\delta \Gamma_{\text{DReg}}}{\delta \bar{L}} \frac{\delta \Gamma_{\text{DReg}}}{\delta \psi} + \frac{\delta \Gamma_{\text{DReg}}}{\delta \bar{R}} \frac{\delta \Gamma_{\text{DReg}}}{\delta \psi'} + \frac{\delta \Gamma_{\text{DReg}}}{\delta L} \frac{\delta \Gamma_{\text{DReg}}}{\delta \bar{\psi}} + \frac{\delta \Gamma_{\text{DReg}}}{\delta R} \frac{\delta \Gamma_{\text{DReg}}}{\delta \bar{\psi}'} \right\} = \\
& = \hat{\Delta} \cdot \Gamma_{\text{DReg}} + \Delta_{\text{ct}} \cdot \Gamma_{\text{DReg}} + \int d^d x \sum_{\Phi} \left\{ \left[\frac{\delta S_{\text{ct}}^{(n)}}{\delta K_\Phi(x)} \cdot \Gamma_{\text{DReg}} \right] \frac{\delta \Gamma_{\text{DReg}}}{\delta \Phi(x)} \right\},
\end{aligned} \tag{40}$$

when applied to the Dimensional Regularization 1PI functional, Γ_{DReg} , obtained from the Dimensional Regularization action $S_{\text{DReg}}^{(n)}$, the latter being given in eq. (35). The symbols Φ and K_Φ in the previous equation stand, respectively, for any field that undergoes non-linear BRS transformation, *i.e.* A_μ , ω , ψ , ψ' , $\bar{\psi}$ and $\bar{\psi}'$ and the external field which couple to the corresponding BRS variation, *i.e.* ρ_μ , ζ , $\bar{L}$, $\bar{R}$, L , and R . The operator $\hat{\Delta}$ is given in eq. (34). The symbol Δ_{ct} is equal to $s_d S_{\text{ct}}^{(n)}$, where $S_{\text{ct}}^{(n)}$ and s_d are defined in eqs. (35) and (31), respectively. Eq. (40) is a particular instance of eq. (19), and it is obtained by considering the concrete realization of the Dimensional Regularization BRS transformations given in eq. (31). This equation is a rigorous equation, valid to all orders in the expansion in powers of $\hbar$ and fields.

Notice that Γ_{DReg} is not BRS invariant since $\mathcal{S}_d(\Gamma_{\text{DReg}})$ does not vanish. Indeed, the far right hand side of eq. (40),

$$\hat{\Delta} \cdot \Gamma_{\text{DReg}} + \Delta_{\text{ct}} \cdot \Gamma_{\text{DReg}} + \int d^d x \sum_{\Phi} \left\{ \left[\frac{\delta S_{\text{ct}}^{(n)}}{\delta K_\Phi(x)} \cdot \Gamma_{\text{DReg}} \right] \frac{\delta \Gamma_{\text{DReg}}}{\delta \Phi(x)} \right\},$$

is not zero. This term is the symmetry breaking term of the dimensionally regularized theory defined by the Dimensional Regularization action $S_{\text{DReg}}^{(n)}$.

We have to ask now about the renormalized counterpart of eq. (40). Since $S_{\text{DReg}}^{(n)}$ contains the singular counterterms needed to render non-singular at $d = 4$ the 1PI functions

of the theory up to order $\hbar^n$, eq. (20) defines the renormalized 1PI functional Γ_{ren} up to order $\hbar^n$:

$$\text{LIM}_{d \rightarrow 4} \Gamma_{\text{DReg}}[\varphi, \Phi; K_\Phi; \mu] = \Gamma_{\text{ren}}[\varphi, \Phi; K_\Phi; \mu]. \quad (41)$$

Let us recall that the limiting process denoted by $\text{LIM}_{d \rightarrow 4}$ is accomplished by taking the ordinary limit $d \rightarrow 4$ first and then replacing with zero every hatted object. The Dimensional Regularization scale has been introduced by hand as explained in *v)* of subsection 2.2.

Now, by particularizing eqs. (18)–(20) to the field theory under study, one concludes that the following equation, which is the renormalized counterpart of eq. (40), holds up to order $\hbar^n$:

$$\begin{aligned} \mathcal{S}(\Gamma_{\text{ren}}) \equiv \int d^4x \left\{ \text{Tr} \frac{\delta \Gamma_{\text{ren}}}{\delta \rho^\mu} \frac{\delta \Gamma_{\text{ren}}}{\delta A_\mu} + \text{Tr} \frac{\delta \Gamma_{\text{ren}}}{\delta \zeta} \frac{\delta \Gamma_{\text{ren}}}{\delta \omega} + \text{Tr} B \frac{\delta \Gamma_{\text{ren}}}{\delta \bar{\omega}} \right. \\ \left. + \frac{\delta \Gamma_{\text{ren}}}{\delta \bar{L}} \frac{\delta \Gamma_{\text{ren}}}{\delta \psi} + \frac{\delta \Gamma_{\text{ren}}}{\delta \bar{R}} \frac{\delta \Gamma_{\text{ren}}}{\delta \psi'} + \frac{\delta \Gamma_{\text{ren}}}{\delta L} \frac{\delta \Gamma_{\text{ren}}}{\delta \bar{\psi}} + \frac{\delta \Gamma_{\text{ren}}}{\delta R} \frac{\delta \Gamma_{\text{ren}}}{\delta \bar{\psi}'} \right\} = \Delta_{\text{breaking}}, \end{aligned} \quad (42)$$

where Γ_{ren} is the dimensionally renormalized 1PI functional as defined by eq. (41). The BRS symmetry breaking term Δ_{breaking} is given up to order $\hbar^n$ by the following “limit”

$$\Delta_{\text{breaking}} = \text{LIM}_{d \rightarrow 4} \left\{ \hat{\Delta} \cdot \Gamma_{\text{DReg}} + \Delta_{\text{ct}} \cdot \Gamma_{\text{DReg}} + \int d^d x \sum_{\Phi} \left\{ \left[\frac{\delta S_{\text{ct}}^{(n)}}{\delta K_\Phi(x)} \cdot \Gamma_{\text{DReg}} \right] \frac{\delta \Gamma_{\text{DReg}}}{\delta \Phi(x)} \right\} \right\}. \quad (43)$$

The fact [18] that if the anomaly cancellation conditions are met [23] the cohomology of the linearized BRS operator b in eq. (29) is trivial on the space of local polynomials of ghost number one, guarantees that $S_{\text{fct}}^{(n)}$ can be chosen so that Δ_{breaking} above vanishes whatever the value of n . Hence, if the BRS symmetry is non-anomalous, non-symmetric counterterms can be added to the Dimensional Regularization classical action so that the resulting renormalized action is BRS invariant: $\mathcal{S}(\Gamma_{\text{ren}}) = 0$. In this paper we shall compute $S_{\text{fct}}^{(1)}$ explicitly (see subsection 3.6.).

Let us close this subsection and express in terms of local operators in four dimensions the order $\hbar$ contribution, $\Delta_{\text{breaking}}^{(1)}$, to the symmetry breaking term given in eq. (43). Adapting eqs. (21) and (23) to the regularized theory at hand we conclude that

$$\begin{aligned} \Delta_{\text{breaking}}^{(1)} &= \text{LIM}_{d \rightarrow 4} \left\{ \left[\hat{\Delta} \cdot \Gamma_{\text{DReg}} \right]_{\text{singular}}^{(1)} + b_d S_{\text{sct}}^{(1)} \right\} + \left[N[\hat{\Delta}] \cdot \Gamma_{\text{ren}} \right]^{(1)} + b S_{\text{fct}}^{(1)} \\ &= \left[N[\hat{\Delta}] \cdot \Gamma_{\text{ren}} \right]^{(1)} + b S_{\text{fct}}^{(1)}. \end{aligned} \quad (44)$$

where $\hat{\Delta}$ and b_d are defined in eqs. (34) and (29), respectively. Indeed, we shall show below that

$$\left[\hat{\Delta} \cdot \Gamma_{\text{DReg}} \right]_{\text{singular}}^{(1)} = \frac{1}{(4\pi)^2} \frac{T_{\text{L+R}}}{2} \frac{4}{3} \frac{\hbar}{\varepsilon} \int d^d x -\omega^a \{ \hat{\square} \bar{\partial}_\mu \bar{A}^{\mu a} + c^{abc} (\hat{\square} \bar{A}_\mu^a) \bar{A}^{\mu b} \} = -b_d S_{\text{sct}}^{(1)}.$$

The order $\hbar$ singular (pole) contribution, $\left[\hat{\Delta} \cdot \Gamma_{\text{DReg}}\right]_{\text{singular}}^{(1)}$, to $\Delta \cdot \Gamma_{\text{DReg}}$ only comes from the diagrams depicted in figs. 5 and 6, after replacing $\hat{\Delta}$ with $\hat{\Delta}$. These contributions read

$$\begin{aligned} \left[\left[\hat{\Delta} \cdot \Gamma_{\text{DReg}}\right]_{Aw}^{\mu ba}\right]_{\text{singular}}^{(1)}(p) &= \frac{i}{(4\pi)^2} T_{L+R} \delta^{ab} \frac{2}{3} \frac{1}{\varepsilon} \bar{p}^\mu \hat{p}^2; \\ \left[\left[\hat{\Delta} \cdot \Gamma_{\text{DReg}}\right]_{AAw}^{\mu\nu bca}\right]_{\text{singular}}^{(1)}(p_1, p_2) &= \frac{i}{(4\pi)^2} i c^{abc} \frac{4}{3} \frac{1}{\varepsilon} (\hat{p}_1^2 - \hat{p}_2^2) \bar{g}^{\mu\nu}. \end{aligned} \quad (45)$$

Now, $b_d L_g = b_d L_A = b_d L_\omega = b_d \bar{L}_g = b_d \bar{L}_{\psi^{(\nu)}} = 0$, and therefore the only non-vanishing b_d -variation of $\Gamma_{0\text{sing}}^{(1)}$ comes from the third line of eq. (39), which in turns is equal to eq. (45). The fact that $S_{\text{sct}}^{(1)} = -\Gamma_{0\text{sing}}^{(1)}$ concludes the proof.

The next task to face is the computation of the order $\hbar$ contribution to the anomalous insertion $N[\hat{\Delta}] \cdot \Gamma_{\text{ren}}$. We shall carry out this calculation in the next section. The computation is somewhat simplified if we bare in mind that the field B , enforcing the gauge-fixing condition, has no dynamics (no interaction vertices) unless forcibly introduced by appropriate choice of $S_{\text{fct}}^{(n)}$ (finite counterterms). Hence, it will be very advisable to impose that the finite counterterms be independent of B , so that the only contribution to Γ_{DReg} involving B is order $\hbar^0$ and given by S_0 in eq. (32). It is thus plain that the so-called gauge-fixing equation

$$\mathcal{B}(\Gamma_{\text{ren}}) \equiv \frac{\delta \Gamma_{\text{ren}}}{\delta B} - \partial_\mu A^\mu - \alpha B = 0, \quad (46)$$

holds for the renormalized theory. Notice that eq. (46) does not clash with restoring the BRS symmetry since Δ_{breaking} defined in eq. (43) does not depend on B . Of course, eq. (46) is the equation of motion of B .

Finally, the ghost equation is another interesting equation. It is just the equation of motion of $\bar{\omega}$:

$$\mathcal{G} \Gamma_{\text{ren}} \equiv \left\{ \frac{\delta}{\delta \bar{\omega}} + \partial_\mu \frac{\delta}{\delta \rho_\mu} \right\} \Gamma_{\text{ren}} = 0, \quad (47)$$

and is a consequence of eq. (42) and the gauge-fixing equation. Some functional differentiation and the fact that Δ_{breaking} does not depend on B lead to eq. (47). A further simplification, the ghost equation implies that the functional which satisfy it depends on ρ_μ and $\bar{\omega}$ through the combination $\tilde{\rho}_\mu = \rho_\mu + \partial_\mu \bar{\omega}$. Finally, it is not difficult to come to the conclusion that this very simplification applies to each term on the right hand side of eq. (43) and $S_{\text{ct}}^{(n)}$ in eq. (35).

3.5. Expansion of the anomalous insertion

The anomalous insertion $N[\hat{\Delta}] \cdot \Gamma_{\text{ren}}$ has ghost number +1 and it is the insertion into the 1PI functional Γ_{ren} of an evanescent integrated polynomial operator of ultraviolet dimension 4 with neither free Lorentz indices nor free indices for the gauge group. This anomalous insertion is a functional of ψ , ψ' , A_μ , $\bar{\omega}$ (only through $\tilde{\rho}_\mu$), ω , but not of B , and

also of the external fields $L, \bar{L}, R, \bar{R}, \rho_\mu$ (only through $\tilde{\rho}_\mu$) and ζ . It is thus compulsory to generalize first eq. (25) so that it includes also Feynman diagrams with the external fields as vertices.

Due to the explicit power-counting in our model, only diagrams with either no or one external fields can be divergent. Consider a diagram with the vertex $K_\Phi s\Phi$ (Φ denotes any field which undergoes non-linear BRS transformations). The Feynman rule in momentum space of this vertex is an integration over a momentum q' , which is absorbed in the definition of the Fourier transform of the function with an insertion of an operator, and the factors $\frac{i}{\hbar} \tilde{K}_\Phi(q')$ times the Feynman rule of the operator insertion $s\Phi$ at momentum q' . The singular part is also a polynomial in q' , therefore it can be expanded, together with the rest of momenta, in a finite Taylor series in q' . A factor $q'^{\mu_1} \dots q'^{\mu_s}$ together with $\tilde{K}(q')$ will lead to a $(-i)^s \partial_{\mu_1} \dots \partial_{\mu_s} K_\Phi(x)$ which multiply the insertion of the monomial operator obtained with the rest of momenta. Therefore the Bonneau identities (eq. (25)) are generalized to:

$$\begin{aligned}
N[\hat{\Delta}](x) \cdot \Gamma_{\text{ren}} = & - \sum_{n=0}^4 \sum_{\{j_1 \dots j_n\}} \left[\sum_{r=0}^{\delta(J)} \sum_{\substack{\{i_1 \dots i_r\} \\ 1 \leq i_j \leq n}} \right. \\
& \left\{ \frac{(-i)^r}{r!} \frac{\partial^r}{\partial p_{i_1}^{\mu_1} \dots \partial p_{i_r}^{\mu_r}} (-i\hbar) \text{r.s.p.} \overline{\langle \tilde{\phi}_{j_1}(p_1) \dots \tilde{\phi}_{j_n}(p_n) N[\check{\Delta}](q=-\sum p_i) \rangle_{K=0}^{\text{1PI}}} \Big|_{p_i=0, \check{g}=0} \right\} \\
& \times N \left[\frac{1}{n!} \prod_{k=n}^1 \left\{ \left(\prod_{\{\alpha/i_\alpha=k\}} \partial_{\mu_\alpha} \right) \phi_{j_k} \right\} \right] (x) \cdot \Gamma_{\text{ren}} + \\
& + \sum_{\Phi} \sum_{\substack{s,t \\ 0 \leq s+t \leq \delta(J;\Phi)}} \sum_{\substack{\{i_1 \dots i_s\} \\ 1 \leq i_j \leq n}} \left\{ \frac{(-i)^{s+t}}{(s+t)!} \frac{\partial^{s+t}}{\partial p_{i_1}^{\mu_1} \dots \partial p_{i_s}^{\mu_s} \partial p_{n+1}^{\nu_1} \dots \partial p_{n+1}^{\nu_t}} \right. \\
& \left. \text{r.s.p.} \overline{\langle \tilde{\phi}_{j_1}(p_1) \dots \tilde{\phi}_{j_n}(p_n); N[s\Phi](p_{n+1}) N[\check{\Delta}](q=-\sum p_i) \rangle_{K=0}^{\text{1PI}}} \Big|_{p_i=0, \check{g}=0} \right\} \\
& \times (\partial_{\nu_1} \dots \partial_{\nu_t} K_\Phi)(x) N \left[\frac{1}{n!} \prod_{k=n}^1 \left\{ \left(\prod_{\{\alpha/i_\alpha=k\}} \partial_{\mu_\alpha} \right) \phi_{j_k} \right\} \right] (x) \cdot \Gamma_{\text{ren}} \quad \Bigg] , \tag{48}
\end{aligned}$$

where $J \equiv \{j_1, \dots, j_n\}$ and the indexes j_k , $k = 1, \dots, n$ label the different types of quantum fields, with the proviso that fields having different values of gauge group indices are taken as different. The symbols $\delta(J)$, $\delta(J; \Phi)$ denote, respectively, the overall degrees of ultraviolet divergence of the 1PI functions $\langle \tilde{\phi}_{j_1}(p_1) \dots \tilde{\phi}_{j_n}(p_n) N[\check{\Delta}](q=-\sum p_i) \rangle_{K=0}^{\text{1PI}}$ and $\langle \tilde{\phi}_{j_1}(p_1) \dots \tilde{\phi}_{j_n}(p_n); N[s\Phi](p_{n+1}) N[\check{\Delta}](q=-\sum p_i) \rangle_{K=0}^{\text{1PI}}$. Notice that since our propagators are massless the only non-vanishing contributions to the left hand side of eq. (48) come from $r = \delta(J)$ and $s + t = \delta(J; \Phi)$. $\prod_{k=n}^1$ means that fields in the product are ordered from left to right according to decreasing values of k .

The previous formula gives rise to an expansion of $N[\hat{\Delta}](x) \cdot \Gamma_{\text{ren}}$ in terms of both standard and evanescent monomials [6,7] (see section 2.4.). If we denote the monomials generically by the letter $\mathcal{M}$, the expansion will have the form

$$\begin{aligned}
N[\hat{\Delta}] \cdot \Gamma_{\text{ren}} &= \int d^4x N[\hat{\Delta}](x) \cdot \Gamma_{\text{ren}} \\
&= \sum_i \beta_i^{abc\dots} N[\mathcal{M}_i^{abc\dots}] \cdot \Gamma_{\text{ren}} + \sum_j \hat{\beta}_j^{abc\dots} N[\hat{\mathcal{M}}_j^{abc\dots}] \cdot \Gamma_{\text{ren}} \\
&= \sum_i \bar{\beta}_i^{abc\dots} N[\bar{\mathcal{M}}_i^{abc\dots}] \cdot \Gamma_{\text{ren}} + \sum_j \hat{\beta}_j^{abc\dots} N[\hat{\mathcal{M}}_j^{abc\dots}] \cdot \Gamma_{\text{ren}}, \tag{49}
\end{aligned}$$

with each coefficient being a formal series in $\hbar$ (starting at order $\hbar^1$). The latin letters abc stand for the gauge group indices. We shall assume that repeated gauge group indices are summed over. The monomials in eq. (49) have no free Lorentz index. The objects $\{\bar{\mathcal{M}}_i^{abc\dots}\}$, called “barred” nonomials, are monomials where all Lorentz contractions are carried out with $\bar{g}^{\mu\nu}$. Notice that there is a one-to-one correspondence between $\{\mathcal{M}_i^{abc\dots}\}$ and $\{\bar{\mathcal{M}}_i^{abc\dots}\}$. The objects $\{\hat{\mathcal{M}}_i^{abc\dots}\}$, called “hatted” nonomials, are evanescent operators.

Notice that in our case the monomials will be integrated local functionals of the fields and its derivatives with ghost number one and ultraviolet dimension 4.

Expanding in the same way all the normal insertions of the evanescent monomials $\hat{\mathcal{M}}_i^{abc\dots}$ on the right hand side of eq. (49) and solving the system of Bonneau identities we would get finally the true expansion of the anomalous insertions in terms of the basis of standard (or barred) normal products (also called standard quantum insertions):

$$\begin{aligned}
N[\hat{\Delta}] \cdot \Gamma_{\text{ren}} &= \sum_i k_i^{abc\dots} N[\mathcal{M}_i^{abc\dots}] \cdot \Gamma_{\text{ren}} \\
&= \sum_i \bar{k}_i^{abc\dots} N[\bar{\mathcal{M}}_i^{abc\dots}] \cdot \Gamma_{\text{ren}};
\end{aligned}$$

where the barred monomials simply means again that all the Lorentz contractions are done with the $\bar{g}$ tensor.

Let us shorthand $\int d^d x$ to $\int$. The list of integrated monomials of ghost number 1, ultraviolet dimension 4 and no free of Lorentz indices which is relevant to our computation is the following:

* a) *Monomials with 1 ω and 1 A*

$$\mathcal{M}_1^{ab} \equiv \int d^d x \omega^a \square \partial_\mu A^{b\mu}.$$

* b) *Monomials with 1 ω and 2 A 's and no $\varepsilon_{\mu\nu\alpha\beta}$ tensor*

$$\begin{aligned}
\mathcal{M}_2^{abc} &\equiv \int \omega^a (\square A_\mu^b) A^{c\mu}, & \mathcal{M}_3^{abc} &\equiv \int \omega^a (\partial_\mu A_\nu^b) (\partial^\mu A^{c\nu}) = \mathcal{M}_3^{acb}, \\
\mathcal{M}_4^{abc} &\equiv \int \omega^a (\partial_\mu \partial_\nu A^{b\mu}) A^{c\nu}, & \mathcal{M}_5^{abc} &\equiv \int \omega^a (\partial_\mu A^{b\mu}) (\partial_\nu A^{c\nu}) = \mathcal{M}_5^{acb}, \\
\mathcal{M}_6^{abc} &\equiv \int \omega^a (\partial_\mu A_\nu^b) (\partial^\nu A^{c\mu}) = \mathcal{M}_6^{acb}.
\end{aligned}$$

* c) *Monomials with 1 ω and 3 A 's and no $\varepsilon_{\mu\nu\alpha\beta}$ tensor*

$$\mathcal{M}_7^{abcd} \equiv \int \omega^a (\partial_\mu A^{b\mu}) A_\nu^c A^{d\nu} = \mathcal{M}_7^{abdc}, \quad \mathcal{M}_8^{abcd} \equiv \int \omega^a (\partial_\mu A_\nu^b) A^{c\mu} A^{d\nu}.$$

* d) *Monomials with 1 ω and 4 A 's and no $\varepsilon_{\mu\nu\alpha\beta}$ tensor*

$$\mathcal{M}_9^{abcde} \equiv \int \omega^a A_\mu^b A^{c\mu} A_\nu^d A^{e\nu} = \mathcal{M}_9^{acbde} = \mathcal{M}_9^{abced} = \mathcal{M}_9^{adebc}.$$

Since the formulae with fermions ψ and ψ' are very similar we collect them in a way obvious to interpret. In general, we will denote with a roman letter L or R the fermionic lines or loops in Feynman diagrams, the monomials, their coefficients, ... corresponding to the fermions ψ , whose interaction is left-handed, or to the fermions ψ' , whose interaction is right-handed.

* e) *Monomials with 1 ω and $\psi, \bar{\psi}$ ($\psi', \bar{\psi}'$)*

$$\mathcal{M}_{10L(R)}^{a,ij} \equiv \int \omega^a \bar{\psi}_i^{(\prime)} \bar{\gamma}^\mu P_{L(R)} \partial_\mu \psi_j^{(\prime)}, \quad \mathcal{M}_{11L(R)}^{a,ij} \equiv \int \omega^a (\partial_\mu \bar{\psi}_i^{(\prime)}) \bar{\gamma}^\mu P_{L(R)} \partial_\mu \psi_j^{(\prime)},$$

i, j denote the group indices of the corresponding fermion fields.

* f) *Monomials with 1 ω , 1 A and $\psi, \bar{\psi}$ ($\psi', \bar{\psi}'$)*

$$\mathcal{M}_{12L(R)}^{ab,ij} \equiv \int \omega^a \bar{\psi}_i^{(\prime)} \bar{\gamma}^\mu P_{L(R)} \psi_j^{(\prime)} A_\mu^b.$$

* g) *Monomials with fermions and external fields*

$$\mathcal{M}_{14L(R)}^{ab,ij} \equiv \int \omega^a \omega^b \bar{L}_i(R_i) P_{L(R)} \psi_j^{(\prime)}, \quad \mathcal{M}_{15L(R)}^{ab,ij} \equiv \int \omega^a \omega^b \bar{\psi}_i^{(\prime)} P_{R(L)} L_j(R_j)$$

We have not considered operators like $\omega^a \bar{\psi}_i \bar{\gamma}^\mu P_R \partial_\mu \psi_j$ as admissible monomials since it will not be generated by the Bonneau identities due to the form of our choice for fermion vertex at order $\hbar^0$.

We shall adopt the notation that the $\{\dots\}$ enclosing indices denotes symmetrization and $[\dots]$ antisymmetrization.

* h) *Monomials with $\varepsilon_{\mu\nu\alpha\beta}$*

$$\mathcal{M}_{50}^{abc} \equiv \int \varepsilon_{\mu\nu\alpha\beta} \omega^a (\partial^\alpha A^{b\mu}) (\partial^\beta A^{c\nu}) = \mathcal{M}_{50}^{acb}, \mathcal{M}_{51}^{abcd} \equiv \int \varepsilon_{\mu\nu\rho\alpha} \omega^a (\partial^\alpha A^{b\mu}) A^{c\nu} A^{d\rho} = -\mathcal{M}_{51}^{abdc},$$

$$\mathcal{M}_{52}^{abc} \equiv \int \varepsilon_{\mu\nu\rho\lambda} \omega^a A^{b\mu} A^{c\nu} A^{d\rho} A^{e\lambda} = \mathcal{M}_{52}^{a[bcd]e};$$

* i) *Monomials with $\tilde{\rho}$ or ζ*

Notice that it is possible to construct other monomials with ghost number +1 and dimension 4: $\tilde{\rho}_\mu^a \omega^b \omega^c A^{d\mu}$, $\tilde{\rho}_\mu^a (\partial^\mu \omega^b) \omega^c$, $\zeta^a \omega^b \omega^c \omega^d$. The formulae for their coefficients in the Bonneau expansion are also easily obtained, but we do not write them because in the one loop computation all this coefficients turn out to be zero. This is clear because all the 1PI functions we would have to compute involve at least two loops.

* *Anomalous monomials, i.e. with some $\hat{g}$*

Take all the monomials written above and write all monomials obtained by adding hats to the Lorentz indices in all possible ways. No hatted index should appear contracted with the $\varepsilon_{\mu\nu\alpha\beta}$, because this contraction vanish. Notice that with the fermionic vertices $\bar{\psi}^{(\prime)} \bar{\gamma}^\mu P_{L(R)} T_{L(R)}^a \psi^{(\prime)} A_\mu^a = \bar{\psi}^{(\prime)} P_{R(L)} \bar{\gamma}^\mu P_{L(R)} T_{L(R)}^a \psi^{(\prime)} A_\mu^a$ in the Dimensional Regularization classical action S_0 , neither $\omega^a \bar{\psi}_i P_L \hat{\gamma}^\mu P_L \partial_\mu \psi_j$ nor $\omega^a \bar{\psi}_i^{(\prime)} P_R \hat{\gamma}^\mu P_R \partial_\mu \psi_j^{(\prime)}$ occur in the

Bonneau expansion we are considering; however, with the vertex $\bar{\psi}^{(\iota)} \gamma^\mu P_{L(R)} T_{L(R)}^a \psi^{(\iota)} A_\mu^a = A_\mu^a (\bar{\psi}^{(\iota)} P_{R(L)} \bar{\gamma}^\mu P_{L(R)} T_{L(R)}^a \psi^{(\iota)} + \bar{\psi}^{(\iota)} P_{L(R)} \hat{\gamma}^\mu T_{L(R)}^a P_{L(R)} \psi^{(\iota)})$, both do.

It is convenient to use the expansion in terms of barred and hatted monomials (so the 1PI functions with $\check{\Delta}$ inserted have to be expressed in terms of $\bar{g}$ and $\hat{g}$). Then, the formulae of the coefficients for all orders and the result for the first order read (see figs. 7-12):

* a) *From 1PI functions with one ω and one A (fig. 5)*

$$\begin{aligned} \beta_1^{ab} &= -(-i)^3 \times \text{coef. in r.s.p. } \overline{\langle \omega^a(p_2 \equiv -p_1) A_\mu^b(p_1) N[\check{\Delta}](q=0) \rangle}_{K=0}^{1\text{PI}} \text{ of } \bar{p}_1^2 \bar{p}_{1\mu} \\ &= -\frac{1}{48} \frac{(-i)^3 \partial^3}{\partial \bar{p}_1^\nu \partial \bar{p}_{1\nu} \partial \bar{p}_{1\mu}} \text{r.s.p. } \overline{\langle \omega^a(-p_1) A_\mu^b(p_1) N[\check{\Delta}](q=0) \rangle}_{K=0}^{1\text{PI}} \Big|_{p_i \equiv 0} \\ &= -\frac{1}{(4\pi)^2} \frac{T_L + T_R}{3} \delta^{ab} \hbar^1 + O(\hbar^2), \end{aligned} \quad (50)$$




Figure 5: 1PI Feynman diagrams needed to compute eq. (50)

* b) *From 1PI functions with one ω and two A s, and no $\varepsilon_{\mu\nu\alpha\beta}$ tensor (fig. 6)*

$$\begin{aligned} \{\beta_2^{abc}, \beta_4^{abc}\} &= \\ &= -(-i)^2 \times \text{coef. in r.s.p. } \overline{\langle \omega^a(p_3 \equiv -p_1 - p_2) A_\mu^b(p_1) A_\nu^c(p_2) N[\check{\Delta}](q=0) \rangle}_{K=0}^{1\text{PI}} \text{ of } \\ &\quad \{\bar{p}_1^2 \bar{g}_{\mu\nu}, \bar{p}_{1\mu} \bar{p}_{1\nu}\} \text{ respectively} \\ &= \frac{1}{(4\pi)^2} (T_L + T_R) c^{abc} \left\{ -\frac{1}{3}, \frac{2}{3} \right\} \hbar^1 + O(\hbar^2), \end{aligned} \quad (51)$$

$$\begin{aligned} \{\beta_3^{abc} = \beta_3^{acb}, \beta_5^{abc} = \beta_5^{acb}, \beta_6^{abc} = \beta_6^{acb}\} &= \\ &= -\frac{(-i)^2}{2} \times \text{coef. in r.s.p. } \overline{\langle \omega^a(p_3 \equiv -p_1 - p_2) A_\mu^b(p_1) A_\nu^c(p_2) N[\check{\Delta}](q=0) \rangle}_{K=0}^{1\text{PI}} \text{ of } \\ &\quad \{\bar{p}_1 \cdot \bar{p}_2 \bar{g}_{\mu\nu}, \bar{p}_{1\mu} \bar{p}_{2\nu}, \bar{p}_{1\nu} \bar{p}_{2\mu}\} \text{ resp.} \\ &= 0 \hbar^1 + O(\hbar^2), \end{aligned} \quad (52)$$




Figure 6: 1PI Feynman diagrams needed to compute eqs. (51), (52) and (61)

* c) From 1PI functions with one ω and three As , and no $\varepsilon_{\mu\nu\alpha\beta}$ tensor (fig. 7)

$$\begin{aligned}
\beta_7^{abcd} &= \beta_7^{abdc} = \\
&= -\frac{(-i)}{2} \times \text{coef. in r.s.p. } \overline{\langle \omega^a(-\sum_i^3 p_i) A_\mu^b(p_1) A_\nu^c(p_2) A_\rho^d(p_3) N[\check{\Delta}](q=0) \rangle}_{K=0}^{1\text{PI}} \text{ of} \\
&\quad \bar{p}_{1\mu} \bar{g}_{\nu\rho} \\
&= \frac{1}{(4\pi)^2} \frac{1}{6} [T_{L+R}^{abcd} + T_{L+R}^{acdb} + T_{L+R}^{abdc} + T_{L+R}^{adcb} - T_{L+R}^{acbd} - T_{L+R}^{adbc}] \hbar^1 + O(\hbar^2), \tag{53}
\end{aligned}$$

$$\begin{aligned}
\beta_8^{abcd} &= -(-i) \times \text{coef. in r.s.p. } \overline{\langle \omega^a(-\sum_i^3 p_i) A_\mu^b(p_1) A_\nu^c(p_2) A_\rho^d(p_3) N[\check{\Delta}](q=0) \rangle}_{K=0}^{1\text{PI}} \text{ of} \\
&\quad (\bar{p}_{1\nu} \bar{g}_{\mu\rho} + \bar{p}_{1\rho} \bar{g}_{\mu\nu}) \\
&= \frac{1}{(4\pi)^2} \frac{1}{3} [T_{L+R}^{abcd} + T_{L+R}^{acbd} + T_{L+R}^{acdb} + T_{L+R}^{adbc} - T_{L+R}^{abdc} - T_{L+R}^{adcb}] \hbar^1 + O(\hbar^2), \tag{54}
\end{aligned}$$




Figure 7: 1PI Feynman diagrams needed to compute eqs. (53), (54) and (62)




Figure 8: 1PI Feynman diagrams needed to compute eqs. (55) and (63)

* d) From 1PI functions with one ω and four A s, and no $\varepsilon_{\mu\nu\alpha\beta}$ tensor (fig. 8)

$$\begin{aligned}
\beta_9^{abcde} &= \beta_9^{a\{bc\}\{de\}} = \beta_9^{a\{de\}\{bc\}} = \\
&= -\frac{1}{8} \times \text{coef. in r.s.p. } \overline{\langle \omega^a A_\mu^b A_\nu^c A_\rho^d A_\lambda^e N[\check{\Delta}](q=0) \rangle}_{K=0}^{1\text{PI}} \text{ of } \bar{g}_{\mu\nu} \\
&= 0 \hbar^1 + O(\hbar^2),
\end{aligned} \tag{55}$$

In principle, due to the presence of γ_5 , there should be also evanescent normal products in the expansion of the anomalous insertion $N[\hat{\Delta}] \cdot \Gamma_{\text{ren}}$. Indeed, the coefficient of $\int N[\omega^a \hat{\square} \bar{\partial}_\mu A^{b\mu}]$ is

$$\begin{aligned}
&- (-i)^3 \times \text{coef. in r.s.p. } \overline{\langle \omega^a(p_2 \equiv -p_1) A_\mu^b(p_1) N[\check{\Delta}](q=0) \rangle}_{K=0}^{1\text{PI}} \text{ of } \hat{p}_1^2 \bar{p}_{1\mu} \\
&= -\frac{1}{8(d-4)} \frac{(-i)^3 \partial^3}{\partial \bar{p}_1^\nu \partial \hat{p}_{1\nu} \partial \bar{p}_{1\mu}} \text{ r.s.p. } \overline{\langle \omega^a(-p_1) A_\mu^b(p_1) N[\check{\Delta}](q=0) \rangle}_{K=0}^{1\text{PI}} \Big|_{p_i \equiv 0} \\
&= -\frac{1}{(4\pi)^2} \frac{T_L + T_R}{3} \delta^{ab} \hbar^1 + O(\hbar^2),
\end{aligned}$$

However, a simplification occur, for the coefficient of $\int N[\omega^a \bar{\square} \hat{\partial}_\mu A^{b\mu}]$ is

$$\begin{aligned}
&- (-i)^3 \times \text{coef. in r.s.p. } \overline{\langle \omega^a(p_2 \equiv -p_1) A_\mu^b(p_1) N[\check{\Delta}](q=0) \rangle}_{K=0}^{1\text{PI}} \text{ of } \bar{p}_1^2 \hat{p}_{1\mu} \\
&= -\frac{1}{8(d-4)} \frac{(-i)^3 \partial^3}{\partial \bar{p}_1^\nu \partial \bar{p}_{1\nu} \partial \hat{p}_{1\mu}} \text{ r.s.p. } \overline{\langle \omega^a(-p_1) A_\mu^b(p_1) N[\check{\Delta}](q=0) \rangle}_{K=0}^{1\text{PI}} \Big|_{p_i \equiv 0} \\
&= 0 \hbar^1 + O(\hbar^2),
\end{aligned}$$

and so on. Due to our expansion in barred and hatted objects rather than in standard and hatted ones, and because of the form of the regularized interaction it turns out that *the rest of anomalous coefficients are 0 in the one-loop approximation*. Nevertheless, as we know

$$\begin{aligned}
& (T_{L(R)}^e T_{L(R)}^a T_{L(R)}^e)_{ij} = \\
& = (C_{L(R)} - \frac{C_A}{2}) (T_{L(R)}^a)_{ij} \\
& c_{bac} (T_{L(R)}^{[c} T_{L(R)}^{b]})_{ij} = \\
& = \frac{i}{2} c_{acb} c_{cbe} (T_{L(R)}^e)_{ij} = \\
& = \frac{i}{2} C_A (T_{L(R)}^a)_{ij} \\
& c_{bac} (T_{L(R)}^{[b} T_{L(R)}^{c]})_{ij} = \\
& = -\frac{i}{2} c_{abc} c_{bce} (T_{L(R)}^e)_{ij} = \\
& = -\frac{i}{2} C_A (T_{L(R)}^a)_{ij}
\end{aligned}$$

Figure 9: 1PI Feynman diagrams needed to compute eqs. (56) and (57). (Notice that the gauge group structure is explicitly shown under these and subsequent diagrams)

from subsection 2.4., these anomalous coefficients only matter at the next perturbative order, so this vanishing of the coefficients is only a simplification relevant to higher order computations.

* e) From 1PI functions with one ω and one $\psi, \bar{\psi}$ ($\psi', \bar{\psi}'$) pair (fig. 9)

$$\begin{aligned}
\beta_{10L(R)}^{a,ij} &= -(-i) \times \text{coef. in r.s.p.} \overline{\langle \omega^a(p_3 \equiv -p_1 - p_2) \psi_{\beta j}^{(i)}(p_2) \bar{\psi}_{\alpha i}^{(j)}(p_1) N[\check{\Delta}](q=0) \rangle}_{K=0}^{1\text{PI}} \text{ of} \\
& (\bar{p}_2 P_{L(R)})_{\alpha\beta} \\
&= -\frac{1}{4 \cdot 2} \frac{(-i) \partial}{\partial \bar{p}_2^\mu} \text{r.s.p.} \overline{\langle \omega^a(-p_1 - p_2) \psi_{\beta j}^{(i)}(p_2) \bar{\psi}_{\alpha i}^{(j)}(p_1) N[\check{\Delta}](q=0) \rangle}_{K=0}^{1\text{PI}} \Big|_{p \equiv 0} \cdot (P_{L(R)} \bar{\gamma}^\mu)_{\beta\alpha} \\
&= \frac{i}{4 \cdot 2} \text{Tr} \left[\frac{\partial}{\partial \bar{p}_2^\mu} \text{r.s.p.} \overline{\langle \omega^a(-p_1 - p_2) \psi_j^{(i)}(p_2) \bar{\psi}_i^{(j)}(p_1) N[\check{\Delta}](q=0) \rangle}_{K=0}^{1\text{PI}} \Big|_{p_i \equiv 0} P_{L(R)} \bar{\gamma}^\mu \right] \\
&= -\frac{1}{(4\pi)^2} g^2 \left\{ (C_{L(R)} - \frac{C_A}{4}) [T_{L(R)}^a]^{ij} + (\alpha' - 1) \left(\frac{C_{L(R)}}{6} - \frac{C_A}{4} \right) [T_{L(R)}^a]^{ij} \right\} \hbar^1 \\
&+ O(\hbar^2). \tag{56}
\end{aligned}$$

$$\begin{aligned}
\beta_{11L(R)}^{a,ij} &= -(-i) \times \text{coef. in r.s.p.} \overline{\langle \omega^a(p_3 \equiv -p_1 - p_2) \psi_{\beta j}^{(i)}(p_2) \bar{\psi}_{\alpha i}^{(j)}(p_1) N[\check{\Delta}](q=0) \rangle}_{K=0}^{1\text{PI}} \text{ of} \\
& (\bar{p}_1 P_{L(R)})_{\alpha\beta} \\
&= \frac{i}{4 \cdot 2} \text{Tr} \left[\frac{\partial}{\partial \bar{p}_1^\mu} \text{r.s.p.} \overline{\langle \omega^a(-p_1 - p_2) \psi_j^{(i)}(p_2) \bar{\psi}_i^{(j)}(p_1) N[\check{\Delta}](q=0) \rangle}_{K=0}^{1\text{PI}} \Big|_{p_i \equiv 0} P_{L(R)} \bar{\gamma}^\mu \right] \\
&= -\frac{1}{(4\pi)^2} g^2 \left\{ (C_{L(R)} - \frac{C_A}{4}) [T_{L(R)}^a]^{ij} + (\alpha' - 1) \left(\frac{C_{L(R)}}{6} - \frac{C_A}{4} \right) [T_{L(R)}^a]^{ij} \right\} \hbar^1 \\
&+ O(\hbar^2). \tag{57}
\end{aligned}$$




Figure 10: 1PI Feynman diagrams needed to compute eq. (58)

* f) From 1PI functions with one ω , one A and one $\psi, \bar{\psi}$ ($\psi', \bar{\psi}'$) pair (fig. 10)

$$\begin{aligned}
 \beta_{12L(R)}^{ab,ij} &= -1 \times \text{coef. in r.s.p. } \overline{\langle \omega^a A_\mu^b \psi_{\beta j}^{(\prime)} \bar{\psi}_{\alpha i}^{(\prime)} N[\check{\Delta}](q=0) \rangle}_{K=0}^{1\text{PI}} \text{ of} \\
 &\quad (\bar{\gamma}^\mu P_{L(R)})_{\alpha\beta} \\
 &= -\frac{1}{4 \cdot 2} \text{Tr} \left[\text{r.s.p. } \overline{\langle \omega^a A_\mu^b \psi_j^{(\prime)} \bar{\psi}_i^{(\prime)} N[\check{\Delta}](q=0) \rangle}_{K=0}^{1\text{PI}} \Big|_{p_i \equiv 0} P_{L(R)} \bar{\gamma}^\mu \right] \\
 &= -\frac{1}{(4\pi)^2} g^2 \frac{1 + (\alpha' - 1)}{4} C_A i [T_{L(R)}^a, T_{L(R)}^b]_{ij} \hbar^1 + O(\hbar^2). \tag{58}
 \end{aligned}$$

Notice that the gauge group structure of the first five diagrams in fig. 10 involves three or more gauge group generator matrices. The gauge group structure of each diagram is rather involved but it turns out that the r.s.p. of each of the first three diagrams vanishes, that the r.s.p. of the fourth cancels exactly against the fifth, and, thanks to the Jacobi identity, that the result of the last four diagrams combine in pairs to give eq. (58), where the gauge group structure is simply a commutator. This is remarkable, because we shall see in Appendix C that the difference between the one-loop 1PI fermionic vertex computed *à la* Breitenhoner and Maison and the same 1PI function evaluated with the “naive”

prescription involves only a single gauge group generator and, hence, it is sensible to expect that the finite counterterm that is needed to restore at the one-loop level the BRS symmetry in the Breitenlöhner and Maison formalism involved only a generator. The BRS variation, b , of this counterterm yields a contribution linear in A_μ^b which will match the gauge group structure of the symmetry breaking diagrams in fig. 10 thanks to the various cancellations and simplifications just described.




Figure 11: 1PI Feynman diagrams needed to compute eq. (59)

* g) From 1PI functions with fermions and external fields (figs. 11-12)

$$\begin{aligned}
 \beta_{14L(R)}^{ab,ij} &= \frac{i}{2\hbar} \times \text{coef. in r.s.p.} \overline{\langle \omega^a \omega^b \psi_{\beta j}^{(\prime)} N[s\psi_{\alpha i}^{(\prime)}] N[\check{\Delta}](q=0) \rangle}_{K=0}^{1\text{PI}} \text{ of} \\
 &\quad (P_{L(R)})_{\alpha\beta} \\
 &= \frac{1}{(4\pi)^2} g^2 [1 + (\alpha' - 1)] \frac{C_A}{8} [T_{L(R)}^a, T_{L(R)}^b]_{ij} \hbar^1 + O(\hbar^2). \tag{59}
 \end{aligned}$$

$$\begin{aligned}
 \beta_{15L(R)}^{ab,ij} &= \frac{i}{2\hbar} \times \text{coef. in r.s.p.} \overline{\langle \omega^a \omega^b \bar{\psi}_{\beta j}^{(\prime)} N[s\bar{\psi}_{\alpha i}^{(\prime)}] N[\check{\Delta}](q=0) \rangle}_{K=0}^{1\text{PI}} \text{ of} \\
 &\quad (P_{R(L)})_{\alpha\beta} \\
 &= -\frac{1}{(4\pi)^2} g^2 [1 + (\alpha' - 1)] \frac{C_A}{8} [T_{L(R)}^a, T_{L(R)}^b]_{ij} \hbar^1 + O(\hbar^2). \tag{60}
 \end{aligned}$$

As in case f), it seems impossible that the gauge group structure of the diagrams be matched by gauge group structure of a simple one-loop finite counterterm. But the last pair of diagrams of each 1PI function vanishes, and the other pair of antisymmetric diagrams conspire with the help of the Jacobi identity to give a simple commutator as result.




Figure 12: 1PI Feynman diagrams needed to compute eq. (60)

* h) From 1PI functions with $\varepsilon_{\mu\nu\alpha\beta}$ (figs. 6-8)

These coefficients are very important, because if they are non-zero, give monomials in the expansion of the breaking that can not be cancelled by finite counterterms; *i.e.* they give the essential non-Abelian chiral anomaly.

$$\begin{aligned}
\beta_{50}^{abc} &= \beta_{50}^{acb} = \\
&= -\frac{(-i)^2}{2} \times \text{coef. in r.s.p. } \overline{\langle \omega^a(p_3 \equiv -p_1 - p_2) A_\mu^b(p_1) A_\nu^c(p_2) N[\check{\Delta}](q=0) \rangle}_{K=0}^{1\text{PI}} \text{ of} \\
&\quad \varepsilon_{\mu\nu\alpha\beta} p_1^\alpha p_2^\beta \\
&= -\frac{1}{(4\pi)^2} \frac{1}{3} d_{L-R}^{abc} \hbar^1 + O(\hbar^2),
\end{aligned} \tag{61}$$

$$\begin{aligned}
\beta_{51}^{abcd} &= -\beta_{51}^{abdc} = \\
&= -\frac{(-i)}{2} \times \text{coef. in r.s.p. } \overline{\langle \omega^a(p_4 \equiv -\sum_i^3 p_i) A_\mu^b(p_1) A_\nu^c A_\rho^d N[\check{\Delta}](q=0) \rangle}_{K=0}^{1\text{PI}} \text{ of} \\
&\quad \varepsilon_{\mu\nu\rho\alpha} \bar{p}_1^\alpha \\
&= -\frac{1}{(4\pi)^2} \frac{1}{6} \mathcal{D}_{L-R}^{abcd} \hbar^1 + O(\hbar^2),
\end{aligned} \tag{62}$$

$$\begin{aligned}
\beta_{52}^{abcde} &= \beta_{52}^{a[bcde]} = -\frac{1}{4!} \times \text{coef. in r.s.p. } \overline{\langle \omega^a A_\mu^b A_\nu^c A_\rho^d A_\lambda^e N[\check{\Delta}](q=0) \rangle}_{K=0}^{1\text{PI}} \text{ of} \\
&\quad \varepsilon_{\mu\nu\rho\lambda} \\
&= O(\hbar^2),
\end{aligned} \tag{63}$$

with the definitions

$$\begin{aligned}
d_L^{abc} &= \text{Tr} [T_L^a \{T_L^b, T_L^c\}] = d_L^{\{abc\}} \\
\mathcal{D}_L^{abcd} &= -i 3! \text{Tr} [T_L^a T_L^b T_L^c T_L^d] = \frac{1}{2} (d_L^{abe} c^{ecd} + d_L^{ace} c^{edb} + d_L^{ade} c^{ebc});
\end{aligned}$$

same for d_R^{abc} and $\mathcal{D}_R^{abcd}$, and $d_{L-R} \equiv d_L - d_R$, $\mathcal{D}_{L-R} \equiv \mathcal{D}_L - \mathcal{D}_R$.

Of course, we write $O(\hbar^2)$ because we have computed at first order in $\hbar$; but due to the Adler-Bardeem theorem for non-Abelian gauge theories [18], all these coefficient should be zero at higher orders if there would not be an anomaly at first order.

Our one-loop computation of these coefficients give the *non-Abelian (essential) chiral gauge anomaly*:

$$\begin{aligned}
& \int d^4x \{ \beta_{50}^{abc} N[\varepsilon_{\mu\nu\alpha\beta} \omega^a (\partial^\alpha A^{\mu b})(\partial^\beta A^{\nu c})] + \beta_{51}^{abcd} N[\varepsilon_{\mu\nu\alpha\beta} \omega^a (\partial^\alpha A^{\mu b}) A^{\nu c} A^{\rho d}] + \\
& \quad + \beta_{52}^{abcde} N[\varepsilon_{\mu\nu\alpha\beta} \omega^a A^{\mu b} A^{\nu c} A^{\rho d} A^{\lambda e}] \} \cdot \Gamma_{\text{ren}} = \\
& = -\frac{1}{(4\pi)^2} \frac{1}{3} \left\{ \varepsilon_{\mu\nu\alpha\beta} d_{L-R}^{abc} \omega^a (\partial^\alpha A^{\mu b})(\partial^\beta A^{\nu c}) + \frac{1}{2} \varepsilon_{\mu\nu\rho\alpha} \mathcal{D}_{L-R}^{abcd} \omega^a (\partial^\alpha A^{\mu b}) A^{\nu c} A^{\rho d} \right\} \\
& \quad + O(\hbar^2) = \\
& = \frac{1}{(4\pi)^2} \frac{1}{3} \varepsilon_{\mu\nu\rho\sigma} \omega^a \partial^\mu \left\{ d_{L-R}^{abc} (\partial^\nu A^{\rho b}) A^{\sigma c} + \frac{1}{6} \mathcal{D}_{L-R}^{abcd} A^{\nu b} A^{\rho c} A^{\sigma d} \right\} + O(\hbar^2) = \\
& = \frac{1}{(4\pi)^2} \frac{2}{3} \varepsilon_{\mu\nu\rho\sigma} \text{Tr}_{L-R} \left\{ \omega \partial^\mu (\partial^\nu A^\rho A^\sigma - \frac{i}{2} A^\nu A^\rho A^\sigma) \right\} + O(\hbar^2)
\end{aligned}$$

Remember that in the formulae for the coefficients, a sum over the different left or right representations is understood. Therefore, the matter representation may be chosen such that the coefficients of the essential anomaly above vanish.

Finally, as explained at the end of section 2.4., the coefficients k_i of the true expansion in the quantum (standard) basis of insertions are equal to the coefficients β_i in the one-loop approximation. Moreover $\bar{k}_i^{(1)} = k_i^{(1)}$.

* *The expression of the breaking at order $\hbar$*

Putting it all together we get the complete form of the integrated breaking, $\left[N[\hat{\Delta}] \cdot \Gamma_{\text{ren}} \right]^{(1)}$, at order $\hbar$ (this formula is a 4 dimensional one !), which reads

$$\begin{aligned}
\left[N[\hat{\Delta}] \cdot \Gamma_{\text{ren}} \right]^{(1)} = & -\frac{1}{(4\pi)^2} \frac{T_L + T_R}{3} \delta^{ab} \mathcal{M}_1^{ab} \hbar^1 + \\
& -\frac{1}{(4\pi)^2} \frac{T_L + T_R}{3} c^{abc} \mathcal{M}_2^{abc} \hbar^1 + \frac{1}{(4\pi)^2} (T_L + T_R) \frac{2}{3} c^{abc} \mathcal{M}_4^{abc} \hbar^1 + \\
& + \frac{1}{(4\pi)^2} \frac{1}{6} [T_{L+R}^{acbd} + T_{L+R}^{adbc} - (T_L + T_R) (c^{ebc} c^{eda} + c^{ebd} c^{eca})] \mathcal{M}_7^{abcd} \hbar^1 \\
& + \frac{1}{(4\pi)^2} \frac{1}{3} [T_{L+R}^{abcd} + T_{L+R}^{adcb} + (T_L + T_R) (c^{ebc} c^{eda} + c^{eba} c^{edc})] \mathcal{M}_8^{abcd} \hbar^1 \\
& - \frac{1}{(4\pi)^2} g^2 \left[C_L - \frac{C_A}{4} + (\alpha' - 1) \left(\frac{C_L}{6} - \frac{C_A}{4} \right) \right] (\mathcal{M}_{10L} + \mathcal{M}_{11L}) \hbar^1 \\
& - \frac{1}{(4\pi)^2} g^2 \left[C_R - \frac{C_A}{4} + (\alpha' - 1) \left(\frac{C_R}{6} - \frac{C_A}{4} \right) \right] (\mathcal{M}_{10R} + \mathcal{M}_{11R}) \hbar^1
\end{aligned}$$

$$\begin{aligned}
& - \frac{i}{(4\pi)^2} g^2 \frac{1 + (\alpha' - 1)}{4} C_A (\mathcal{M}_{12L} - \mathcal{M}_{13L} + \mathcal{M}_{12R} - \mathcal{M}_{13R}) \hbar^1 \\
& + \frac{i}{(4\pi)^2} g^2 \frac{1 + (\alpha' - 1)}{4} \frac{C_A}{8} c^{abc} (\mathcal{M}_{14L}^{abc} - \mathcal{M}_{15L}^{abc} + \mathcal{M}_{14R}^{abc} - \mathcal{M}_{15R}^{abc}) \hbar^1 \\
& + \frac{1}{(4\pi)^2} \frac{2}{3} \int d^4x \varepsilon_{\mu\nu\rho\sigma} \text{Tr}_{L-R} [\omega \partial^\mu (\partial^\nu A^\rho A^\sigma - \frac{i}{2} A^\nu A^\rho A^\sigma)] \hbar^1, \quad (64)
\end{aligned}$$

where we have defined

$$\begin{aligned}
\mathcal{M}_{10L(R)} &= (T_{L(R)}^a)^{ij} \mathcal{M}_{10L(R)}^{a,ij} = \int d^4x \omega^a \bar{\psi}^{(\prime)} \bar{\gamma}^\mu P_{L(R)} T_{L(R)}^a \partial_\mu \psi^{(\prime)} \\
\mathcal{M}_{11L(R)} &= (T_{L(R)}^a)^{ij} \mathcal{M}_{11L(R)}^{a,ij} = \int d^4x \omega^a (\partial_\mu \bar{\psi}^{(\prime)}) \bar{\gamma}^\mu P_{L(R)} T_{L(R)}^a \psi^{(\prime)} \\
\mathcal{M}_{12L(R)} &= (T_{L(R)}^a T_{L(R)}^b)^{ij} \mathcal{M}_{12L(R)}^{ab,ij} = \int d^4x \omega^a \bar{\psi}^{(\prime)} \bar{\gamma}^\mu P_{L(R)} T_{L(R)}^a T_{L(R)}^b \psi^{(\prime)} A_\mu^b \\
\mathcal{M}_{13L(R)} &= (T_{L(R)}^b T_{L(R)}^a)^{ij} \mathcal{M}_{13L(R)}^{ab,ij} = \int d^4x \omega^a \bar{\psi}^{(\prime)} \bar{\gamma}^\mu P_{L(R)} T_{L(R)}^b T_{L(R)}^a \psi^{(\prime)} A_\mu^b \\
\mathcal{M}_{14L(R)}^{abc} &= (T_{L(R)}^c)^{ij} \mathcal{M}_{14L(R)}^{ab,ij} = \int d^4x \omega^a \omega^b \bar{L}(\bar{R}) P_{L(R)} T_{L(R)}^c \psi^{(\prime)}, \\
\mathcal{M}_{15L(R)}^{abc} &= (T_{L(R)}^c)^{ij} \mathcal{M}_{15L(R)}^{ab,ij} = \int d^4x \omega^a \omega^b \bar{\psi}^{(\prime)} P_{R(L)} T_{L(R)}^c L(R),
\end{aligned}$$

and we have used the following equation:

$$-2T_{L+R}^{abcd} - 2T_{L+R}^{adcb} + T_{L+R}^{abdc} + T_{L+R}^{acbd} + T_{L+R}^{acdb} + T_{L+R}^{adbc} = (T_L + T_R) (c^{ebc} c^{eda} + c^{eba} c^{edc}). \quad (65)$$

Notice that if the classical theory were CP invariant, the combination of monomials obtained have the same CP definite value as the breaking, as we would have expected from the fact that Dimensional Regularization respects discrete symmetries. But we have not assumed any discrete symmetry property when writing the basis of Bonneau monomials because we work with general representations.

The result in eq. (64) can be obtained by computing the left hand side of eq. (64) to the one-loop order, *i.e.* calculating the renormalized part of all possible one-loop 1PI diagrams with the insertion $\hat{\Delta}$; their finite part has to be a local term, because when going to the 4-dimensional space all $(d-4)$ -objects are set to be 0, so the only remaining finite part of such a diagram must come from a polynomial singular part formed by a 4-dimensional object and a contracted $(d-4)$ -object which cancel the singularity in $d-4$. Therefore, being local this one-loop renormalized 1PI functions, there are operators (in the 4-dimensional space) whose Feynman rules attached with a $\hbar^1$ give the same result. The sum of this operators is precisely the right hand side of eq. (64).

In fact, the one-loop calculation of the coefficients of the Bonneau identities via r.s.p. of diagrams with the insertion of $\tilde{\Delta}$ are exactly the same as the finite part of diagrams with the insertion of $\hat{\Delta}$, excepting some factors in the Bonneau coefficients which directly give the operators whose Feynman rules lead to the same result.

But the Bonneau identities unravel the problem in higher order computations, and the formulae of the coefficients given above would be the same (although with new Feynman rules coming from the finite counterterms we would have added in the previous recursive step).

3.6. Restoring the BRS symmetry: the computation of $S_{\text{fct}}^{(1)}$

We know from algebraic renormalization theory and BRS cohomology [18] that there exists an integrated local functional of the fields and their derivatives, let us call it $X^{(1)}$, such that the anomalous contribution $\left[N[\hat{\Delta}] \cdot \Gamma_{\text{ren}} \right]^{(1)}$, given in eq. (64), can be cast into the form

$$\left[N[\hat{\Delta}] \cdot \Gamma_{\text{ren}} \right]^{(1)} = \frac{1}{(4\pi)^2} \frac{2}{3} \int d^4x \varepsilon_{\mu\nu\rho\sigma} \text{Tr}_{\text{L-R}} \left[\omega \partial^\mu (\partial^\nu A^\rho A^\sigma - \frac{i}{2} A^\nu A^\rho A^\sigma) \right] \hbar^1 + bX^{(1)}.$$

The integrated local functional, $X^{(1)}$, has ghost number zero and ultraviolet dimension 4. Hence, by choosing $S_{\text{fct}}^{(1)}$ in eq. (35) so that it verifies

$$S_{\text{fct}}^{(1)} = -X^{(1)},$$

and recalling that eqs. (44) and (42) hold, we conclude that the BRS identity is broken at order $\hbar$ by the essential non-Abelian gauge anomaly only:

$$\mathcal{S}(\Gamma_{\text{ren}}) = \frac{1}{(4\pi)^2} \frac{2}{3} \int d^4x \varepsilon_{\mu\nu\rho\sigma} \text{Tr}_{\text{L-R}} \left[\omega \partial^\mu (\partial^\nu A^\rho A^\sigma - \frac{i}{2} A^\nu A^\rho A^\sigma) \right] \hbar^1 + O(\hbar^2).$$

The previous equation leads in turn to the result we sought for; namely, that if the fermion representations meet the anomaly cancellation criterion [23], the BRS symmetry can be restored (up to order $\hbar$) by an appropriate choice of finite counterterms $S_{\text{fct}}^{(1)}$.

Our next task will be to compute $S_{\text{fct}}^{(1)} = -X^{(1)}$. We shall demand that $S_{\text{fct}}^{(1)}$ do not depend neither on B nor $\tilde{\rho}$. Recall that these fields do not occur in eq. (64).

The more general approach to this computation, similar to our study of the expansion of $\left[N[\hat{\Delta}] \cdot \Gamma_{\text{ren}} \right]^{(1)}$, would be: 1. Write the complete list of possible monomials of ultraviolet dimension 4, ghost number 0 and with no free Lorentz indices $\dagger$; 2. Write the general expression of the finite counterterms as a linear combination of those monomials with coefficients depending on gauge group indices. 3. Take their b -variation and finally 4. impose that this variation cancels the expansion of the breaking term $\left[N[\hat{\Delta}] \cdot \Gamma_{\text{ren}} \right]^{(1)}$,

$\dagger$ This list is a list in the four dimensional space-time. Of course, some generalization of them to the d -dimensional space-time of Dimensional Regularization has to be chosen while writing them in the Dimensional Regularization action; but the differences, being $(d-4)$ -objects, will begin to produce any effect at the following order in $\hbar$ (because they require loops to yield a non-zero contribution and there is already an explicit $\hbar$ in $S_{\text{fct}}^{(1)}$)

excepting the essential non-Abelian anomaly. This should lead to a compatible indeterminate system with the coefficients as variables and with a degree of indeterminacy exactly equal to the number of b -invariant operators in the classical theory.

But, writing such list and taking its b variation is a tedious task: lots of indices would come in and the resolution of the final system would appear a bit messy.

It is plain that the gauge group structure of the finite counterterms which would be needed in the one-loop computation to restore the BRS symmetry is not arbitrary. A natural ansatz for the gauge group structure of these counterterms is to take the one-loop gauge group structure which can appear while computing the Feynman diagrams. Therefore, we write

$$\begin{aligned}
S_{\text{fct}}^{(1)} = & a_1 \int (\partial_\mu A^\mu)^2 + a_2 \int A_\mu \square A^\mu + c \int i c^{abc} (\partial_\mu A_\nu^a) A^{b\mu} A^{c\nu} \\
& + (d_L d_L^{abc} + d_R d_R^{abc}) \int (\partial_\mu A_\nu^a) A^{b\mu} A^{c\nu} + (f_L T_L^{abcd} + f_R T_R^{abcd}) \int A^{a\mu} A_\mu^b A^{c\nu} A_\nu^d \\
& + e c^{abc} c^{ade} \int A^{b\mu} A^{c\nu} A_\mu^d A_\nu^e + n_L \int \frac{i}{2} \bar{\psi} \gamma^\mu P_L \overset{\leftrightarrow}{\partial}_\mu \psi + n_R \int \frac{i}{2} \bar{\psi}' \gamma^\mu P_R \overset{\leftrightarrow}{\partial}_\mu \psi' \\
& + p_L \int \bar{\psi} \gamma^\mu P_L T_L^a \psi A_\mu^a + p_R \int \bar{\psi}' \gamma^\mu P_R T_R^a \psi' A_\mu^a \\
& + \int [u_{1L} \bar{L} s \psi + u_{1R} \bar{R} s \psi' + u_{2L} L s \bar{\psi} + u_{2R} R s \bar{\psi}'] \quad (66)
\end{aligned}$$

Notice that because of the reason given above we have not written terms with ρ or $\bar{\omega}$.

Moreover, we have the combinations of terms given in eq. (30) which are b -invariant. Obviously, these combinations are not relevant to our computation. Therefore, we can reduce the basis of possible finite counterterms to a minimum number taking into account these invariant counterterms.

L_A and L_ω contain $\tilde{\rho}^\mu$, which we supposed already not to appear in the finite counterterms, so these invariant terms can not be used to reduce more the number of finite counterterms. But, using L_g we can impose $e = 0$ and by taking advantage of $L_{\psi^{(\prime)}}$ we can set $p_{L(R)} = 0$, without loss of generality.

Then, there should be a *unique* solution for the problem of cancellation of spurious anomalies in terms of the rest of variables $a_1, a_2, c, d_{L(R)}, f_{L(R)}, n_{L(R)}, u_{1L(R)}$ and $u_{2L(R)}$.

We recast eq. (66) in the following form:

$$\begin{aligned}
S_{\text{fct}}^{(1)} = & a_1 \tilde{\Delta}_1 + a_2 \tilde{\Delta}_2 + c \tilde{\Delta}_3 + d_L \tilde{\Delta}_{4L} + d_R \tilde{\Delta}_{4R} + f_L \tilde{\Delta}_{5L} + f_R \tilde{\Delta}_{5R} \\
& + n_L \tilde{\Delta}_{6L} + n_R \tilde{\Delta}_{6R} + u_{1L} \tilde{\Delta}_{7L} + u_{1R} \tilde{\Delta}_{7R} + u_{2L} \tilde{\Delta}_{8L} + u_{2R} \tilde{\Delta}_{8R}. \quad (67)
\end{aligned}$$

We need the b variations of the $\tilde{\Delta}$ s expanded in terms of the integrated monomials we called $\mathcal{M}$ s. Moreover, the coefficients of these expansions should be symmetrized if the gauge group indices of the corresponding monomials have some symmetry properties.

This is necessary because, for example, $\mathcal{M}_3^{abc} = \mathcal{M}_3^{acb}$ and then not all $\mathcal{M}_3^{abc}$ are linearly independent. The result reads :

$$\begin{aligned}
b \tilde{\Delta}_1 &= 2 \delta^{ab} \mathcal{M}_1^{ab} - 2 c^{abc} \mathcal{M}_4^{abc}, & b \tilde{\Delta}_2 &= -2 \delta^{ab} \mathcal{M}_1^{ab} + 2 c^{abc} \mathcal{M}_2^{abc}, \\
b \tilde{\Delta}_3 &= -i c^{abc} \mathcal{M}_4^{abc} + i c^{abc} \mathcal{M}_2^{abc} \\
&\quad - \frac{i}{2} [c^{ebc} c^{eda} + c^{ebd} c^{eca}] \mathcal{M}_7^{abcd} + i [c^{ebc} c^{eda} + c^{eba} c^{edc}] \mathcal{M}_8^{abcd}, \\
b \tilde{\Delta}_{4L(R)} &= -d_{L(R)}^{abc} \mathcal{M}_2^{abc} - d_{L(R)}^{abc} \mathcal{M}_3^{abc} + d_{L(R)}^{abc} \mathcal{M}_4^{abc} + d_{L(R)}^{abc} \mathcal{M}_5^{abc} \\
&\quad - (d_{L(R)}^{ebc} c^{eda} + d_{L(R)}^{ebd} c^{eca}) \mathcal{M}_7^{abcd} + d_{L(R)}^{ebd} c^{eca} \mathcal{M}_8^{abcd}, \\
b \tilde{\Delta}_{5L(R)} &= - [T_{L(R)}^{abcd} + T_{L(R)}^{acdb} + T_{L(R)}^{abdc} + T_{L(R)}^{adcb}] \mathcal{M}_7^{abcd} \\
&\quad - 2 [T_{L(R)}^{acbd} + T_{L(R)}^{abdc} + T_{L(R)}^{acdb} + T_{L(R)}^{adbc}] \mathcal{M}_8^{abcd} \\
&\quad - \frac{1}{4} [(T_{L(R)}^{mcde} + T_{L(R)}^{mdce} + T_{L(R)}^{mced} + T_{L(R)}^{medc}) c^{mab} \\
&\quad + (T_{L(R)}^{mbde} + T_{L(R)}^{mdbe} + T_{L(R)}^{mbed} + T_{L(R)}^{medb}) c^{mac} \\
&\quad + (T_{L(R)}^{mebc} + T_{L(R)}^{mbce} + T_{L(R)}^{mecb} + T_{L(R)}^{mcbe}) c^{mad} \\
&\quad + (T_{L(R)}^{mdbc} + T_{L(R)}^{mbcd} + T_{L(R)}^{mdcb} + T_{L(R)}^{mcdb}) c^{mae}] \mathcal{M}_9^{abcde}, \\
&= - [2T_{L(R)}^{acbd} + 2T_{L(R)}^{adbc} - (T_L + T_R) (c^{ebc} c^{eda} + c^{ebd} c^{eca})] \mathcal{M}_7^{abcd} \\
&\quad - 2 [2T_{L(R)}^{abcd} + 2T_{L(R)}^{adcb} + (T_L + T_R) (c^{ebc} c^{eda} + c^{eba} c^{edc})] \mathcal{M}_8^{abcd} \\
&\quad + 0 \mathcal{M}_9^{abcde}, \\
b \tilde{\Delta}_{6L(R)} &= \mathcal{M}_{10L(R)} + \mathcal{M}_{11L(R)}, \\
b \tilde{\Delta}_{7L(R)} &= \mathcal{M}_{11L(R)} + i \mathcal{M}_{13L(R)} + \frac{i}{2} c^{abc} \mathcal{M}_{14L(R)}, \\
b \tilde{\Delta}_{8L(R)} &= \mathcal{M}_{10L(R)} - i \mathcal{M}_{12L(R)} - \frac{i}{2} c^{abc} \mathcal{M}_{15L(R)},
\end{aligned} \tag{68}$$

The coefficient in $b \cdot \tilde{\Delta}_{5L(R)}$ of $\mathcal{M}_9^{abcde}$ turns out to be zero!, which can be proved by expressing the c^{abc} 's as commutators of the corresponding matrices and then using the cyclicity of the trace. If this coefficient would not be zero the system of equations below would be incompatible! We have also used the relationship given in eq. (65).

Now, we impose the that the sum of the terms on the right hand side of eq. (68) matches the right hand side of eq. (64), after removal, of course, of the term carrying the essential non-Abelian anomaly. The following linear system of equations is thus obtained:

$$\begin{aligned}
* \text{ from } \mathcal{M}_1^{ab} : \quad 2a_1 - 2a_2 &= \frac{1}{(4\pi)^2} \frac{T_L + T_R}{3}, \\
* \text{ from } \mathcal{M}_2^{abc} : \quad -2a_2 - i c &= -\frac{1}{(4\pi)^2} \frac{T_L + T_R}{3} \quad \text{and} \quad d_{L(R)} = 0,
\end{aligned}$$

- * from $\mathcal{M}_4^{abc}$: $2a_1 + i c = \frac{1}{(4\pi)^2} (T_L + T_R) \frac{2}{3}$ and $d_{L(R)} = 0$,
- * from $\mathcal{M}_5^{abc}$: $d_{L(R)} = 0$, we will not write this unknown anymore
- * from $\mathcal{M}_7^{abcd}$: $-\frac{i}{2} c + T_L f_L + T_R f_R = \frac{1}{(4\pi)^2} \frac{T_L + T_R}{6}$ and $-2 f_{L(R)} = -\frac{1}{(4\pi)^2} \frac{1}{6}$
- * from $\mathcal{M}_8^{abcd}$: $i c - 2T_L f_L - 2T_R f_R = -\frac{1}{(4\pi)^2} \frac{T_L + T_R}{3}$ and $-4 f_{L(R)} = -\frac{1}{(4\pi)^2} \frac{1}{3}$
- * from $\mathcal{M}_{9L(R)}^{abcd}$: $\frac{f_{L(R)}}{2} \times 0 = 0$
- * from $\mathcal{M}_{10L(R)}$: $n_{L(R)} + u_{2L(R)} = \frac{1}{(4\pi)^2} g^2 \left[C_{L(R)} - \frac{C_A}{4} + (\alpha' - 1) \left(\frac{C_{L(R)}}{6} - \frac{C_A}{4} \right) \right]$
- * from $\mathcal{M}_{11L(R)}$: $n_{L(R)} + u_{1L(R)} = \frac{1}{(4\pi)^2} g^2 \left[C_{L(R)} - \frac{C_A}{4} + (\alpha' - 1) \left(\frac{C_{L(R)}}{6} - \frac{C_A}{4} \right) \right]$
- * from $\mathcal{M}_{12L(R)}$: $-i u_{2L(R)} = \frac{i}{(4\pi)^2} g^2 [1 + (\alpha' - 1)] \frac{C_A}{4}$
- * from $\mathcal{M}_{13L(R)}$: $i u_{1L(R)} = -\frac{i}{(4\pi)^2} g^2 [1 + (\alpha' - 1)] \frac{C_A}{4}$
- * from $\mathcal{M}_{14L(R)}^{abc}$: $-\frac{i}{2} u_{1L(R)} = \frac{i}{(4\pi)^2} g^2 [1 + (\alpha' - 1)] \frac{C_A}{8}$
- * from $\mathcal{M}_{15L(R)}^{abc}$: $\frac{i}{2} u_{1L(R)} = -\frac{i}{(4\pi)^2} g^2 [1 + (\alpha' - 1)] \frac{C_A}{8}$

Notice that there are more constraints than unknowns. In fact, several equations appear repeated and the system has 14 different equations and 13 unknowns, which turns to be *compatible* whose *unique solution* is:

$$\begin{aligned}
a_1 &= \frac{1}{(4\pi)^2} \frac{5}{12} (T_L + T_R), & a_2 &= \frac{1}{(4\pi)^2} \frac{1}{4} (T_L + T_R), \\
c &= \frac{i}{(4\pi)^2} \frac{1}{6} (T_L + T_R), & d_{L(R)} &= 0, \\
f_{L(R)} &= \frac{1}{(4\pi)^2} \frac{1}{12}, & n_{L(R)} &= \frac{1}{(4\pi)^2} g^2 \left[C_{L(R)} + (\alpha' - 1) \frac{C_{L(R)}}{6} \right] \\
u_{1L(R)} &= u_{2L(R)} = -\frac{1}{(4\pi)^2} g^2 \frac{C_A}{4} \alpha'.
\end{aligned}$$

In summary, the following choice of $S_{\text{fct}}^{(1)}$ removes up to order $\hbar$ the spurious anomaly terms from the BRS equation (eq. (42))

$$S_{\text{fct}}^{(1)} = \int d^d x \left\{ \frac{\hbar^1}{(4\pi)^2} (T_L + T_R) \left[\frac{5}{12} (\partial_\mu A^\mu)^2 + \frac{1}{4} A_\mu \square A^\mu \right] \right\}$$

$$\begin{aligned}
& - \frac{\hbar^1}{(4\pi)^2} \frac{T_L + T_R}{6} c^{abc} (\partial_\mu A_\nu^a) A^{b\mu} A^{c\nu} \\
& + \frac{\hbar^1}{(4\pi)^2} \frac{T_L^{abcd} + T_R^{abcd}}{12} A_\mu^a A^{b\mu} A_\nu^c A^{d\nu} \\
& + \frac{\hbar^1}{(4\pi)^2} g^2 [C_L + (\alpha' - 1)C_L/6] \frac{i}{2} \bar{\psi} \bar{\gamma}^\mu P_L \overleftrightarrow{\partial}_\mu \psi \\
& + \frac{\hbar^1}{(4\pi)^2} g^2 [C_R + (\alpha' - 1)C_R/6] \frac{i}{2} \bar{\psi}' \bar{\gamma}^\mu P_R \overleftrightarrow{\partial}_\mu \psi' \\
& - \frac{\hbar^1}{(4\pi)^2} g^2 \frac{C_A}{4} [1 + (\alpha' - 1)] (\bar{L}s\psi + Ls\bar{\psi} + \bar{R}s\psi' + Rs\bar{\psi}') \Big\} \\
& + \hbar^1 l_g^{(1)} L_g + \hbar^1 l_\psi^{(1)} L_\psi + \hbar^1 l_{\psi'}^{(1)} L_{\psi'} + \hbar^1 l_A^{(1)} L_A + \hbar^1 l_\omega^{(1)} L_\omega,
\end{aligned} \tag{69}$$

where $l_g^{(1)}$, $l_\psi^{(1)}$, $l_{\psi'}^{(1)}$, $l_A^{(1)}$ and $l_\omega^{(1)}$ are arbitrary coefficients which will determine the renormalization conditions at the one-loop order and L_g , L_ψ , $L_{\psi'}$, L_A , L_ω are any of the generalizations to the d -dimensional space-time of Dimensional Regularization of the corresponding b -invariant four-dimensional operators. We have written $S_{\text{fct}}^{(1)}$ in the d -dimensional space of Dimensional Regularization so that diagrammatic computation can be readily done.

Notice that no terms in $\bar{\omega}$ or ρ are added, excepting the combination $\tilde{\rho}$ in the invariant counterterms; therefore the ghost equation holds up to order $\hbar$.

Notice also that, because of finite counterterms depending of the external fields L , $\bar{L}$, R and $\bar{R}$ have been added, we have now, *e.g.* (set $l_\omega^{(1)} = 0$),

$$\begin{aligned}
\left. \frac{\delta \Gamma_{\text{ren}}}{\delta \bar{L}(x)} \right|_{L \equiv 0} & \neq N[s\psi](x) \cdot \Gamma_{\text{ren}} \\
\left. \frac{\delta \Gamma_{\text{ren}}}{\delta \bar{L}(x)} \right|_{L \equiv 0} & = N[s\psi](x) \cdot \Gamma_{\text{ren}} + \hbar^1 u_{1L}^{(1)} s\psi(x) + O(\hbar^2) = N[s\psi + \hbar^1 u_{1L}^{(1)} s\psi](x) \cdot \Gamma_{\text{ren}} + O(\hbar^2),
\end{aligned}$$

which can be interpreted as a “non-minimal” renormalization of the operator insertion (we always apply minimal subtraction to the regularized diagrams, but the action have explicit higher order finite counterterms). Moreover, with $l_A^{(1)} \neq 0$ or $l_\omega^{(1)} \neq 0$, the insertion of the non-linear BRS-variations of the fields can be renormalized in different ways, all of them compatible with the BRS identities (*i.e.* there are several renormalization conditions for that insertions compatible with the identities).

Of course, the finite counterterms given in eq. (69), are strongly dependent on the choice for the $O(\hbar^0)$ action, S_0 (the Dimensional Regularization classical action in eq. (35)). If we would have chosen the other mentioned fermionic vertex, the counterterms would have certainly been completely different (and more complicated). But anyway the procedure would have been equally correct.

Notice that if the classical original theory is CP invariant the finite counterterms of eq. (69) are also CP invariant. This was expected, due to the comments in the second paragraph below eq. (34).

3.7. The renormalization group equation

It is plain that the standard textbook techniques usually employed to derive the renormalization group equation for dimensionally renormalized vector theories like Q.E.D and Q.C.D are of no use here. The formalism of bare fields, bare coupling constants and multiplicative renormalization of the classical action breaks down as shows eq. (37). The generalization of this formalism to include evanescent operators, and the corresponding renormalized coupling constants, becomes a bit awkward when the regularized theory does not possess the symmetries that should hold in the quantum field theory. Indeed a host of bare evanescent non-symmetric operators, each bringing in a new renormalized coupling constant, may enter the game. This is in addition to the finite non-evanescent non-symmetric counterterms needed to restore the symmetry broken in the dimensional renormalization process. These finite counterterms should also come from bare operators. Hence, the construction of an appropriate bare action and the derivation of the true (*i.e.* only involving as many coupling constants and anomalous dimensions as there are classical couplings and fields) renormalization group equation is a rather involved task (for related information see ref. [32]). We shall abandon this quest altogether. Indeed, thanks to the *Quantum Action Principle*, Bonneau identities and the formalism of algebraic renormalization there is no need to introduce bare fields and bare couplings to derive the true renormalization group equation. We only need the renormalized 1PI functional, Γ_{ren} , which is symmetric up to the order in $\hbar$ demanded, obtained by the method explained in this paper.

Let $\Gamma_{\text{ren}}[\varphi, \Phi, K_\Phi; g, \alpha', \mu]$ be the dimensionally renormalized up to order $\hbar^n$ 1PI functional of our theory, which we will take to be anomaly free all along the current subsection. Γ_{ren} depends explicitly on the fields, collectively denoted by φ and Φ , the external fields, collectively denoted by K_Φ , the coupling constant g , the gauge fixing parameter α' and the Dimensional Regularization scale μ , which is introduced along the lines laid in *v)* of subsection 2.2. We shall assume that finite counterterms, $S_{\text{fct}}^{(n)}$ (see eq. (35)), has been chosen so that the BRS equation $\mathcal{S}(\Gamma_{\text{ren}}) = 0$, the gauge-fixing condition $\mathcal{B}(\Gamma_{\text{ren}}) = 0$ (eq. (46)), and the ghost equation $\mathcal{G} \Gamma_{\text{ren}} = 0$ (eq. (47)) hold up to order $\hbar^n$.

The renormalization group equation of the theory gives the expansion of the functional $\mu \frac{\partial \Gamma_{\text{ren}}}{\partial \mu}$ in terms of a certain basis of quantum insertions of ultraviolet dimension 4 and ghost number 0 [7,18]. The coefficients of this expansion, which are formal power series in $\hbar$, are the beta functions and anomalous dimensions of the theory. Since Γ_{ren} satisfies, up to order $\hbar^n$, the equations mentioned in the previous paragraph, the elements of basis of insertions we are seeking is not only constrained by power-counting and ghost number. Indeed, the following three equations hold up to order $\hbar^n$:

$$\begin{aligned} \mathcal{S}_{\Gamma_{\text{ren}}} \mu \frac{\partial \Gamma_{\text{ren}}}{\partial \mu} &= 0, \\ \frac{\delta}{\delta B} \mu \frac{\partial \Gamma_{\text{ren}}}{\partial \mu} &= 0, \\ \mathcal{G} \mu \frac{\partial \Gamma_{\text{ren}}}{\partial \mu} &= 0. \end{aligned} \tag{70}$$

Notice that the operator $\mu \frac{\partial}{\partial \mu}$ commutes with the functional operators $\mathcal{B}$, $\mathcal{G}$ and that

$$\mu \frac{\partial}{\partial \mu} (\mathcal{S}(\Gamma_{\text{ren}})) = \mathcal{S}_{\Gamma_{\text{ren}}} \mu \frac{\partial \Gamma_{\text{ren}}}{\partial \mu}.$$

$\mathcal{S}_{\Gamma_{\text{ren}}}$ stands for the linearized BRS operator:

$$\begin{aligned} \mathcal{S}_{\Gamma_{\text{ren}}} = \int d^4x \left\{ \text{Tr} \frac{\delta \Gamma_{\text{ren}}}{\delta \rho^\mu} \frac{\delta}{\delta A_\mu} + \text{Tr} \frac{\delta \Gamma_{\text{ren}}}{\delta A_\mu} \frac{\delta}{\delta \rho^\mu} + \text{Tr} \frac{\delta \Gamma_{\text{ren}}}{\delta \zeta} \frac{\delta}{\delta \omega} + \text{Tr} \frac{\delta \Gamma_{\text{ren}}}{\delta \omega} \frac{\delta}{\delta \zeta} + \text{Tr} B \frac{\delta \Gamma_{\text{ren}}}{\delta \bar{\omega}} \right. \\ \left. + \frac{\delta \Gamma_{\text{ren}}}{\delta \bar{L}} \frac{\delta}{\delta \psi} + \frac{\delta \Gamma_{\text{ren}}}{\delta \psi} \frac{\delta}{\delta \bar{L}} + \frac{\delta \Gamma_{\text{ren}}}{\delta \bar{R}} \frac{\delta}{\delta \psi'} + \frac{\delta \Gamma_{\text{ren}}}{\delta \psi'} \frac{\delta}{\delta \bar{R}} + \frac{\delta \Gamma_{\text{ren}}}{\delta L} \frac{\delta}{\delta \bar{\psi}} + \frac{\delta \Gamma_{\text{ren}}}{\delta \bar{\psi}} \frac{\delta}{\delta L} \right. \\ \left. + \frac{\delta}{\delta R} \frac{\delta \Gamma_{\text{ren}}}{\delta \bar{\psi}'} + \frac{\delta \Gamma_{\text{ren}}}{\delta \bar{\psi}'} \frac{\delta}{\delta R} \right\}. \end{aligned}$$

The functionals with ultraviolet dimension 4 and ghost number 0 which satisfy up to order $\hbar^n$ the set of equations in eq. (70) form a linear space. Let us construct a basis of this space. In the classical approximation (*i.e.* at order $\hbar^0$), a basis of this space is given [18] by the $b \equiv \mathcal{S}_{\text{cl}}$ -invariant terms L_g , L_ψ^R , $L_{\psi'}^L$, L_ψ^L , $L_{\psi'}^R$, L_A and L_ω , defined as follows:

$$\begin{aligned} L_g &= g \frac{\partial \mathcal{S}_{\text{cl}}}{\partial g} = -2 (S_{\text{cl}AA} + S_{\text{cl}AAA} + S_{\text{cl}AAA}), \\ L_\psi^R &= \mathcal{N}_\psi^R S_{\text{cl}} = 2S_{\text{cl}\bar{\psi}\psi}^R, & L_{\psi'}^L &= \mathcal{N}_{\psi'}^L S_{\text{cl}} = 2S_{\text{cl}\bar{\psi}'\psi'}^L, \\ L_\psi^L &= \mathcal{N}_\psi^L S_{\text{cl}} = 2S_{\text{cl}\bar{\psi}\psi}^L + 2S_{\text{cl}\bar{\psi}\psi A}, & L_{\psi'}^R &= \mathcal{N}_{\psi'}^R S_{\text{cl}} = 2S_{\text{cl}\bar{\psi}'\psi'}^R + 2S_{\text{cl}\bar{\psi}'\psi' A}, \\ L_A &= \mathcal{N}_A S_{\text{cl}} = 2S_{\text{cl}AA} + 3S_{\text{cl}AAA} + 4S_{\text{cl}AAAA} + S_{\text{cl}\bar{\psi}\psi A} + S_{\text{cl}\bar{\psi}'\psi' A} - S_{\text{cl}\rho\omega} - S_{\text{cl}\bar{\omega}\omega}, \\ L_\omega &= \mathcal{N}_\omega S_{\text{cl}} = S_{\text{cl}\rho\omega} + S_{\text{cl}\bar{\omega}\omega} + S_{\text{cl}\rho\omega A} + S_{\text{cl}\zeta\omega\omega} + S_{\text{cl}\bar{L}s\psi} + S_{\text{cl}\bar{R}s\psi'} + S_{\text{cl}Ls\bar{\psi}} + S_{\text{cl}Rs\bar{\psi}'}, \end{aligned} \quad (71)$$

where the $S_{\text{cl}}\dots$ terms are the corresponding B -independent terms of the classical action $\mathcal{S}_{\text{cl}}$ given in eq. (27). The symmetric differential operators [18] $\mathcal{N}_\psi^R$, $\mathcal{N}_\psi^L$, $\mathcal{N}_{\psi'}^R$, $\mathcal{N}_{\psi'}^L$, $\mathcal{N}_A$ and $\mathcal{N}_\omega$ in eq. (71) are given by the following identities:

$$\begin{aligned} \mathcal{N}_\psi^L &\equiv N_\psi^L + N_{\bar{\psi}}^L - N_L - N_{\bar{L}}, & \mathcal{N}_\psi^R &\equiv N_\psi^R + N_{\bar{\psi}}^R, \\ \mathcal{N}_{\psi'}^L &\equiv N_{\psi'}^L + N_{\bar{\psi}'}^L, & \mathcal{N}_{\psi'}^R &\equiv N_{\psi'}^R + N_{\bar{\psi}'}^R - N_R - N_{\bar{R}}, \\ \mathcal{N}_A &\equiv \text{Tr}(N_A - N_\rho - N_B - N_{\bar{\omega}}) + 2\alpha \frac{\partial}{\partial \alpha}, \\ \mathcal{N}_\omega &\equiv \text{Tr}(N_\omega - N_\zeta). \end{aligned} \quad (72)$$

The field-counting operators $N_\psi^{\text{L(R)}}$, $N_{\psi'}^{\text{L(R)}}$, N_L , $N_{\bar{L}}$, N_R , $N_{\bar{R}}$, N_A , N_ω , $N_{\bar{\omega}}$ and N_ζ are the operators in eq. (38) for $d = 4$.

We shall work out now a basis (up to order $\hbar^n$, of course) of the space of joint solutions of eqs. (70) with the help of the statement made in the second paragraph of subsection 2.4. (see ref. [18] for further details). Up to order $\hbar^n$, a quantum extension of the classical basis

in eq. (71) is furnished [18] by the action on Γ_{ren} of the symmetric differential operators defined above along with symmetric differential operator $g\partial_g$. Indeed, the functionals

$$g\frac{\partial\Gamma_{\text{ren}}}{\partial g}, \mathcal{N}_{\psi}^{\text{R}}\Gamma_{\text{ren}}, \mathcal{N}_{\psi'}^{\text{L}}\Gamma_{\text{ren}}, \mathcal{N}_{\psi}^{\text{L}}\Gamma_{\text{ren}}, \mathcal{N}_{\psi'}^{\text{R}}\Gamma_{\text{ren}}, \mathcal{N}_A\Gamma_{\text{ren}}, \mathcal{N}_{\omega}\Gamma_{\text{ren}},$$

which are to be understood as formal series expansions in $\hbar$, match the corresponding functionals in eq. (71) at order $\hbar^0$ and the symmetric differential operators $g\partial_g, \mathcal{N}_{\psi}^{\text{R}}, \mathcal{N}_{\psi}^{\text{L}}, \mathcal{N}_{\psi'}^{\text{R}}, \mathcal{N}_{\psi'}^{\text{L}}, \mathcal{N}_A$ and $\mathcal{N}_{\omega}$ are compatible (this is why they are symmetric), up to order $\hbar^n$, with the BRS equation, the gauge-fixing condition and the ghost equation, in the sense we spell out next.

Let $\mathcal{F}$ be the linear space of functionals with ultraviolet dimension 4 and ghost number 0 which satisfy (up to order $\hbar^n$) the BRS equation, the gauge-fixing condition and the ghost equation: $\mathcal{S}(\Gamma) = 0, \mathcal{B}(\Gamma) = 0, \mathcal{G}(\Gamma) = 0, \forall \Gamma \in \mathcal{F}$. An operator $\mathcal{D}$ acting on $\mathcal{F}$ is said to be compatible with the BRS equation, the gauge-fixing condition and the ghost equation if, by definition, the following set of equations hold:

$$\mathcal{S}_{\Gamma}(\mathcal{D}\Gamma) = 0, \quad \frac{\delta}{\delta B}(\mathcal{D}\Gamma) = 0, \quad \mathcal{G}(\mathcal{D}\Gamma) = 0.$$

In summary, up to order $\hbar^n$, any joint solution of eqs. (70) is a linear combination with $\hbar$ -dependent coefficients of the following functionals:

$$g\frac{\partial\Gamma_{\text{ren}}}{\partial g}, \mathcal{N}_{\psi}^{\text{R}}\Gamma_{\text{ren}}, \mathcal{N}_{\psi'}^{\text{L}}\Gamma_{\text{ren}}, \mathcal{N}_{\psi}^{\text{L}}\Gamma_{\text{ren}}, \mathcal{N}_{\psi'}^{\text{R}}\Gamma_{\text{ren}}, \mathcal{N}_A\Gamma_{\text{ren}}, \mathcal{N}_{\omega}\Gamma_{\text{ren}}. \quad (73)$$

In particular, if μ is the arbitrary “scale” introduced by hand for each loop integration in minimal Dimensional Renormalization (see v) in subsection 2.2.) , then, $\mu\frac{\partial\Gamma_{\text{ren}}}{\partial\mu}$ has an expansion in the quantum basis given above:

$$\left[\mu\frac{\partial}{\partial\mu} + \beta g\frac{\partial}{\partial g} - \gamma_{\psi}^{\text{L}}\mathcal{N}_{\psi}^{\text{L}} - \gamma_{\psi'}^{\text{R}}\mathcal{N}_{\psi'}^{\text{R}} - \gamma_{\psi}^{\text{R}}\mathcal{N}_{\psi}^{\text{R}} - \gamma_{\psi'}^{\text{L}}\mathcal{N}_{\psi'}^{\text{L}} - \gamma_A\mathcal{N}_A - \gamma_{\omega}\mathcal{N}_{\omega} \right] \Gamma_{\text{ren}} = 0. \quad (74)$$

This equation holds up to order $\hbar^n$ and it is the renormalization group equation of our theory. Notice that the expansion $\mu\frac{\partial\Gamma_{\text{ren}}}{\partial\mu}$ in terms of the basis in eq. (73) is only possible if the renormalized functional satisfies order by order the BRS identity (hence, there is anomaly cancellation, in particular) and the gauge-fixing condition. It is important to notice that the regularized 1PI functional does not generally satisfy an equation such as eq. (74).

Let us notice that the coefficients of the expansions of $\mu\frac{\partial\Gamma_{\text{ren}}}{\partial\mu}$ in the quantum basis of insertions in eq. (73) are the beta functions and anomalous dimensions of the theory. These coefficients are to be understood as formal expansions in $\hbar$. The first non-trivial contribution to these expansions is always of order $\hbar$, since the lowest order contribution to Γ_{ren} is the classical action, S_{cl} , and $\mu\frac{\partial S_{\text{cl}}}{\partial\mu} = 0$.

We next proceed to the computation of the beta function and anomalous dimensions of our theory. This is done [6] by expressing first $\mu\frac{\partial\Gamma_{\text{ren}}}{\partial\mu}$, $g\frac{\partial\Gamma_{\text{ren}}}{\partial g}$ and $\mathcal{N}_{\phi}\Gamma_{\text{ren}}$, where

$\mathcal{N}_\phi$ denotes any of the differential operators in eq. (72), as insertions in Γ_{ren} of linearly independent local normal products of the fields and their derivatives; then, these insertions are substituted into the left hand side of the renormalization group equation (eq. (74)) and, finally, one solves for β , γ_ψ^L , $\gamma_{\psi'}^R$, γ_ψ^R , $\gamma_{\psi'}^L$, γ_A and γ_ω , the set of equations involving only numbers which results from the linear independence of the aforementioned local normal products.

That $g \frac{\partial \Gamma_{\text{ren}}}{\partial g}$ and $\mathcal{N}_\phi \Gamma_{\text{ren}}$, where $\mathcal{N}_\phi$ denotes any of the differential operators in eq. (72), can be expressed as insertions in Γ_{ren} of normal products is a mere consequence of the *Quantum Action Principle*: QAP1 and QAP3, given by eqs. (14) and (16), lead to

$$\frac{\partial \Gamma_{\text{ren}}}{\partial g} = \text{N} \left[\frac{\partial (S_0 + S_{\text{fct}}^{(n)})}{\partial g} \right] \cdot \Gamma_{\text{ren}}, \quad (75)$$

and

$$\mathcal{N}_\phi \Gamma_{\text{ren}} = \text{N} [\mathcal{N}_\phi (S_0 + S_{\text{fct}}^{(n)})] \cdot \Gamma_{\text{ren}}, \quad (76)$$

respectively. Hence, every element of the symmetric quantum basis in eq. (73) is a local combination of quantum insertions of integrated standard (*i.e.* non-evanescent) monomials with ultraviolet dimension 4 and ghost number 0 (see subsection 2.4.). It should not be overlooked the fact $S_0 + S_{\text{fct}}^{(n)}$ in eqs. (75) and (76) is the non-singular contribution to the Dimensional Regularization action, $S_{DReg}^{(n)}$ (see subsection 3.2.), employed to obtain the minimal dimensionally renormalized 1PI functional up to order $\hbar^n$, Γ_{ren} , by following the algorithm spell out in subsection 2.2. The formal functional $S_0 + S_{\text{fct}}^{(n)}$ is therefore an object in the d -dimensional space-time of Dimensional Regularization. If $n > 2$, it would not do to use some other functional, S' , such that $(S_0 + S_{\text{fct}}^{(n)}) - S' = \hat{S}'$ is an integrated evanescent operator whose $\hbar$ -expansion has contributions of order $\hbar^m$, with $m \leq n-2$. Recall that the contribution $O(\hbar^{(n-1)})$ in $\hat{S}'$ is not be relevant here since, being an evanescent object, the lowest order contribution coming from its insertion in Γ_{ren} is $O(\hbar^n)$ and local, so that it does not contribute to the renormalization group equation at order $\hbar^n$; a similar argument holds for the term of order $\hbar^n$ in $\hat{S}'$.

Now, by substituting eqs. (75) and (76) in eq. (74) one obtains the following equation

$$\mu \frac{\partial \Gamma_{\text{ren}}}{\partial \mu} = -\beta g \text{N} \left[\frac{\partial (S_0 + S_{\text{fct}}^{(n)})}{\partial g} \right] \cdot \Gamma_{\text{ren}} + \sum_{\phi} \gamma_{\phi} \text{N} [\mathcal{N}_{\phi} (S_0 + S_{\text{fct}}^{(n)})] \cdot \Gamma_{\text{ren}}, \quad (77)$$

where ϕ labels the symmetric differential operators in eq. (72). Hence, to compute the beta functions and anomalous dimensions of the theory we only need an independent way of computing $\mu \frac{\partial \Gamma_{\text{ren}}}{\partial \mu}$ in terms of the insertions on the right hand side of eq. (77). We shall set up this independent way of computation next.

Unlike for the subtraction algorithm employed in ref. [3], in minimal dimensional renormalization, one can not use (see ref. [6]) the *Quantum Action Principle* to obtain an equation for $\mu \frac{\partial \Gamma_{\text{ren}}}{\partial \mu}$ analogous to eqs. (75), (76). Indeed, in minimal dimensional renormalization, μ is not a parameter of the action ($S_0 + S_{\text{fct}}^{(n)}$), but a parameter which is introduced

by hand in each loop-momentum integration (as a factor μ^{4-d}) before computing the renormalized value of the corresponding Feynman diagram.

The problem of expressing $\mu \frac{\partial \Gamma_{\text{ren}}}{\partial \mu}$ as an insertion in Γ_{ren} of a linear combination normal product operators was also solved by Bonneau [6] in the framework of rigorous minimal dimensional renormalization. He obtain a formula similar to eq. (25), *i.e.* a Zimmermann-like identity, for a scalar case with no added finite counterterms in the action. Here we need the generalization to the presence of several type of fields, including external fields, the presence of hatted objects and also the presence of finite counterterms up to order $\hbar^n$, in the action ($S_0 + S_{\text{fct}}^{(n)}$). The needed generalization reads

$$\begin{aligned}
\mu \frac{\partial \Gamma_{\text{ren}}}{\partial \mu} = & \sum_{n=0}^4 \sum_{\{j_1, \dots, j_n\}} \sum_{N_l > 0} N_l \left[\right. \\
& \sum_{r=0}^{\omega(J)} \sum_{\substack{\{i_1 \dots i_r\} \\ 1 \leq i_j \leq n-1}} \left\{ \text{r.s.p.} \frac{(-i)^r}{r!} \frac{\partial^r}{\partial p_{i_1}^{\mu_1} \dots \partial p_{i_r}^{\mu_r}} (-i\hbar) \overline{\langle \tilde{\phi}_{j_1}(p_1) \dots \tilde{\phi}_{j_n}(p_n = -\sum p_i) \rangle_{K=0}^{\text{1PI}, (N_l)}} \Big|_{p_i=0} \right\} \\
& \times \frac{1}{n!} \int d^4x \, \text{N} \left[\phi_{j_n} \prod_{k=n-1}^1 \left\{ \left(\prod_{\{\alpha/i_\alpha=k\}} \partial_{\mu_\alpha} \right) \phi_{j_k} \right\} \right] (x) \cdot \Gamma_{\text{ren}} + \\
& \sum_{\Phi} \sum_{r=0}^{\omega(J; \Phi)} \sum_{\substack{\{i_1 \dots i_r\} \\ 1 \leq i_j \leq n}} \left\{ \text{r.s.p.} \frac{(-i)^r}{(r)!} \frac{\partial^r}{\partial p_{i_1}^{\mu_1} \dots \partial p_{i_r}^{\mu_r}} \right. \\
& \left. \overline{\langle \tilde{\phi}_{j_1}(p_1) \dots \tilde{\phi}_{j_n}(p_n) \text{N}[s\Phi](p_{n+1} = -\sum p_i) \rangle_{K=0}^{\text{1PI}, (N_l)}} \Big|_{p_i=0} \right\} \\
& \times \frac{1}{n!} \int d^4x \, (K_\Phi(x) \text{N} \left[\prod_{k=n}^1 \left\{ \left(\prod_{\{\alpha/i_\alpha=k\}} \partial_{\mu_\alpha} \right) \phi_{j_k} \right\} \right] (x) \cdot \Gamma_{\text{ren}} \left. \right] , \tag{78}
\end{aligned}$$

where $J = \{j_1, \dots, j_n\}$, and, $\omega(J)$ and $\omega(J; \Phi)$ stand, respectively, for the overall ultraviolet degree of divergence of $\langle \tilde{\phi}_{j_1}(p_1) \dots \tilde{\phi}_{j_n}(p_n) \rangle_{K=0}^{\text{1PI}}$ and $\langle \tilde{\phi}_{j_1}(p_1) \dots \tilde{\phi}_{j_n}(p_n) \text{N}[s\Phi](p_{n+1}) \rangle_{K=0}^{\text{1PI}}$. The bar upon the 1PI Green functions means that all subdivergences has been subtracted from the Feynman diagrams which contribute to them. This is as in eq. (48). The novel feature here is the presence of the superscript N_l on the upper right of the Green function, which indicates that only Feynman diagrams with precisely N_l loops are to be considered. Notice that one then sums over all number loops upon multiplication by N_l of the N_l -contribution.

The r.s.p. of the subtracted (subdivergences only) 1PI Feynman diagrams contributing to the right hand side of eq. (78) is a local polynomial in the d -dimensional space of Dimensional Regularization of external momenta associated with the fields of the corresponding 1PI function; so, in general, it will contain hatted and barred objects (metrics, momenta, gamma matrices). Therefore, the formula will generate automatically also evanescent (also called anomalous) insertions. The meaning of $\times$ in last formula is the same as in eq. (25).

Hence, the expansion in eq. (78) can be recast into the form

$$\mu \frac{\partial \Gamma_{\text{ren}}}{\partial \mu} = \sum_i \bar{r}_i N[\bar{\mathcal{W}}_i] \cdot \Gamma_{\text{ren}} + \sum_j \hat{r}_j N[\hat{\mathcal{W}}_j] \cdot \Gamma_{\text{ren}} \quad (79)$$

where $\bar{\mathcal{W}}_i$ and $\hat{\mathcal{W}}_j$ denote, respectively, the barred (non-evanescent) and hatted (evanescent) elements of a basis of integrated monomials with ghost number 0 and ultraviolet dimension 4. The symbols $\bar{r}_i$ and $\hat{r}_j$ in eq. (79) are coefficients defined as formal expansions in $\hbar$ (starting at $\hbar$, since their computation involves that of the r.s.p. of the divergent part of the appropriate 1PI function). The order $\hbar^m$, $0 < m \leq n$, contribution to these coefficients is obtained as follows:

- * Take a divergent, by power-counting, 1PI function with some number of fields of definite type (maybe including some external field coupled to a BRS variation).

- * Compute the residue of the simple pole (r.s.p.) of every $O(\hbar^m)$ -graph constructed with the Feynman rules derived from the action $(S_0 + S_{\text{fct}}^{(n)})$ and contributing to that 1PI function, with all their subdivergences subtracted.

- * Multiply the last quantity by the number of the loops of the graph

- * Sum over all graphs contributing to the 1PI function

- * Being the sum a local expression, their different terms can be interpreted as Feynman rules of tree-level integrated insertions of Lorentz invariant (normal or anomalous) composite operators, formed by just the fields of the original 1PI function and some different combination of metrics, derivatives and gamma matrices.

- * Do the same steps for every divergent 1PI function. Then you will obtain the expansion eq. (79) with the $O(\hbar^m)$ coefficients.

Explicit expressions for each $\bar{r}_i$ and $\hat{r}_j$ similar to that ones for the expansion of the anomalous breaking can be given. For example, if

$$\begin{aligned} \bar{\mathcal{W}}_2^{ab} &= \int \bar{\square} A_{\bar{\mu}}^a A^{b\bar{\mu}}, & \hat{\mathcal{W}}_{21}^{ab} &= \int \bar{\square} A_{\bar{\mu}}^a A^{b\hat{\mu}}, \\ \hat{\mathcal{W}}_{22}^{ab} &= \int \hat{\square} A_{\bar{\mu}}^a A^{b\bar{\mu}}, & \hat{\mathcal{W}}_{23}^{ab} &= \int \hat{\square} A_{\hat{\mu}}^a A^{b\hat{\mu}}, \end{aligned}$$

then

$$\begin{aligned} \bar{r}_2^{ab} &= \frac{-(-i)^2}{2} N_l \times \text{coef. in r.s.p. } (-i\hbar) \overline{\langle A_{\bar{\mu}}^a(p_1) A_{\bar{\nu}}^b(p_2 \equiv -p_1) \rangle}^{1\text{PI}} \text{ of } \bar{p}_1^2 \bar{g}^{\mu\nu} \\ \hat{r}_{21}^{ab} &= \frac{-(-i)^2}{2} N_l \times \text{coef. in r.s.p. } (-i\hbar) \overline{\langle A_{\bar{\mu}}^a(p_1) A_{\bar{\nu}}^b(p_2 \equiv -p_1) \rangle}^{1\text{PI}} \text{ of } \bar{p}_1^2 \hat{g}^{\mu\nu}, \\ \hat{r}_{22}^{ab} &= \frac{-(-i)^2}{2} N_l \times \text{coef. in r.s.p. } (-i\hbar) \overline{\langle A_{\bar{\mu}}^a(p_1) A_{\bar{\nu}}^b(p_2 \equiv -p_1) \rangle}^{1\text{PI}} \text{ of } \hat{p}_1^2 \bar{g}^{\mu\nu}, \\ \hat{r}_{23}^{ab} &= \frac{-(-i)^2}{2} N_l \times \text{coef. in r.s.p. } (-i\hbar) \overline{\langle A_{\bar{\mu}}^a(p_1) A_{\bar{\nu}}^b(p_2 \equiv -p_1) \rangle}^{1\text{PI}} \text{ of } \hat{p}_1^2 \hat{g}^{\mu\nu}; \end{aligned}$$

and so on.

Next, by using the Bonneau identities technique we shall express (see subsection 2.4.) every evanescent insertion $N[\hat{\mathcal{W}}_j] \cdot \Gamma_{\text{ren}}$ on the right hand side of eq. (79) as a linear combination of standard (*i.e.* non-evanescent) insertions $N[\bar{\mathcal{W}}_i] \cdot \Gamma_{\text{ren}}$:

$$N[\hat{\mathcal{W}}_j] \cdot \Gamma_{\text{ren}} = \sum_i c_{ji} N[\bar{\mathcal{W}}_i] \cdot \Gamma_{\text{ren}}. \quad (80)$$

The coefficients c_{ji} are formal expansions in powers of $\hbar$, having no order $\hbar^0$ contribution. The use of the previous equation turns eq. (79) into the following equation:

$$\mu \frac{\partial \Gamma_{\text{ren}}}{\partial \mu} = \sum_i r_i N[\bar{\mathcal{W}}_i] \cdot \Gamma_{\text{ren}}, \quad (81)$$

where $r_i = \bar{r}_i + \sum_j \hat{r}_j c_{ji}$, so that $r_i = \bar{r}_i$ at order $\hbar$. Notice that, because the evanescent insertions are to be expanded according to the Bonneau identities in eq. (25), the computation of r_i up to order $\hbar^n$ only involves the contributions to $\hat{r}_j$ up to order $\hbar^{(n-1)}$, whereas the contributions to $\bar{r}_i$ are needed up to order $\hbar^n$.

Now, since $g \frac{\partial}{\partial g} [\bar{S}_0 + \bar{S}_{\text{fct}}^{(n)}]$, $\mathcal{N}_\phi [\bar{S}_0 + \bar{S}_{\text{fct}}^{(n)}]$, $g \frac{\partial}{\partial g} [(S_0 - \bar{S}_0) + (S_{\text{fct}}^{(n)} - \bar{S}_{\text{fct}}^{(n)})]$ and $\mathcal{N}_\phi [(S_0 - \bar{S}_0) + (S_{\text{fct}}^{(n)} - \bar{S}_{\text{fct}}^{(n)})]$ are linear combinations of monomials of ultraviolet dimension 4 and ghost number 0, it is advisable to choose the basis of monomials $\{\bar{\mathcal{W}}_i, \hat{\mathcal{W}}_j\}$, which occurs in eq. (79), in such a way that it contains the former set of monomials. In other words, the basis $\{\bar{\mathcal{W}}_i, \hat{\mathcal{W}}_j\}$ so chosen makes it possible to obtain the coefficients on the right hand side of the following equations

$$\begin{aligned} g \frac{\partial}{\partial g} [\bar{S}_0 + \bar{S}_{\text{fct}}^{(n)}] &= \sum_i \bar{w}_{gi} \bar{\mathcal{W}}_i, \quad \mathcal{N}_\phi [\bar{S}_0 + \bar{S}_{\text{fct}}^{(n)}] = \sum_i \bar{w}_{\phi i} \bar{\mathcal{W}}_i \\ g \frac{\partial}{\partial g} [(S_0 - \bar{S}_0) + (S_{\text{fct}}^{(n)} - \bar{S}_{\text{fct}}^{(n)})] &= \sum_j \hat{w}_{gj} \hat{\mathcal{W}}_j, \\ \mathcal{N}_\phi [(S_0 - \bar{S}_0) + (S_{\text{fct}}^{(n)} - \bar{S}_{\text{fct}}^{(n)})] &= \sum_j \hat{w}_{\phi j} \hat{\mathcal{W}}_j, \end{aligned}$$

by reading them from the left hand side of the corresponding equation. The coefficients $\bar{w}_{gi}$, $\hat{w}_{gi}$, $\bar{w}_{\phi i}$ and $\hat{w}_{\phi i}$ have expansions in powers of $\hbar$; these expansions, though, have now contributions of order $\hbar^0$.

It is a trivial exercise to check that eq. (77) can be recast into the form

$$\mu \frac{\partial \Gamma_{\text{ren}}}{\partial \mu} = \sum_i (-\beta \bar{w}_{gi} + \sum_\phi \gamma_\phi \bar{w}_{\phi i}) N[\bar{\mathcal{W}}_i] \cdot \Gamma_{\text{ren}} + \sum_j (-\beta \hat{w}_{gj} + \sum_\phi \gamma_\phi \hat{w}_{\phi j}) N[\hat{\mathcal{W}}_j] \cdot \Gamma_{\text{ren}}$$

The substitution of eq. (80) into the last equation gives the following result

$$\mu \frac{\partial \Gamma_{\text{ren}}}{\partial \mu} = \sum_i (-\beta w_{gi} + \sum_\phi \gamma_\phi w_{\phi i}) N[\bar{\mathcal{W}}_i] \cdot \Gamma_{\text{ren}}, \quad (82)$$

where

$$w_{gi} = \bar{w}_{gi} + \sum_j \hat{w}_{gj} c_{ji} \quad \text{and} \quad w_{\phi i} = \bar{w}_{\phi i} + \sum_j \hat{w}_{\phi j} c_{ji},$$

where the index j labels the elements of the “evanescent” basis $\{\hat{\mathcal{W}}_j\}$.

Finally, by comparing eqs. (81) and (82), one obtains an overdeterminate system of linear equations whose unknowns are the coefficients β and γ_ϕ :

$$r_i = -\beta w_{gi} + \sum_\phi \gamma_\phi w_{\phi i}, \quad (83)$$

where the index i runs over the elements of the standard (*i.e.* with no evanescent operator) basis $\{\bar{\mathcal{W}}_i\}$.

Several remarks about eq. (78) are now in order.

* This equation is only correct if the finite counterterms in $S_{\text{fct}}^{(n)}$ are independent of μ . This poses no problem, because the finite counterterms we need to add to the action to restore the BRS identities do not depend on μ , thanks to the fact that in the Bonneau identities we have to compute only r.s.p.s of 1PI functions with all subdivergences minimally subtracted and these r.s.p. are always mass and μ independent.

* The counterpart of eq. (78) in ref. [6] replaces the factor N_l , which counts the number of loops, with the operator $\hbar \frac{\partial}{\partial \hbar}$. This is because, *if no terms of order $\hbar^m$, $m > 0$ are present in the action*, then the powers in $\hbar$ of a graph counts its number of loops. But now we have added higher order terms in $\hbar$ to the action and so we cannot use $\hbar$ as a loop meter. This can be superseded by introducing a new parameter l , which divides the whole action, including the finite counterterms. Then l^{N_l-1} would be the factor attached to a N_l -loop Feynman diagram. Note that only in the limit $l \rightarrow 1$ the finite counterterms restore the BRS identities. Therefore, the factor N_l in eq. (78) is replaced by $\lim_{l \rightarrow 1} \{1 + l \frac{\partial}{\partial l}\}$. Thanks to the action principles, the action of the operator $\lim_{l \rightarrow 1} \{1 + l \frac{\partial}{\partial l}\}$ on a 1PI Green function computed for arbitrary l can be replaced with the insertion of the integrated operator $1 - N[i(S_0 + S_{\text{fct}}^{(n)})]$ into that 1PI function at $l = 1$. It should be stressed that this is true only if there is no explicit μ -dependence in the finite counterterms.

Let us move on and compute the beta function and anomalous dimensions of our theory at order $\hbar$. At $\hbar$ -order, eq. (83) reads

$$\bar{r}_i^{(1)} = -\beta^{(1)} \bar{w}_{gi}^{(0)} + \sum_\phi \gamma_\phi^{(1)} \bar{w}_{\phi i}^{(0)}, \quad (84)$$

for $r_i = \bar{r}_i^{(1)} \hbar + O(\hbar^2)$, $\beta = \beta^{(1)} \hbar + O(\hbar^2)$, $w_{gi} = \bar{w}_{gi}^{(0)} + O(\hbar)$, $\gamma_\phi^{(1)} = \gamma_\phi^{(1)} \hbar + O(\hbar^2)$ and $w_{\phi i} = \bar{w}_{\phi i}^{(0)} + O(\hbar)$. To obtain $\{\bar{r}_i^{(1)}\}$, we just need a suitable basis of integrated barred monomials $\{\bar{\mathcal{W}}_i\}$ with ultraviolet dimension 4 and ghost number 0: the evanescent monomials $\{\hat{\mathcal{W}}_j\}$ are not needed at this order in $\hbar$. The basis $\{\bar{\mathcal{W}}_i\}$ is furnished by the following list of monomials with ghost number 0, dimension 4, free gauge indices and fully contracted Lorentz indices:

* a) *Monomials with only As*

$$\begin{aligned}
\bar{\mathcal{W}}_1^{ab} &\equiv \int (\partial_{\bar{\mu}} \partial_{\bar{\nu}} A^{a\bar{\mu}}) A^{b\bar{\nu}}, & \bar{\mathcal{W}}_2^{ab} &\equiv \int (\bar{\square} A_{\bar{\mu}}^a) A^{b\bar{\mu}}, \\
\bar{\mathcal{W}}_3^{abc} &\equiv \int (\partial_{\bar{\mu}} A_{\bar{\nu}}^a) A^{b\bar{\mu}} A^{c\bar{\nu}}, & \bar{\mathcal{W}}_4^{abc} &\equiv \int (\partial_{\bar{\mu}} A^{a\bar{\mu}}) A_{\bar{\nu}}^b A^{c\bar{\nu}}, \\
\bar{\mathcal{W}}_5^{abc} &\equiv \int \varepsilon_{\mu\nu\rho\alpha} (\partial^\alpha A^{b\bar{\mu}}) A^{a\bar{\nu}} A^{c\bar{\rho}}, \\
\bar{\mathcal{W}}_6^{a_1 a_2 a_3 a_4} &\equiv \int A_{\bar{\mu}}^{a_1} A^{a_2 \bar{\mu}} A_{\bar{\nu}}^{a_3} A^{a_4 \bar{\nu}} = \bar{\mathcal{W}}_6^{\{a_1 a_2\} \{a_3 a_4\}}, \\
\bar{\mathcal{W}}_7^{a_1 a_2 a_3 a_4} &\equiv \int \varepsilon_{\mu\nu\rho\delta} A^{a_1 \mu} A^{a_2 \nu} A^{a_3 \rho} A^{a_4 \delta}.
\end{aligned}$$

* b) *Monomials involving fermions and no external fields*

$$\begin{aligned}
\bar{\mathcal{W}}_8^{ij} &\equiv \int \frac{i}{2} \bar{\psi}_i \bar{\gamma}^\mu P_L \overset{\leftrightarrow}{\partial}_{\bar{\mu}} \psi_j, & \bar{\mathcal{W}}_9^{ij a} &\equiv \int \bar{\psi}_i \bar{\gamma}^\mu P_L \psi_j A_{\bar{\mu}}^a, \\
\bar{\mathcal{W}}_{10}^{ij} &\equiv \int \frac{i}{2} \bar{\psi}'_i \bar{\gamma}^\mu P_R \overset{\leftrightarrow}{\partial}_{\bar{\mu}} \psi'_j, & \bar{\mathcal{W}}_{11}^{ij a} &\equiv \int \bar{\psi}'_i \bar{\gamma}^\mu P_R \psi'_j A_{\bar{\mu}}^a, \\
\bar{\mathcal{W}}_{12}^{ij} &\equiv \int \frac{i}{2} \bar{\psi}_i \bar{\gamma}^\mu P_R \overset{\leftrightarrow}{\partial}_{\bar{\mu}} \psi_j, & \bar{\mathcal{W}}_{13}^{ij a} &\equiv \int \bar{\psi}_i \bar{\gamma}^\mu P_R \psi_j A_{\bar{\mu}}^a, \\
\bar{\mathcal{W}}_{14}^{ij} &\equiv \int \frac{i}{2} \bar{\psi}'_i \bar{\gamma}^\mu P_L \overset{\leftrightarrow}{\partial}_{\bar{\mu}} \psi'_j, & \bar{\mathcal{W}}_{15}^{ij a} &\equiv \int \bar{\psi}'_i \bar{\gamma}^\mu P_L \psi'_j A_{\bar{\mu}}^a.
\end{aligned}$$

* c) *Monomials involving ghost and no external fields*

$$\begin{aligned}
\bar{\mathcal{W}}_{16}^{ab} &\equiv \int -\bar{\omega}^a \bar{\square} \omega^b = \int (\partial_{\bar{\mu}} \bar{\omega}^a) (\partial^{\bar{\mu}} \omega^b), \\
\bar{\mathcal{W}}_{17}^{abc} &\equiv \int \bar{\omega}^a \partial^{\bar{\mu}} (\omega^c A_{\bar{\mu}}^b), & \bar{\mathcal{W}}_{18}^{abc} &\equiv \int \bar{\omega}^a (\partial^{\bar{\mu}} A^{b\bar{\mu}}) \omega^c, \\
\bar{\mathcal{W}}_{19}^{abcd} &\equiv \int \bar{\omega}^a \omega^b A^{c\bar{\mu}} A_{\bar{\mu}}^d, & \bar{\mathcal{W}}_{20}^{abcd} &\equiv \int \bar{\omega}^a \bar{\omega}^b \omega^c \omega^d.
\end{aligned}$$

* d) *Monomials involving external fields*

$$\begin{aligned}
\bar{\mathcal{W}}_{21}^{ij a} &\equiv \int \omega^a \bar{L}_i P_L \psi_j, & \bar{\mathcal{W}}_{22}^{ij a} &\equiv \int \bar{\psi}_i P_R \omega^a L_j, \\
\bar{\mathcal{W}}_{23}^{ij a} &\equiv \int \omega^a \bar{R}_i P_R \psi'_j, & \bar{\mathcal{W}}_{24}^{ij a} &\equiv \int \bar{\psi}'_i P_L \omega^a R_j, \\
\bar{\mathcal{W}}_{25}^{ij a} &\equiv \int \omega^a \bar{L}_i P_R \psi_j, & \bar{\mathcal{W}}_{26}^{ij a} &\equiv \int \bar{\psi}_i P_L \omega^a L_j, \\
\bar{\mathcal{W}}_{27}^{ij a} &\equiv \int \omega^a \bar{R}_i P_L \psi'_j, & \bar{\mathcal{W}}_{28}^{ij a} &\equiv \int \bar{\psi}'_i P_R \omega^a R_j, \\
\bar{\mathcal{W}}_{29}^{ab} &\equiv \int \rho_{\bar{\mu}}^a \partial^{\bar{\mu}} \omega^b, & \bar{\mathcal{W}}_{30}^{abc} &\equiv \int \rho_{\bar{\mu}}^a \omega^b A^{c\bar{\mu}}, \\
\bar{\mathcal{W}}_{31}^{abc} &\equiv \int \zeta^a \omega^b \omega^c.
\end{aligned}$$

Next, eq. (78) leads to

$$\mu \frac{\partial \Gamma_{\text{ren}}}{\partial \mu} = \sum_{i=1}^{31} \sum_{A_i} \bar{r}_i^{(1), A_i} \mathcal{W}_i^{A_i} + O(\hbar^2), \quad (85)$$

where A_i denotes the set of gauge indices, $\mathcal{W}_i^{A_i}$ denotes henceforth the counterpart of the operator $\bar{\mathcal{W}}_i^{A_i}$ in four dimensional space-time and the coefficients $\bar{r}_i^{(1), A_i}$ are defined as follows:

$$\bar{r}_1^{(1) ab} = \frac{(-i)^2}{2} \text{coef. in r.s.p. } (-i) \overline{\langle A_{\mu}^a(p_1) A_{\nu}^b(p_2 \equiv -p_1) \rangle}^{1\text{PI}(1)} \text{ of } \bar{p}_1^\mu \bar{p}_1^\nu =$$

$$\begin{aligned}
&= \frac{1}{(4\pi)^2} \delta^{ab} \left[T_{L+R} \frac{2}{3} - C_A \left(\frac{5}{3} + \frac{1-\alpha'}{2} \right) \right], \\
\bar{r}_2^{(1)ab} &= \frac{(-i)^2}{2} \text{coef. in r.s.p. } (-i) \overline{\langle A_\mu^a(p_1) A_\nu^b(p_2 \equiv -p_1) \rangle}^{1\text{PI}(1)} \text{ of } \bar{p}_1^2 \bar{g}^{\mu\nu} = \\
&= \frac{1}{(4\pi)^2} \delta^{ab} \left[T_{L+R} \frac{2}{3} - C_A \left(\frac{5}{3} + \frac{1-\alpha'}{2} \right) \right], \\
\bar{r}_3^{(1)abc} &= \frac{(-i)}{3} \text{coef. in r.s.p. } (-i) \overline{\langle A_\mu^a(p_1) A_\nu^b(p_2) A_\rho^c(p_3 \equiv -p_1 - p_2) \rangle}^{1\text{PI}(1)} \text{ of } \bar{p}_1^\nu \bar{g}^{\mu\rho} + \\
&\quad + \frac{(-i)}{3} \text{coef. in r.s.p. } (-i) \overline{\langle A_\mu^a(p_1) A_\nu^c(p_2) A_\rho^b(p_3 \equiv -p_1 - p_2) \rangle}^{1\text{PI}(1)} \text{ of } \bar{p}_1^\rho \bar{g}^{\mu\nu} = \\
&= \frac{1}{(4\pi)^2} \frac{2}{3} c^{abc} \left[\frac{2}{3} T_{L+R} - C_A \left(\frac{2}{3} + \frac{3(1-\alpha')}{4} \right) \right], \\
\bar{r}_4^{(1)abc} &= \frac{(-i)}{3} \text{coef. in r.s.p. } (-i) \overline{\langle A_\mu^a(p_1) A_\nu^b(p_2) A_\rho^c(p_3 \equiv -p_1 - p_2) \rangle}^{1\text{PI}(1)} \text{ of } \bar{p}_1^\mu \bar{g}^{\nu\rho} = \\
&= \frac{1}{(4\pi)^2} \frac{2}{3} c^{abc} \left[\frac{2}{3} T_{L+R} - C_A \left(\frac{2}{3} + \frac{3(1-\alpha')}{4} \right) \right], \\
\bar{r}_5^{(1)abc} &= \frac{(-i)}{3} \text{coef. in r.s.p. } (-i) \overline{\langle A_\mu^a(p_1) A_\nu^b(p_2) A_\rho^c(p_3 \equiv -p_1 - p_2) \rangle}^{1\text{PI}(1)} \text{ of } \bar{p}_1^\alpha \varepsilon_{\mu\nu\rho\alpha} = \\
&= 0, \\
\bar{r}_6^{(1)\{a_1 a_2\}\{a_3 a_4\}} &= r_6^{(1)\{a_3 a_4\}\{a_1 a_2\}} = \\
&= \frac{1}{8} \text{coef. in r.s.p. } (-i) \overline{\langle A_{\mu_1}^{a_1} A_{\mu_2}^{a_2} A_{\mu_3}^{a_3} A_{\mu_4}^{a_4} \rangle}^{1\text{PI}(1)} \text{ of } \bar{g}^{\mu_1 \mu_2} \bar{g}^{\mu_3 \mu_4} = \\
&= \frac{1}{(4\pi)^2} \frac{1}{8} \left[\frac{4}{3} T_{L+R} + 2 C_A \left(\frac{1}{3} - (1-\alpha') \right) \right] \\
&\quad (c^{ea_1 a_3} c^{ea_4 a_2} - c^{ea_1 a_4} c^{ea_2 a_3}), \tag{86} \\
\bar{r}_7^{(1)a_1 a_2 a_3 a_4} &= r_7^{(1)\{a_1 a_2 a_3 a_4\}} = \\
&= \frac{1}{4!} \text{coef. in r.s.p. } (-i) \overline{\langle A_{\mu_1}^{a_1} A_{\mu_2}^{a_2} A_{\mu_3}^{a_3} A_{\mu_4}^{a_4} \rangle}^{1\text{PI}(1)} \text{ of } \varepsilon_{\mu_1 \mu_2 \mu_3 \mu_4} = 0, \\
\bar{r}_8^{(1)ij} &= i \text{coef. in r.s.p. } \overline{\langle \psi_{\beta j}(-p) \bar{\psi}_{\alpha i}(p) \rangle}^{1\text{PI}(1)} \text{ of } (\bar{p} P_L)_{\alpha\beta} = \frac{i}{(4\pi)^2} 2g^2 \alpha' C_L, \\
\bar{r}_9^{(1)ij a} &= i \text{coef. in r.s.p. } (-i) \overline{\langle \psi_{\beta j} \bar{\psi}_{\alpha i} A_a^\mu \rangle}^{1\text{PI}(1)} \text{ of } (\bar{\gamma}^\mu P_L)_{\alpha\beta} = \\
&= \frac{1}{(4\pi)^2} 2g^2 C_L \left[\left(1 - \frac{1-\alpha'}{4} \right) C_A + \alpha' C_L \right] (T_L^a)_{ij}, \\
\bar{r}_{10}^{(1)ij} &= i \text{coef. in r.s.p. } \overline{\langle \psi'_{\beta j}(-p) \bar{\psi}'_{\alpha i}(p) \rangle}^{1\text{PI}(1)} \text{ of } (\bar{p} P_R)_{\alpha\beta} = \frac{i}{(4\pi)^2} 2g^2 \alpha' C_R, \\
\bar{r}_{11}^{(1)ij a} &= i \text{coef. in r.s.p. } (-i) \overline{\langle \psi'_{\beta j} \bar{\psi}'_{\alpha i} A_a^\mu \rangle}^{1\text{PI}(1)} \text{ of } (\bar{\gamma}^\mu P_R)_{\alpha\beta} = \\
&= \frac{1}{(4\pi)^2} 2g^2 C_R \left[\left(1 - \frac{1-\alpha'}{4} \right) C_A + \alpha' C_R \right] (T_R^a)_{ij},
\end{aligned}$$

$$\begin{aligned}
\bar{r}_{12}^{(1)ij} &= i \text{ coef. in r.s.p. } \overline{\langle \psi_{\beta j}(-p) \bar{\psi}_{\alpha i}(p) \rangle}^{1\text{PI}(1)} \text{ of } (\bar{p} P_R)_{\alpha\beta} = 0, \\
\bar{r}_{13}^{(1)ija} &= i \text{ coef. in r.s.p. } (-i) \overline{\langle \psi_{\beta j} \bar{\psi}_{\alpha i} A_a^\mu \rangle}^{1\text{PI}(1)} \text{ of } (\bar{\gamma}^\mu P_R)_{\alpha\beta} = 0, \\
\bar{r}_{14}^{(1)ij} &= i \text{ coef. in r.s.p. } \overline{\langle \psi'_{\beta j}(-p) \bar{\psi}'_{\alpha i}(p) \rangle}^{1\text{PI}(1)} \text{ of } (\bar{p} P_L)_{\alpha\beta} = 0, \\
\bar{r}_{15}^{(1)ija} &= i \text{ coef. in r.s.p. } (-i) \overline{\langle \psi'_{\beta j} \bar{\psi}'_{\alpha i} A_a^\mu \rangle}^{1\text{PI}(1)} \text{ of } (\bar{\gamma}^\mu P_L)_{\alpha\beta} = 0, \\
\bar{r}_{16}^{(1)ab} &= i \text{ coef. in r.s.p. } (-i) \overline{\langle \omega^b(-p) \bar{\omega}^a(-p) \rangle}^{1\text{PI}(1)} \text{ of } \bar{p}^2 = -\frac{1}{(4\pi)^2} C_A g^2 (1 + \frac{1-\alpha'}{2}), \\
\bar{r}_{17}^{(1)abc} &= (-i) \text{ coef. in r.s.p. } (-i) \overline{\langle \omega^c(-p_1 - p_2) \bar{\omega}^a(p_1) A_\mu^b(p_2) \rangle}^{1\text{PI}(1)} \text{ of } \bar{p}_1^\mu = \\
&= \frac{1}{(4\pi)^2} C_A g^2 \alpha' c^{abc}, \\
\bar{r}_{18}^{(1)abc} &= (-i) \text{ coef. in r.s.p. } (-i) \overline{\langle \omega^c(-p_1 - p_2) \bar{\omega}^a(p_1) A_\mu^b(p_2) \rangle}^{1\text{PI}(1)} \text{ of } \bar{p}_2^\mu = 0, \\
\bar{r}_{19}^{(1)abcd} &= \frac{1}{2} \text{ coef. in r.s.p. } (-i) \overline{\langle \omega^b \bar{\omega}^a A_\mu^b A_\nu^d \rangle}^{1\text{PI}(1)} \text{ of } \bar{g}^{\mu\nu} = 0, \\
\bar{r}_{20}^{(1)abcd} &= \frac{1}{4} \text{ r.s.p. } (-i) \overline{\langle \omega^d \omega^c \bar{\omega}^b \bar{\omega}^a \rangle}^{1\text{PI}(1)} = 0, \\
\bar{r}_{21}^{(1)ija} &= \text{coef. in r.s.p. } \overline{\langle \psi_{j\beta} \omega^a; s \psi_{i\alpha} \rangle}^{1\text{PI}(1)} \text{ of } (P_L)_{\alpha\beta} = \frac{i}{(4\pi)^2} g^2 \alpha' C_A (T_L^a)_{ij}, \\
\bar{r}_{22}^{(1)ija} &= \text{coef. in r.s.p. } \overline{\langle \omega^a \bar{\psi}_{i\alpha}; s \bar{\psi}_{j\beta} \rangle}^{1\text{PI}(1)} \text{ of } (P_R)_{\alpha\beta} = \frac{i}{(4\pi)^2} g^2 \alpha' C_A (T_L^a)_{ij}, \\
\bar{r}_{23}^{(1)ija} &= \text{coef. in r.s.p. } \overline{\langle \psi'_{j\beta} \omega^a; s \psi'_{i\alpha} \rangle}^{1\text{PI}(1)} \text{ of } (P_R)_{\alpha\beta} = \frac{i}{(4\pi)^2} g^2 \alpha' C_A (T_R^a)_{ij}, \\
\bar{r}_{24}^{(1)ija} &= \text{coef. in r.s.p. } \overline{\langle \omega^a \bar{\psi}'_{i\alpha}; s \bar{\psi}'_{j\beta} \rangle}^{1\text{PI}(1)} \text{ of } (P_L)_{\alpha\beta} = \frac{i}{(4\pi)^2} g^2 \alpha' C_A (T_R^a)_{ij}, \\
\bar{r}_{25}^{(1)ija} &= \text{coef. in r.s.p. } \overline{\langle \psi_{j\beta} \omega^a; s \psi_{i\alpha} \rangle}^{1\text{PI}(1)} \text{ of } (P_R)_{\alpha\beta} = 0 \\
\bar{r}_{26}^{(1)ija} &= \text{coef. in r.s.p. } \overline{\langle \omega^a \bar{\psi}_{i\alpha}; s \bar{\psi}_{j\beta} \rangle}^{1\text{PI}(1)} \text{ of } (P_L)_{\alpha\beta} = 0 \\
\bar{r}_{27}^{(1)ija} &= \text{coef. in r.s.p. } \overline{\langle \psi'_{j\beta} \omega^a; s \psi'_{i\alpha} \rangle}^{1\text{PI}(1)} \text{ of } (P_L)_{\alpha\beta} = 0 \\
\bar{r}_{28}^{(1)ija} &= \text{coef. in r.s.p. } \overline{\langle \omega^a \bar{\psi}'_{i\alpha}; s \bar{\psi}'_{j\beta} \rangle}^{1\text{PI}(1)} \text{ of } (P_R)_{\alpha\beta} = 0, \\
\bar{r}_{29}^{(1)ab} &= (-i) \text{ coef. in r.s.p. } \overline{\langle \omega^b(p_1); s A_\mu^a(p_2 = -p_1) \rangle}^{1\text{PI}(1)} \text{ of } \bar{p}_1^\mu = \\
&= -\frac{1}{(4\pi)^2} \delta^{ab} C_A g^2 (1 + \frac{1-\alpha'}{2}), \\
\bar{r}_{30}^{(1)abc} &= (-i) \text{ coef. in r.s.p. } \overline{\langle \omega^b A_\mu^c; s A_\nu^a \rangle}^{1\text{PI}(1)} \text{ of } \bar{g}_{\mu\nu} = -\frac{1}{(4\pi)^2} c^{abc} C_A g^2, \\
r_{31}^{(1)abc} &= \frac{1}{2} \text{ r.s.p. } \overline{\langle \omega^c \omega^b; s \omega^a \rangle}^{1\text{PI}(1)} = -\frac{1}{2} \frac{1}{(4\pi)^2} c^{abc} C_A g^2.
\end{aligned}$$

To remove much of the redundancy in the linear system in eq. (84) we shall rather express $\mu \frac{\partial \Gamma_{\text{ren}}}{\partial \mu}$ in terms of the functionals in eq. (71). By using the following equations

$$\begin{aligned}
S_{\text{cl } AA} &= \frac{\delta^{ab}}{2g^2} (\mathcal{W}_1^{ab} - \mathcal{W}_2^{ab}), & S_{\text{cl } AAA} &= -\frac{c^{abc}}{g^2} \mathcal{W}_3^{abc}, \\
S_{\text{cl } AAA} &= -\frac{1}{8g^2} (c_{ace}c_{bde} + c_{bce}c_{ade}) \mathcal{W}_6^{\{ab\}\{cd\}}, \\
S_{\text{cl } \bar{\psi}\psi}^{\text{L}} &= \delta_{ij} \mathcal{W}_8^{ij}, & S_{\text{cl } \bar{\psi}\psi A} &= (T_{\text{L}}^a)_{ij} \mathcal{W}_9^{ij}, \\
S_{\text{cl } \bar{\psi}'\psi'}^{\text{R}} &= \delta_{ij} \mathcal{W}_{10}^{ij}, & S_{\text{cl } \bar{\psi}'\psi' A} &= (T_{\text{R}}^a)_{ij} \mathcal{W}_{11}^{ij}, \\
S_{\text{cl } \bar{\omega}\omega} &= \delta^{ab} \mathcal{W}_{16}^{ab}, & S_{\text{cl } \bar{\omega}\omega A} &= -c^{abc} \mathcal{W}_{17}^{abc}, \\
S_{\text{cl } \bar{L}s\psi} &= i (T_{\text{L}}^a)_{ij} \mathcal{W}_{21}^{ij a}, & S_{\text{cl } Ls\bar{\psi}} &= i (T_{\text{L}}^a)_{ij} \mathcal{W}_{22}^{ij a}, \\
S_{\text{cl } \bar{R}s\psi'} &= i (T_{\text{R}}^a)_{ij} \mathcal{W}_{23}^{ij a}, & S_{\text{cl } Rs\bar{\psi}'} &= i (T_{\text{L(R)}}^a)_{ij} \mathcal{W}_{24}^{ij a}, \\
S_{\text{cl } \rho\omega} &= \delta^{ab} \mathcal{W}_{29}^{ab}, & S_{\text{cl } \rho\omega A} &= -c^{abc} \mathcal{W}_{30}^{abc}, \\
S_{\text{cl } \zeta\omega\omega} &= -\frac{1}{2} c^{abc} \mathcal{W}_{31}^{abc},
\end{aligned}$$

one easily obtains the contribution to $\mu \frac{\partial \Gamma_{\text{ren}}}{\partial \mu}$ at first order in $\hbar$:

$$\begin{aligned}
\mu \frac{\partial \Gamma_{\text{ren}}}{\partial \mu} &= \frac{1}{(4\pi)^2} g^2 \left[\frac{8}{3} \frac{T_{\text{L+R}}}{2} - C_A \left(\frac{10}{3} + (1 - \alpha') \right) \right] S_{\text{cl } AA} \\
&+ \frac{1}{(4\pi)^2} g^2 \left[\frac{8}{3} \frac{T_{\text{L+R}}}{2} - C_A \left(\frac{4}{3} + \frac{3}{2}(1 - \alpha') \right) \right] S_{\text{cl } AAA} \\
&+ \frac{1}{(4\pi)^2} g^2 \left[\frac{8}{3} \frac{T_{\text{L+R}}}{2} - C_A \left(-\frac{2}{3} + 2(1 - \alpha') \right) \right] S_{\text{cl } AAAA} \\
&+ \frac{1}{(4\pi)^2} g^2 C_{\text{L}} 2 \alpha' S_{\text{cl } \bar{\psi}\psi}^{\text{L}} + \frac{1}{(4\pi)^2} g^2 C_{\text{R}} 2 \alpha' S_{\text{cl } \bar{\psi}'\psi'}^{\text{R}} \\
&+ \frac{1}{(4\pi)^2} g^2 \left[\left(2 - \frac{(1 - \alpha')}{2} \right) C_A + 2 \alpha' C_{\text{L}} \right] S_{\text{cl } \bar{\psi}\psi A} \\
&+ \frac{1}{(4\pi)^2} g^2 \left[\left(2 - \frac{(1 - \alpha')}{2} \right) C_A + 2 \alpha' C_{\text{R}} \right] S_{\text{cl } \bar{\psi}'\psi' A} \\
&- \frac{1}{(4\pi)^2} g^2 \left(1 + \frac{(1 - \alpha')}{2} \right) C_A S_{\text{cl } \bar{\omega}\omega} + \frac{1}{(4\pi)^2} g^2 \alpha' C_A S_{\text{cl } \bar{\omega}\omega A} \\
&+ \frac{1}{(4\pi)^2} g^2 \alpha' C_A [S_{\text{cl } \bar{L}s\psi} + S_{\text{cl } \bar{R}s\psi'} + S_{\text{cl } Ls\bar{\psi}} + S_{\text{cl } Rs\bar{\psi}'}] \\
&- \frac{1}{(4\pi)^2} g^2 \left(1 + \frac{(1 - \alpha')}{2} \right) C_A S_{\text{cl } \rho\omega} + \frac{1}{(4\pi)^2} g^2 \alpha' C_A S_{\text{cl } \rho\omega A} \\
&+ \frac{1}{(4\pi)^2} g^2 \alpha' C_A S_{\text{cl } \zeta\omega\omega} + O(\hbar^2)
\end{aligned} \tag{87}$$

where, $T_{\text{L+R}}$ means sum over all left and right representations (*e.g.* number of “flavours”) in the theory.

Now, it is plain that eq. (77) reads

$$\begin{aligned} \mu \frac{\partial \Gamma_{\text{ren}}}{\partial \mu} &= -\beta^{(1)} g \frac{\partial S_{\text{cl}}}{\partial g} + \sum_{\phi} \gamma_{\phi}^{(1)} \mathcal{N}_{\phi} S_{\text{cl}} + O(\hbar^2) = \\ &= -\beta^{(1)} L_g + \sum_{\phi} \gamma_{\phi}^{(1)} L_{\phi} + O(\hbar^2), \end{aligned} \quad (88)$$

which in turn can be expressed in terms of the functionals $S_{\text{cl}} \dots$ given in eq. (71).

Finally, the fact that the left hand sides of eqs. (87) and (88) should match gives rise to the following system of equations:

$$\begin{aligned} * \text{ From } S_{\text{cl AA}} &\rightarrow 2\beta^{(1)} + 2\gamma_A^{(1)} = \frac{1}{(4\pi)^2} g^2 \left[\frac{8}{3} \frac{T_{\text{L+R}}}{2} - C_A \left(\frac{10}{3} + (1 - \alpha') \right) \right], \\ * \text{ From } S_{\text{cl AAA}} &\rightarrow 2\beta^{(1)} + 3\gamma_A^{(1)} = \frac{1}{(4\pi)^2} g^2 \left[\frac{8}{3} \frac{T_{\text{L+R}}}{2} - C_A \left(\frac{4}{3} + \frac{3}{2}(1 - \alpha') \right) \right], \\ * \text{ From } S_{\text{cl AAAA}} &\rightarrow 2\beta^{(1)} + 4\gamma_A^{(1)} = \frac{1}{(4\pi)^2} g^2 \left[\frac{8}{3} \frac{T_{\text{L+R}}}{2} - C_A \left(-\frac{2}{3} + 2(1 - \alpha') \right) \right], \\ * \text{ From } S_{\text{cl } \bar{\psi}\psi}^{\text{L}} &\rightarrow 2\gamma_{\psi}^{\text{L}(1)} = \frac{1}{(4\pi)^2} g^2 C_{\text{L}} 2\alpha', \\ * \text{ From } S_{\text{cl } \bar{\psi}'\psi'}^{\text{R}} &\rightarrow 2\gamma_{\psi'}^{\text{R}(1)} = \frac{1}{(4\pi)^2} g^2 C_{\text{R}} 2\alpha', \\ * \text{ From } S_{\text{cl } \bar{\psi}\psi_A}^{\text{L}} &\rightarrow 2\gamma_{\psi}^{\text{L}(1)} + \gamma_A^{(1)} = \frac{1}{(4\pi)^2} g^2 \left[\left(2 - \frac{(1-\alpha')}{2} \right) C_A + 2\alpha' C_{\text{L}} \right], \\ * \text{ From } S_{\text{cl } \bar{\psi}'\psi'_A}^{\text{R}} &\rightarrow 2\gamma_{\psi'}^{\text{R}(1)} + \gamma_A^{(1)} = \frac{1}{(4\pi)^2} g^2 \left[\left(2 - \frac{(1-\alpha')}{2} \right) C_A + 2\alpha' C_{\text{R}} \right], \\ * \text{ From } S_{\text{cl } \bar{\omega}\omega} &\rightarrow -\gamma_A^{(1)} + \gamma_{\omega}^{(1)} = -\frac{1}{(4\pi)^2} g^2 \left(1 + \frac{(1-\alpha')}{2} \right) C_A, \\ * \text{ From } S_{\text{cl } \bar{\omega}\omega_A} &\rightarrow \gamma_{\omega}^{(1)} = \frac{1}{(4\pi)^2} g^2 \alpha' C_A, \\ * \text{ From } S_{\text{cl } \bar{\text{L}}\text{S}\psi} &\rightarrow \gamma_{\omega}^{(1)} = \frac{1}{(4\pi)^2} g^2 \alpha' C_A, \\ * \text{ From } S_{\text{cl } \bar{\text{R}}\text{S}\psi'} &\rightarrow \gamma_{\omega}^{(1)} = \frac{1}{(4\pi)^2} g^2 \alpha' C_A, \\ * \text{ From } S_{\text{cl } \text{L}\text{S}\bar{\psi}} &\rightarrow \gamma_{\omega}^{(1)} = \frac{1}{(4\pi)^2} g^2 \alpha' C_A, \\ * \text{ From } S_{\text{cl } \text{R}\text{S}\bar{\psi}'} &\rightarrow \gamma_{\omega}^{(1)} = \frac{1}{(4\pi)^2} g^2 \alpha' C_A, \\ * \text{ From } S_{\text{cl } \rho\omega} &\rightarrow -\gamma_A^{(1)} + \gamma_{\omega}^{(1)} = -\frac{1}{(4\pi)^2} g^2 \left(1 + \frac{(1-\alpha')}{2} \right) C_A, \\ * \text{ From } S_{\text{cl } \rho\omega_A} &\rightarrow \gamma_{\omega}^{(1)} = \frac{1}{(4\pi)^2} g^2 \alpha' C_A, \\ * \text{ From } S_{\text{cl } \zeta\omega\omega} &\rightarrow \gamma_{\omega}^{(1)} = \frac{1}{(4\pi)^2} g^2 \alpha' C_A; \end{aligned} \quad (89)$$

which is a simplified version of eq. (84). This simplification takes place since the non-symmetrical finite counterterms, $S_{\text{fct}}^{(n)}$, needed to restore the BRS symmetry and the evanescent monomials, $\hat{\mathcal{W}}_j$, only begin to contribute to the renormalization group equation at order $\hbar^2$. At order $\hbar^n$, $n > 1$, it is the linear system in eq. (83) which is to be dealt with. The system in eq. (89) is compatible and overdeterminate and its solution reads:

$$\beta^{(1)} = \frac{1}{(4\pi)^2} g^2 \left(\frac{4}{3} \frac{T_{\text{L+R}}}{2} - \frac{11}{3} C_A \right),$$

$$\begin{aligned}
\gamma_A^{(1)} &= \frac{1}{(4\pi)^2} g^2 \left(2 - \frac{1 - \alpha'}{2} \right) C_A, \\
\gamma_\omega^{(1)} &= \frac{1}{(4\pi)^2} g^2 \alpha' C_A, \\
\gamma_\psi^L{}^{(1)} &= \frac{1}{(4\pi)^2} g^2 \alpha' C_L, \quad \gamma_{\psi'}^R{}^{(1)} = \frac{1}{(4\pi)^2} g^2 \alpha' C_R.
\end{aligned} \tag{90}$$

The reader should notice that to actually compute the beta function and anomalous dimensions of the theory one does not need to evaluate the thirtyone coefficients in eq. (85) (computed in eq. (86)) but just a few of them, if appropriately chosen. Indeed, the computation of, say, $\bar{r}_1^{(1)ab}$, $\bar{r}_8^{(1)ij}$, $\bar{r}_{10}^{(1)ij}$, $\bar{r}_{16}^{(1)ab}$ and $\bar{r}_{17}^{(1)abc}$ would do the job. However, a thorough check of the formalism demands the computation of the thirtyone coefficients in question. And this we have done.

We have finished the computation of the renormalization group equation at order $\hbar$ for the theory with classical action in eq. (27). Our expansions were expansions in $\hbar$, as suits the cohomology problems which arise in connection with the BRS symmetry, rather than in the coupling constant g . Our parametrizations of the wave functions were such that g only occurs, in the classical action, in the Yang-Mills term, so that $g\partial_g S_{\text{cl}}$ is the only element (modulo multiplication by a constant) of the local cohomology of the BRS operator b (see eq. (29)) over the space of local integrated functionals of ghost number 0 and ultraviolet dimension 4 (Lorentz invariance and rigid gauge symmetry are also assumed). The other contributions to the renormalization group equation are b -exact and have to do with the anomalous dimensions of the theory.

It is often the case that one chooses a parametrization of the wave function such that, at the tree-level, the fermionic and the three-boson vertices carry the coupling factor g , and the four-boson vertex is proportional to g^2 . The renormalization group equation for this parametrization of the wave function is easily retrieved from our results. Let us denote by $\Gamma_{\text{ren}}^*[A^*, \omega^*, \bar{\omega}^*, B^*, \rho^*, \zeta^*, \psi, \bar{\psi}, \psi', \bar{\psi}', \alpha^*, g]$ the 1PI functional for this new parametrization. Then,

$$\begin{aligned}
&\Gamma_{\text{ren}}^*[A^*, \omega^*, \bar{\omega}^*, B^*, \rho^*, \zeta^*, \psi, \bar{\psi}, \psi', \bar{\psi}', \alpha^*, g] \equiv \\
&\Gamma_{\text{ren}}[gA^*, g\omega^*, g^{-1}\bar{\omega}^*g^{-1}, g^{-1}B^*, g^{-1}\rho^*, g^{-1}\zeta^*, \psi, \bar{\psi}, \psi', \bar{\psi}', g^2\alpha^*, g] \equiv \\
&\Gamma_{\text{ren}}[A, \omega, \bar{\omega}, B, \rho, \zeta, \psi, \bar{\psi}, \psi', \bar{\psi}', \alpha, g],
\end{aligned}$$

where $\Gamma_{\text{ren}}[A, \omega, \bar{\omega}, B, \rho, \zeta, \psi, \bar{\psi}, \psi', \bar{\psi}', \alpha, g]$ is the 1PI functional for the parametrization of eq. (27). Now, since

$$\begin{aligned}
g \frac{\partial \Gamma_{\text{ren}}}{\partial g} &= \frac{\partial \Gamma_{\text{ren}}^*}{\partial g} - \frac{1}{g} N_{A^*} \Gamma_{\text{ren}}^* - N_{\omega^*} \Gamma_{\text{ren}}^* + N_{\bar{\omega}^*} \Gamma_{\text{ren}}^* + N_{B^*} \Gamma_{\text{ren}}^* \\
&\quad + N_{\rho^*} \Gamma_{\text{ren}}^* + N_{\zeta^*} \Gamma_{\text{ren}}^* - 2\alpha^* \frac{\partial \Gamma_{\text{ren}}^*}{\partial \alpha^*},
\end{aligned}$$

the renormalization group equation eq. (74) becomes (in terms of the usual combination $\alpha_S \equiv g^2/(4\pi)$):

$$\left[\mu \frac{\partial}{\partial \mu} + \beta_S \alpha_S \frac{\partial}{\partial \alpha_S} + \delta_{\alpha^*} \alpha^* \frac{\partial}{\partial \alpha^*} - \gamma_\psi^L \mathcal{N}_\psi^L - \gamma_{\psi'}^R \mathcal{N}_{\psi'}^R - \gamma_{A^*} N_{A^*} - \gamma_{\omega^*} N_{\omega^*} \right]$$

$$- \gamma_{\bar{\omega}^*} N_{\bar{\omega}^*} - \gamma_{B^*} N_{B^*} - \gamma_{\rho^*} N_{\rho^*} - \gamma_{\zeta^*} N_{\zeta^*} \Big] \Gamma_{\text{ren}}^* = 0,$$

with

$$\begin{aligned} \beta_S &= 2\beta, \\ \gamma_{A^*} &= -\gamma_{\rho^*} = -\gamma_{\bar{\omega}^*} = -\gamma_{B^*} = \gamma_A + \beta, \\ \gamma_{\omega^*} &= -\gamma_{\zeta^*} = \gamma_{\omega} + \beta, \\ \delta_{\alpha^*} &= -2\gamma_{A^*} = -2(\gamma_A + \beta). \end{aligned}$$

finally, by using the results in eq. (90), one obtains

$$\begin{aligned} \beta_S^{(1)} &= \frac{\alpha_S}{\pi} \left[\frac{2}{3} \frac{T_{L+R}}{2} - \frac{11}{6} C_A \right], \\ \gamma_{A^*}^{(1)} &= \frac{\alpha_S}{\pi} \left[\frac{1}{3} \frac{T_{L+R}}{2} - \frac{C_A}{4} \left(\frac{13}{6} - \frac{\alpha^*}{2} \right) \right], \\ \gamma_{\omega^*}^{(1)} &= \frac{\alpha_S}{\pi} \left[\frac{1}{3} \frac{T_{L+R}}{2} - \frac{C_A}{4} \left(\frac{31}{6} + \frac{\alpha^*}{2} \right) \right]. \end{aligned}$$

4. Chiral Yang-Mills theories: non-simple gauge groups

4.1. The tree level action

In this section, the gauge group will be a compact Lie group which is the direct product of N_S simple groups and N_A Abelian factors. Its (real) Lie algebra is a direct sum: $\mathbf{G} = \mathbf{G}_1 \oplus \cdots \oplus \mathbf{G}_{N_A} \oplus \mathbf{G}_{N_A+1} \oplus \cdots \oplus \mathbf{G}_{N_A+N_S}$ of dimension $d_{\mathbf{G}} = N_A + d_{\mathbf{G}_{N_A+1}} + \cdots + d_{\mathbf{G}_{N_A+N_S}}$. The index G , which labels the group factors, will run from 1 to $N_A + N_S$, the index $\mathcal{A}$, which labels the Abelian factors, from 1 to N_A and the index $\mathcal{S}$, which labels the simple factors, from 1 to N_S .

Therefore, there exists a basis for the Lie algebra which can be split and enumerated as follows:

$$\begin{aligned} \{X_a\}_{a=1}^{d_{\mathbf{G}}} &= \{X_1, \dots, X_{N_A}, X_{N_A+1}, \dots, X_{N_A+d_{\mathbf{G}_1}}, \dots\} \\ &\equiv \{X_1, \dots, X_{N_A}\} \cup \left(\bigcup_{\mathcal{S}=1}^{N_S} \{X_{a_{\mathcal{S}}}^{(\mathcal{S})}\}_{a_{\mathcal{S}}=1}^{d_{\mathbf{G}_{N_A+\mathcal{S}}}} \right); \end{aligned}$$

and such that the Killing form of the semisimple part is diagonal and positive definite; so, we choose δ_a^b to lower and rise indices. For any such basis, the structure constants, given by $[X_a, X_b] = i c_{ab}^c X_c$, are completely antisymmetric, and $c_{a_G de} c_{b_{G^*} de} = 0$ if $G \neq G^*$, $c_{a_G de} c_{b_G de} = c_{a_G d_G e_G} c_{b_G d_G e_G} = C_A^{(G)} \delta_{ab}$, with $C_A^{(\mathcal{A})} \equiv 0$ (clearly, $c_{a_A bc} = 0$).

Let us display the field content of the theory and introduce the Dimensional Regularization classical action, S_0 . We have first the Lie-algebra-valued fields. The pure Yang-Mills part of the tree-level action in the d -dimensional space of Dimensional Regularization reads:

$$S_{\text{YM}} = \int d^d x \quad -\frac{1}{4} F_{\mu\nu}^a C_{ab} F^{b\mu\nu} \quad (91)$$

where C is a diagonal matrix with

$$\{g_1^{-2}, \dots, g_{N_A}^{-2}, \underbrace{g_{N_A+1}^{-2}, \dots, g_{N_A+1}^{-2}}_{d_{G_{N_A+1}} \text{ times}}, \dots, \underbrace{g_{N_A+N_S}^{-2}, \dots, g_{N_A+N_S}^{-2}}_{d_{G_{N_A+N_S}} \text{ times}}\}$$

in the diagonal. Hence,

$$S_{\text{YM}} = \int d^d x \sum_G -\frac{1}{4g_G^2} F_{\mu\nu}^{(G)a_G} F_{a_G \mu\nu}^{(G)},$$

where $F_{\mu\nu}^{(G)a_G} = \partial_\mu A_\nu^{(G)a_G} - \partial_\nu A_\mu^{(G)a_G} + c^{a_G b_G c_G} A_\mu^{(G)b_G} A_\nu^{(G)c_G}$. Obviously, all the remaining Lie algebra-valued fields are indexed in the same way yielding:

$$\begin{aligned} S_{\text{gf}} &= \int d^d x \frac{1}{2} B_a \Lambda^{ab} B_b + B_a (\partial^\mu A_\mu^a) - \bar{\omega}_a \partial^\mu \nabla_\mu \omega^a, \\ &= \int d^d x \sum_G \frac{\alpha_G}{2} B^{(G)a_G} B_{a_G}^{(G)} + B_{a_G}^{(G)} (\partial^\mu A_\mu^{(G)a_G}) - \bar{\omega}_{a_G}^{(G)} \partial^\mu \nabla_\mu \omega^{(G)a_G}. \end{aligned}$$

The free boson propagator reads now:

$$\left(C^{-1} \frac{-i}{k^2 + i\epsilon} \left[(g^{\mu\nu} - \frac{k^\mu k^\nu}{k^2}) + C \Lambda \frac{k^\mu k^\nu}{k^2} \right] \right)^{ab};$$

i.e. diagonal but no longer multiple of the identity.

Notice that we have also introduced ghosts to control the Abelian variations. Since ghosts in Abelian theories do not interact with the rest of fields ($\nabla_\mu \omega^A = \partial_\mu \omega^A$), they can occur in a non-tree-level 1PI diagram only as a part of a vertex insertion (BRS-variations and breakings); so that, by differentiating with respect to ω^A the Slavnov-Taylor identities, the usual Abelian Ward identities are easily recovered. But we insist on keeping the Abelian ghost in order to have an unified notation.

The matter content of the theory will be only non-Majorana fermions. The corresponding fermion fields, $\psi(\psi')$, carrying a left-handed (right-handed) fully reducible representation of the gauge group, so that it can be descomposed in a sum of irreducible representations. From now on, until otherwise stated, we will take both the left-handed and right-handed fermionic to be irreducible, the general case just involve a sum over all the irreducible representations.

We shall index the fermion fields carrying the irreducible representation of the gauge Lie algebra as follows $\psi^{i_1, \dots, i_{N_S}}$ ($\psi'^{i_1, \dots, i_{N_S}}$), with i_S running from 1 to $d_S(d'_S)$; so that the generators of the Lie algebra are given by

$$\{T_{L(R)}^A\}_{A=1}^{N_A} = \{t_{L(R)}^A\}_{A=1}^{N_A} \otimes 1_{d_1^{(\prime)}} \cdots \otimes 1_{d_{N_S}^{(\prime)}},$$

where each $t_{L(R)}^A$ is a real number and

$$\{T_{\text{L(R)}}^a\}_{a=N_{\mathcal{A}}+1}^{N_{\mathcal{A}}+d_{\mathbf{G}}} = 1 \otimes \{1_{d_1^{(\prime)}} \otimes \cdots \otimes \{T_{\text{L(R)}}^{(\mathcal{S}) a_{\mathcal{S}}}\}_{a_{\mathcal{S}}=1}^{d_{\mathbf{G}_{N_{\mathcal{A}}+\mathcal{S}}}} \otimes \cdots 1_{d_{N_{\mathcal{S}}}^{(\prime)}}\}_{\mathcal{S}=1}^{N_{\mathcal{S}}},$$

where $T_{\text{L(R)}}^{(\mathcal{S}) a_{\mathcal{S}}} \in \text{M}_{d_{\mathcal{S}}^{(\prime)}}(\mathbb{C})$ are complex matrices of dimension $d_{\mathcal{S}}^{(\prime)}$ which furnish an irreducible representation of dimension $d_{\mathcal{S}}^{(\prime)}$ of the Lie algebra $\mathbf{G}_{N_{\mathcal{A}}+\mathcal{S}}$, $\mathcal{S} = 1 \cdots N_{\mathcal{S}}$.

The BRS variations of the fermion fields in the d -dimensional space of Dimensional Regularization are defined to be

$$\begin{aligned} s\psi^{(\prime)} &= i\omega^a T_{\text{L(R)}}^a P_{\text{L(R)}} \psi^{(\prime)} = \\ &= \sum_{\mathcal{A}=1}^{N_{\mathcal{A}}} i\omega^{\mathcal{A}} t_{\text{L(R)}}^{\mathcal{A}} 1_{d_1^{(\prime)}} \otimes \cdots \otimes 1_{d_{N_{\mathcal{S}}}^{(\prime)}} P_{\text{L(R)}} \psi^{(\prime)} + \\ &\quad + \sum_{\mathcal{S}=1}^{N_{\mathcal{S}}} i \sum_{a_{\mathcal{S}}=1}^{d_{\mathbf{G}_{N_{\mathcal{A}}+\mathcal{S}}}} \omega^{(\mathcal{S}) a_{\mathcal{S}}} 1_{d_1^{(\prime)}} \otimes \cdots \otimes T_{\text{L(R)}}^{(\mathcal{S}) a_{\mathcal{S}}} \otimes \cdots \otimes 1_{d_{N_{\mathcal{S}}}^{(\prime)}} P_{\text{L(R)}} \psi^{(\prime)}, \end{aligned}$$

and the regularized interaction is introduced in the same way.

The following results will be useful

$$\begin{aligned} \text{Tr}_{\text{M}_{d_{\mathcal{S}}^{(\prime)}}(\mathbb{C})} [T_{\text{L(R)}}^{(\mathcal{S}) a_{\mathcal{S}}} T_{\text{L(R)}}^{(\mathcal{S}) b_{\mathcal{S}}}] &\equiv T_{\text{L(R)}}^{\mathcal{S}} \delta^{a_{\mathcal{S}} b_{\mathcal{S}}}, \quad \text{Tr}_{\text{M}_{d_{\mathcal{S}}^{(\prime)}}(\mathbb{C})} [T_{\text{L(R)}}^{(\mathcal{S}) a_{\mathcal{S}}}] = 0, \\ \text{Tr} [T_{\text{L(R)}}^a T_{\text{L(R)}}^b] &= \begin{cases} \left(\prod_{K \neq \mathcal{S}}^{N_{\mathcal{S}}} d_K^{(\prime)} \right) T_{\text{L(R)}}^{\mathcal{S}} \delta^{ab}, & \text{if } a = a_{\mathcal{S}} \text{ and } b = b_{\mathcal{S}}, \text{ with } \mathcal{S} = 1, \dots, N_{\mathcal{S}} \\ \left(\prod_{\mathcal{S}=1}^{N_{\mathcal{S}}} d_{\mathcal{S}}^{(\prime)} \right) t_{\text{L(R)}}^a t_{\text{L(R)}}^b, & \text{if } a \leq N_{\mathcal{A}} \text{ and } b \leq N_{\mathcal{A}} \\ 0 & \text{in the remainder} \end{cases} \\ \sum_{e_{\mathcal{S}}}^{d_{\mathcal{S}}^{(\prime)}} T_{\text{L(R)}}^{(\mathcal{S}) e_{\mathcal{S}}} T_{\text{L(R)}}^{(\mathcal{S}) e_{\mathcal{S}}} &\equiv C_{\text{L(R)}}^{(\mathcal{S})} 1_{d_{\mathcal{S}}^{(\prime)}}, \quad (t_{\text{L(R)}}^{\mathcal{A}})^2 \equiv C_{\text{L(R)}}^{(\mathcal{A})} \\ \sum_a T_{\text{L(R)}}^a T_{\text{L(R)}}^a &= \sum_{G=1}^{N_{\mathcal{A}}+N_{\mathcal{S}}} C_{\text{L(R)}}^{(G)} 1_{d_1^{(\prime)}} \otimes \cdots \otimes 1_{d_{N_{\mathcal{S}}}^{(\prime)}} \equiv \sum_G C_{\text{L(R)}}^{(G)} \\ \sum_{a,b} T_{\text{L(R)}}^a (C^{-1})_b^a T_{\text{L(R)}}^b &= \sum_{G=1}^{N_{\mathcal{A}}+N_{\mathcal{S}}} g_G^2 C_{\text{L(R)}}^{(G)} \equiv C_{\text{L(R)}}^g, \\ \sum_{a,b} T_{\text{L(R)}}^a \Lambda_{ab} T_{\text{L(R)}}^b &= \sum_{G=1}^{N_{\mathcal{A}}+N_{\mathcal{S}}} \alpha_G C_{\text{L(R)}}^{(G)} \equiv C_{\text{L(R)}}^{\alpha}. \end{aligned}$$

4.2. The one-loop singular counterterms

To obtain the one-loop singular counterterms, it is necessary to make in the non-hatted terms in eq. (37) the following substitutions

$$\begin{aligned}
g^2 C_A S_{0X\dots XY\dots Y} &\longrightarrow \sum_S g_S^2 C_A^{(S)} S_{0X\dots XY_S\dots Y_S}, \\
g^2 \alpha' C_A S_{0X\dots XY\dots Y} &\longrightarrow \sum_S g_S^2 \alpha'^S C_A^{(S)} S_{0X\dots XY_S\dots Y_S}, \\
g^2 C_{L(R)} &\longrightarrow C_{L(R)}^g, \\
g^2 C_{L(R)} \alpha' &\longrightarrow C_{L(R)}^\alpha, \\
g^2 T_{L(R)} S_{0A\dots A} &\longrightarrow \sum_{A=1}^{N_A} g_A^2 \left(\prod_{S=1}^{N_S} d_S^{(\prime)} \right) (t_{L(R)}^A)^2 S_{0A_A\dots A_A} \\
&\quad + \sum_{S=1}^{N_S} g_{N_A+S}^2 \left(\prod_{K \neq S}^{N_S} d_K^{(\prime)} \right) T_{L(R)S} S_{0A_S\dots A_S}.
\end{aligned}$$

Above, the symbol Y stands for any Lie-algebra-valued field. A new type of term should also be added. This term is the Abelian mixing term, which is given by

$$\frac{\hbar}{\varepsilon} \frac{1}{(4\pi)^2} \sum_{\substack{A_1, A_2 \\ A_1 \neq A_2}}^{N_A} \left(\prod_{S=1}^{N_S} d_S^{(\prime)} \right) t_{L(R)}^{A_1} t_{L(R)}^{A_2} \int d^d x - \frac{1}{4} (\partial_\mu A_\nu^{A_1} - \partial_\nu A_\mu^{A_1}) (\partial_\mu A_\nu^{A_2} - \partial_\nu A_\mu^{A_2}).$$

The last line of eq. (37), *i.e.* the hatted contribution, has to be modified in the same manner.

The reader should bear in mind that $g^2 C_{L(R)} \alpha' S_{0\bar{\psi}^{(\prime)} \psi^{(\prime)} A} \rightarrow C_{L(R)}^\alpha \sum_G S_{0\bar{\psi}^{(\prime)} \psi^{(\prime)} A_G}$.

4.3. Expression of the breaking

To obtain the generalization of eq. (64) to current case, one applies first the same kind of modifications as in the previous subsection. In addition, the following replacements should be made

$$T_{L(R)} c^{abc} \mathcal{M}_4^{abc} \longrightarrow \sum_{S=1}^{N_S} \left(\prod_{K \neq S}^{N_S} d_K^{(\prime)} \right) T_{L(R)S} c^{asbscs} \mathcal{M}_4^{asbscs},$$

and similar substitutions for the terms of the same type as $T_{L(R)} c^{ebc} c^{eda} \mathcal{M}^{abcd}$.

The traces of the product of three and four generators, which appear for example in the expression of anomaly, keep its form, but with the understanding that Tr is a matrix trace over the full fermion representation of the Gauge Lie algebra $\mathbf{G}$.

4.4. Restoration of Slavnov-Taylor Identities

Apart from modifications of the same type as describe above, we have to be careful with the resolution of the equations involving $\mathcal{M}_1^{ab}$ (and also $\mathcal{M}_7$ and $\mathcal{M}_8$, but they actually do not give rise to any new result). Now, we substitute in eq. (67) $a_1 \tilde{\Delta}_i \rightarrow a_1^G \tilde{\Delta}_i^G + a_1^{\mathcal{A}_1 \mathcal{A}_2} \tilde{\Delta}_i^{\mathcal{A}_1 \mathcal{A}_2}$, $i = 1, 2$, $\mathcal{A}_1 \neq \mathcal{A}_2$, with the obvious meaning.

Apart from the equation in $\mathcal{M}_1^{a_G b_G}$, which will give together with the rest of the system an unique solution for

$$\begin{aligned} a_1^{\mathcal{A}} &= \frac{1}{(4\pi)^2} \frac{5}{12} \left[\left(\prod_{S=1}^{N_S} d_S \right) (t_L^{\mathcal{A}})^2 + \left(\prod_{S=1}^{N_S} d'_S \right) (t_R^{\mathcal{A}})^2 \right], \\ a_1^{\mathcal{S}} &= \frac{1}{(4\pi)^2} \frac{5}{12} \left[\left(\prod_{K \neq \mathcal{S}} d_K \right) T_{LS} + \left(\prod_{K \neq \mathcal{S}} d'_K \right) T_{RS} \right], \\ a_2^{\mathcal{A}} &= \frac{1}{(4\pi)^2} \frac{1}{4} \left[\left(\prod_{S=1}^{N_S} d_S \right) (t_L^{\mathcal{A}})^2 + \left(\prod_{S=1}^{N_S} d'_S \right) (t_R^{\mathcal{A}})^2 \right], \\ a_2^{\mathcal{S}} &= \frac{1}{(4\pi)^2} \frac{1}{4} \left[\left(\prod_{K \neq \mathcal{S}} d_K \right) T_{LS} + \left(\prod_{K \neq \mathcal{S}} d'_K \right) T_{RS} \right], \end{aligned}$$

we have new equations in $\mathcal{M}_1^{\mathcal{A}_1 \mathcal{A}_2}$ and $\mathcal{M}_1^{\mathcal{A}_2 \mathcal{A}_1}$, $\mathcal{A}_1 \neq \mathcal{A}_2$:

$$\begin{aligned} a_1^{\mathcal{A}_1 \mathcal{A}_2} - a_2^{\mathcal{A}_1 \mathcal{A}_2} &= \frac{1}{(4\pi)^2} \frac{1}{3} \left[\left(\prod_{S=1}^{N_S} d_S \right) t_L^{\mathcal{A}_1} t_L^{\mathcal{A}_2} + \left(\prod_{S=1}^{N_S} d'_S \right) t_R^{\mathcal{A}_1} t_R^{\mathcal{A}_2} \right] \\ a_1^{\mathcal{A}_2 \mathcal{A}_1} - a_2^{\mathcal{A}_2 \mathcal{A}_1} &= \frac{1}{(4\pi)^2} \frac{1}{3} \left[\left(\prod_{S=1}^{N_S} d_S \right) t_L^{\mathcal{A}_1} t_L^{\mathcal{A}_2} + \left(\prod_{S=1}^{N_S} d'_S \right) t_R^{\mathcal{A}_1} t_R^{\mathcal{A}_2} \right] \end{aligned}$$

whose solution is not unique. This was expected because we have now a new invariant: $F_{\mu\nu}^{\mathcal{A}_1} F^{\mathcal{A}_2 \mu\nu}$, $\mathcal{A}_1 \neq \mathcal{A}_2$, mixing Abelian bosons. We can impose, for example, $a_2^{\mathcal{A}_1 \mathcal{A}_2} \equiv 0$.

Therefore, from eq. (69), with the help of the modifications given above (do not forget the new terms), one obtains following result:

$$\begin{aligned} S_{\text{fct}}^{(1)} &= \frac{\hbar^1}{(4\pi)^2} \int \sum_{\mathcal{A}=1}^{N_A} \left[\left(\prod_{S=1}^{N_S} d_S \right) (t_L^{\mathcal{A}})^2 + \left(\prod_{S=1}^{N_S} d'_S \right) (t_R^{\mathcal{A}})^2 \right] \left[\frac{5}{12} (\partial^\mu A_\mu^{\mathcal{A}})^2 + \frac{1}{4} A_\mu^{\mathcal{A}} \square A^{\mathcal{A} \mu} \right] \\ &+ \sum_{S=1}^{N_S} \left[\left(\prod_{K \neq \mathcal{S}} d_K \right) T_{LS} + \left(\prod_{K \neq \mathcal{S}} d'_K \right) T_{RS} \right] \left[\frac{5}{12} (\partial^\mu A_\mu^{(\mathcal{S}) a_S}) (\partial^\nu A_\nu^{(\mathcal{S}) a_S}) + \frac{1}{4} A_\mu^{(\mathcal{S}) a_S} \square A^{(\mathcal{S}) a_S \mu} \right] \\ &- \sum_{S=1}^{N_S} \left[\left(\prod_{K \neq \mathcal{S}} d_K \right) \frac{T_{LS}}{6} + \left(\prod_{K \neq \mathcal{S}} d'_K \right) \frac{T_{RS}}{6} \right] c^{a_S b_S c_S} (\partial_\mu A_\nu^{(\mathcal{S}) a_S}) A^{(\mathcal{S}) b_S \mu} A^{(\mathcal{S}) c_S \nu} \\ &+ \frac{\text{Tr}_{\text{full}}[T_L^a T_L^b T_L^c T_L^d] + \text{Tr}_{\text{full}}[T_R^a T_R^b T_R^c T_R^d]}{12} A_\mu^a A^{b\mu} A_\nu^c A^{d\nu} \end{aligned}$$

$$\begin{aligned}
& + \sum_{G=1}^{N_A+N_S} g_G^2 C_L^{(G)} \left[1 + (\alpha'^{(G)} - 1)/6 \right] \frac{i}{2} \bar{\psi} \bar{\gamma}^\mu P_L \overleftrightarrow{\partial}_\mu \psi \\
& + \sum_{G=1}^{N_A+N_S} g_G^2 C_R^{(G)} \left[1 + (\alpha'^{(G)} - 1)/6 \right] \frac{i}{2} \bar{\psi}' \bar{\gamma}^\mu P_R \overleftrightarrow{\partial}_\mu \psi' \\
& - \sum_{S=1}^{N_S} g_S^2 \frac{C_A^{(S)}}{4} [1 + (\alpha'^{(S)} - 1)] (\bar{L}s\psi + Ls\bar{\psi} + \bar{R}s\psi' + Rs\bar{\psi}') \\
& + \hbar^1 \sum_{G=1}^{N_A+N_S} \left(l_{g_G}^{(1)} L_g^{(G)} + l_{A^{(G)}}^{(1)} L_A^{(G)} + l_{\omega^{(G)}}^{(1)} L_\omega^{(G)} \right) + \hbar^1 l_\psi^{(1)} L_\psi^L + \hbar^1 l_{\psi'}^{(1)} L_{\psi'}^R \\
& + \frac{\hbar^1}{(4\pi)^2} \frac{1}{3} \sum_{\substack{\mathcal{A}_1, \mathcal{A}_2 \\ \mathcal{A}_1 \neq \mathcal{A}_2}}^{N_A} \left[\left(\prod_{S=1}^{N_S} d_S \right) t_L^{\mathcal{A}_1} t_L^{\mathcal{A}_2} + \left(\prod_{S=1}^{N_S} d'_S \right) t_R^{\mathcal{A}_1} t_R^{\mathcal{A}_2} \right] \int d^d x (\partial_\mu A^{\mathcal{A}_1 \mu}) (\partial_\mu A^{\mathcal{A}_2 \mu}) \\
& + \hbar^1 \sum_{\substack{\mathcal{A}_1, \mathcal{A}_2 \\ \mathcal{A}_1 \neq \mathcal{A}_2}}^{N_A} l_{\mathcal{A}_1 \mathcal{A}_2}^{(1)} \int d^d x \frac{1}{2g_{\mathcal{A}_1} g_{\mathcal{A}_2}} F_{\mu\nu}^{\mathcal{A}_1} F^{\mathcal{A}_2 \mu\nu},
\end{aligned} \tag{92}$$

where $l_{g_G}^{(1)}$, $l_{A^{(G)}}^{(1)}$, $l_{\omega^{(G)}}^{(1)}$, $l_\psi^{(1)}$, $l_{\psi'}^{(1)}$, and each $l_{\mathcal{A}_1 \mathcal{A}_2}(1)$ for $\mathcal{A}_1 < \mathcal{A}_2 \leq N_A$ are arbitrary coefficients which will determine the renormalization conditions at the one-loop order and $L_g^{(G)}$, $L_A^{(G)}$, $L_\omega^{(G)}$, L_ψ^L , $L_{\psi'}^R$, are any of the generalizations to the d -dimensional space-time of Dimensional Regularization of the corresponding b -invariant four-dimensional operators which are written down explicitly in next subsection.

4.5. Renormalization group equation

The $O(\hbar^0)$ Lorentz invariant functionals L_i with ultraviolet dimension 4 and ghost number 0 which satisfy the following restrictions

$$b L_i = 0 \quad \frac{\delta}{\delta B_a} L_i = 0, \quad \mathcal{G} L_i = 0,$$

form a linear space. It is very convenient to have a basis for such space constituted by elements which are derived by the action of a compatible differential operator over the classical action, because, as explained in section 3.7., then, a quantum basis can be properly defined. The expansion of $\mu \partial_\mu \Gamma_{\text{ren}}$ in such a quantum basis, is just the renormalization group equation. Let us construct a basis for that linear space.

* a) *Cohomologically trivial terms*

We define for each field

$$N_\phi^a{}_b \equiv \int d^4 x \phi^a \frac{\delta}{\delta \phi^b},$$

a, b being group indices.

These are terms which are b -exact:

$$L_A{}^a{}_b \equiv b \cdot \int d^4x A_\mu^a \tilde{\rho}_b^\mu = \left[N_A{}^a{}_b - N_{\rho b}{}^a - N_{\bar{\omega} b}{}^a - N_{B b}{}^a + 2\Lambda^{ae} \frac{\partial}{\partial \Lambda^{eb}} \right] S_{\text{cl}} \equiv \mathcal{N}_A{}^a{}_b S_{\text{cl}},$$

$$L_\omega{}^a{}_b \equiv -b \cdot \int d^4x \omega^a \zeta_b = [N_\omega{}^a{}_b - N_{\zeta b}{}^a] S_{\text{cl}} \equiv \mathcal{N}_\omega{}^a{}_b S_{\text{cl}},$$

$$L_{\psi^{(\prime)}}^{\text{L(R)}i}{}_j \equiv -b \cdot \int d^4x \bar{L}_j(\bar{R}_j) P_{\text{L(R)}} \psi^{(\prime)i} = \left[N_{\psi^{(\prime)}}^{\text{L(R)}i}{}_j - N_{\bar{L}(\bar{R})j}{}^i \right] S_{\text{cl}},$$

$$L_{\bar{\psi}^{(\prime)}}^{\text{R(L)}i}{}_j \equiv -b \cdot \int d^4x \bar{\psi}_i^{(\prime)} P_{\text{R(L)}} L^j(R^j) = \left[N_{\bar{\psi}^{(\prime)}}^{\text{R(L)}i}{}_j - N_{L(R)^j}{}^i \right] S_{\text{cl}},$$

$$L_{\psi^{(\prime)}}^{\text{R(L)}i}{}_j \equiv -b \cdot \int d^4x \bar{L}_j(\bar{R}_j) P_{\text{R(L)}} \psi^{(\prime)i} = N_{\psi^{(\prime)}}^{\text{R(L)}i}{}_j S_{\text{cl}},$$

$$L_{\bar{\psi}^{(\prime)}}^{\text{L(R)}i}{}_j \equiv -b \cdot \int d^4x \bar{\psi}_i P_{\text{L(R)}} L^j(R^j) = N_{\bar{\psi}^{(\prime)}}^{\text{L(R)}i}{}_j S_{\text{cl}},$$

but not all are linearly independent:

$$\delta_i^j L_{\psi^{(\prime)}}^{\text{L(R)}i}{}_j = \delta_j^i L_{\bar{\psi}^{(\prime)}}^{\text{R(L)}i}{}_j, \quad \delta_i^j L_{\psi^{(\prime)}}^{\text{R(L)}i}{}_j = \delta_j^i L_{\bar{\psi}^{(\prime)}}^{\text{L(R)}i}{}_j.$$

In the expression of L_A in terms of field counting operators, it is used the fact that S_{cl} satisfies the gauge condition and the ghost equation.

Usually, not only the BRS symmetry is imposed to the 1PI functional, but also other symmetry of the classical action, the rigid symmetry:

$$\delta_e^{\text{rig}} \phi^a = i[T_e^{(\phi)}]_b^a \phi^b,$$

where $T_e^{(\phi)}$ is the adjoint representation for $A_\mu, \omega, \bar{\omega}, \rho_\mu, \zeta$ and B ; $T_{\text{L(R)}e}$ for $\psi^{(\prime)}, L(R)$; and $-T_{\text{L(R)}e}$ for $\bar{\psi}^{(\prime)}, \bar{L}(\bar{R})$.

Then $\delta_e^{\text{rig}} S_{\text{cl}} = 0$, which has the consequence $[\delta_e^{\text{rig}}, b] = 0$.

Also, our 1PI renormalized functional constructed from eq. (92) with Breitenlohner and Maison minimal dimensional renormalization realizes this symmetry at one-loop level.

If the 1PI functional satisfies this new requirement, the set of admissible b -invariants terms is obviously smaller. Thanks to proposition 5.9 of ref. [18], the trivial subset formed by terms which are b -exact and rigidly symmetric is fully determined by studying the restricted cohomology, *i.e.* by studying the set of terms which are b -variations of rigidly symmetric terms.

Some of them are obvious generalizations of the trivial elements of eq. (30):

$$\begin{aligned} L_A^{(\mathcal{S})} &= b \cdot \int d^4x \tilde{\rho}_{a\mathcal{S}}^\mu A_\mu^{(\mathcal{S})a\mathcal{S}}, & L_\omega^{(\mathcal{S})} &= -b \cdot \int d^4x \zeta_{a\mathcal{S}}^{(\mathcal{S})} \omega^{(\mathcal{S})a\mathcal{S}} \\ L_\psi^{\text{L}} &= -b \cdot \int d^4x \bar{L} P_{\text{L}} \psi + \bar{\psi} P_{\text{R}} L = 2 \int d^4x \frac{i}{2} \bar{\psi} \overleftrightarrow{\partial} P_{\text{L}} \psi + \bar{\psi} \gamma^\mu P_{\text{L}} T_{\text{L}}^a \psi A_\mu^a, \\ L_{\psi'}^{\text{R}} &= -b \cdot \int d^4x \bar{R} P_{\text{R}} \psi' + \bar{\psi}' P_{\text{L}} R = 2 \int d^4x \frac{i}{2} \bar{\psi}' \overleftrightarrow{\partial} P_{\text{R}} \psi' + \bar{\psi}' \gamma^\mu P_{\text{R}} T_{\text{R}}^a \psi' A_\mu^a, \\ L_\psi^{\text{R}} &= -b \cdot \int d^4x \bar{L} P_{\text{R}} \psi + \bar{\psi} P_{\text{L}} L = 2 \int d^4x \frac{i}{2} \bar{\psi} \overleftrightarrow{\partial} P_{\text{R}} \psi, \\ L_{\psi'}^{\text{L}} &= -b \cdot \int d^4x \bar{R} P_{\text{L}} \psi' + \bar{\psi}' P_{\text{R}} R = 2 \int d^4x \frac{i}{2} \bar{\psi}' \overleftrightarrow{\partial} P_{\text{L}} \psi'; \end{aligned}$$

where $\mathcal{S}$ runs from 1 to $N_{\mathcal{S}}$.

The rest are terms which are b -exact but involve possibly different Abelian components:

$$\begin{aligned}
L_A^{\mathcal{A}_1 \mathcal{A}_2} &= b \cdot \int \tilde{\rho}^{\mathcal{A}_1 \mu} A_{\mu}^{\mathcal{A}_2} = \\
&= \int d^4 x \, 2 \frac{(-1)}{4g_{\mathcal{A}_1}^2} F_{\mu\nu}^{\mathcal{A}_2} F^{\mathcal{A}_1 \mu\nu} + \bar{\psi} \bar{\gamma}^{\mu} t_L^{\mathcal{A}_1} P_L \psi A_{\mu}^{\mathcal{A}_2} + \bar{\psi}' \bar{\gamma}^{\mu} t_R^{\mathcal{A}_1} P_R \psi' A_{\mu}^{\mathcal{A}_2} - \tilde{\rho}_{\mu}^{\mathcal{A}_1} \partial^{\mu} \omega^{\mathcal{A}_2}, \\
L_{\omega}^{\mathcal{A}_1 \mathcal{A}_2} &= -b \cdot \int \zeta^{\mathcal{A}_1} \omega^{\mathcal{A}_2} = \\
&= \int d^4 x \left\{ \tilde{\rho}_{\mu}^{\mathcal{A}_1} \partial^{\mu} \omega^{\mathcal{A}_2} \bar{L} i \omega^{\mathcal{A}_2} t_L^{\mathcal{A}_1} P_L \psi + \bar{\psi} P_R t_L^{\mathcal{A}_1} i \omega^{\mathcal{A}_2} L + \bar{R} i \omega^{\mathcal{A}_2} t_R^{\mathcal{A}_1} P_R \psi' + \bar{\psi}' P_L t_R^{\mathcal{A}_1} i \omega^{\mathcal{A}_2} R \right\}.
\end{aligned}$$

Notice that similar non-diagonal terms are not allowed for non-Abelian components because of rigid invariance ($L_A^{\mathcal{A}_1 \mathcal{A}_2}$ and $L_{\omega}^{\mathcal{A}_1 \mathcal{A}_2}$ are trivially rigid invariant even if $\mathcal{A}_1 \neq \mathcal{A}_2$).

$L_A^{\mathcal{A}_1 \mathcal{A}_2}$ and $L_{\omega}^{\mathcal{A}_1 \mathcal{A}_2}$ are related with the following multiplicative field renormalizations:

$$\begin{aligned}
A_{\mu}^{\mathcal{A}_1} &= (Z_A)_{\mathcal{A}_2}^{\mathcal{A}_1} A_{\mu}^{\star \mathcal{A}_2}, & B_A &= (Z_A^{-1})_A^{\mathcal{A}_2} B_{\mathcal{A}_2}^{\star}, \\
\omega^{\mathcal{A}_1} &= (Z_{\omega})_{\mathcal{A}_2}^{\mathcal{A}_1} \omega^{\star \mathcal{A}_2}, & \bar{\omega}_{\mathcal{A}_1} &= (Z_A^{-1})_{\mathcal{A}_1}^{\mathcal{A}_2} \bar{\omega}_{\mathcal{A}_2}^{\star}, \\
\rho_{\mathcal{A}_1}^{\mu} &= (Z_A^{-1})_{\mathcal{A}_1}^{\mathcal{A}_2} \rho_{\mathcal{A}_2}^{\star \mu}, & \zeta_{\mathcal{A}_1} &= (Z_{\omega}^{-1})_{\mathcal{A}_1}^{\mathcal{A}_2} \zeta_{\mathcal{A}_2}^{\star}, \\
\Lambda^{\mathcal{A}_1 \mathcal{A}_2} &= (Z_A)_{\mathcal{A}_1'}^{\mathcal{A}_1} (Z_A)_{\mathcal{A}_2'}^{\mathcal{A}_2} \Lambda^{\star \mathcal{A}_1' \mathcal{A}_2'}
\end{aligned}$$

But, they will not be needed in renormalization group computations, because in minimal subtraction of loop diagrams this kind of terms can not appear due to the fact that they involve Abelian ghosts.

* b) *Elements of the cohomology*

Imposing the rigid invariance, we have the generalization of the unique element of the cohomology of the simple group case:

$$L_g^{(G)} = g_G \frac{\partial S_{\text{cl}}}{\partial g_G} = \frac{1}{2g_G^2} \int d^4 x \, F_{\mu\nu}^{(G) a_G} F_{a_G}^{(G) \mu\nu},$$

and the new elements (for $\mathcal{A}_1 \neq \mathcal{A}_2$):

$$L_{\mathcal{A}_1 \mathcal{A}_2} = \int d^4 x \, \frac{1}{2 g_{\mathcal{A}_1} g_{\mathcal{A}_2}} F_{\mu\nu}^{\mathcal{A}_1} F^{\mathcal{A}_2 \mu\nu}$$

with $\mathcal{A}_1, \mathcal{A}_2 \leq N_{\mathcal{A}}$.

Note that $L_{\mathcal{A}\mathcal{A}} = L_{g^{\mathcal{A}}} = g^{\mathcal{A}} \partial_{g^{\mathcal{A}}} S_{\text{cl}}$

These terms come from the renormalization of the Abelian components of the matrix C of eq. (91). At tree level, it is a diagonal matrix, but, more generally, it can be changed to a non-diagonal matrix in the Abelian part while the right hand side of eq. (91) continue being a rigidly invariant term:

$$C_{\mathcal{A}_1 \mathcal{A}_2} = (Z_C)^{\mathcal{A}'_1 \mathcal{A}'_2}_{\mathcal{A}_1 \mathcal{A}_2} C_{\mathcal{A}'_1 \mathcal{A}'_2}^*.$$

We define:

$$L_C^{\mathcal{A}_1 \mathcal{A}_2} = \frac{\partial}{\partial C_{\mathcal{A}_1 \mathcal{A}_2}} S_{\text{cl}} = -\frac{L^{\mathcal{A}_1 \mathcal{A}_2}}{2}$$

Therefore, eq. (74) becomes

$$\left[\mu \frac{\partial}{\partial \mu} + \sum_G^{N_A + N_S} \beta^G g_G \frac{\partial}{\partial g_G} - \gamma_\psi^L \mathcal{N}_\psi^L - \gamma_{\psi'}^R \mathcal{N}_{\psi'}^R - \gamma_\psi^R \mathcal{N}_\psi^R - \gamma_{\psi'}^L \mathcal{N}_{\psi'}^L \right. \\ \left. - \sum_{S=1}^{N_S} \gamma_A^S \mathcal{N}_A^{(S)} - \sum_{S=1}^{N_S} \gamma_\omega^S \mathcal{N}_\omega^{(S)} - \sum_{\substack{\mathcal{A}_1, \mathcal{A}_2 \\ \mathcal{A}_1 \neq \mathcal{A}_2}}^{N_A} \gamma_C^{\mathcal{A}_1 \mathcal{A}_2} \frac{\partial}{\partial C_{\mathcal{A}_1 \mathcal{A}_2}} \right] \Gamma_{\text{ren}} = 0,$$

which at order $\hbar$ reads

$$\mu \frac{\partial \Gamma_{\text{ren}}}{\partial \mu} = - \sum_{G=1}^{N_A + N_S} \beta^{G(1)} L_g^{(G)} + \gamma_\psi^{L(1)} L_\psi^L + \gamma_{\psi'}^{R(1)} L_{\psi'}^R \\ + \sum_{S=1}^{N_S} \gamma_A^{S(1)} L_A^{(S)} + \sum_{S=1}^{N_S} \gamma_\omega^{S(1)} L_\omega^{(S)} + \sum_{\substack{\mathcal{A}_1, \mathcal{A}_2 \\ \mathcal{A}_1 \neq \mathcal{A}_2}}^{N_A} \gamma_C^{\mathcal{A}_1 \mathcal{A}_2(1)} L_C^{\mathcal{A}_1 \mathcal{A}_2} + O(\hbar^2).$$

Eq. (88) is thus generalized.

Eq. (87) should be modified by applying the rules stated above and adding the new term:

$$+ \frac{1}{(4\pi)^2} \frac{8}{3} \sum_{\substack{\mathcal{A}_1, \mathcal{A}_2 \\ \mathcal{A}_1 \neq \mathcal{A}_2}}^{N_A} \left(\prod_{S=1}^{N_S} d_S^{(\prime)} \right) t_{L(R)}^{\mathcal{A}_1} t_{L(R)}^{\mathcal{A}_2} \int d^4 x - \frac{1}{4} F_{\mu\nu}^{\mathcal{A}_1} F^{\mathcal{A}_2 \mu\nu} S_{\text{cl} \mathcal{A} \mathcal{A}}$$

Therefore,

$$\beta^{\mathcal{A}(1)} = \frac{1}{(4\pi)^2} g_{\mathcal{A}}^2 \frac{2}{3} \left[\left(\prod_{S=1}^{N_S} d_S \right) (t_L^{\mathcal{A}})^2 + \left(\prod_{S=1}^{N_S} d'_S \right) (t_R^{\mathcal{A}})^2 \right], \\ \beta^{S(1)} = \frac{1}{(4\pi)^2} g_S^2 \left(\frac{2}{3} \left[\left(\prod_{K \neq S}^{N_S} d_K \right) T_{LS} + \left(\prod_{K \neq S}^{N_S} d'_K \right) T_{RS} \right] - \frac{11}{3} C_A^{(S)} \right), \\ \gamma_A^{S(1)} = \frac{1}{(4\pi)^2} g_S^2 \left(2 - \frac{1 - \alpha'^S}{2} \right) C_A^{(S)}, \\ \gamma_\omega^{S(1)} = \frac{1}{(4\pi)^2} g_S^2 \alpha'^S C_A^{(S)}, \\ \gamma_\psi^{L(1)} = \frac{1}{(4\pi)^2} \sum_{G=1}^{N_A + N_S} g_G^2 \alpha'^G C_L^{(G)}, \quad \gamma_{\psi'}^{R(1)} = \frac{1}{(4\pi)^2} \sum_{G=1}^{N_A + N_S} g_G^2 \alpha'^G C_R^{(G)}, \\ \gamma_C^{\mathcal{A}_1 \mathcal{A}_2(1)} = \frac{1}{(4\pi)^2} \frac{4}{3} (t_L^{\mathcal{A}_1} t_L^{\mathcal{A}_2} + t_R^{\mathcal{A}_1} t_R^{\mathcal{A}_2}).$$

where $\alpha'^G \equiv \alpha^G/g_G^2$, $1 \leq \mathcal{A}, \mathcal{A}_1, \mathcal{A}_2 \leq N_{\mathcal{A}}$, $1 \leq \mathcal{S} \leq N_{\mathcal{S}}$.

6. Conclusions

We have shown by performing explicit thorough one-loop computations that Dimensional Regularization *à la* Breitenlohner and Maison can be used blindly to carry out calculations in chiral gauge theories. Up to best of our knowledge it is the first time that such a complete one-loop study has been carried out. It shows that the renormalization method at hand has all the properties that on general grounds were expected. Notice that here the cancellation of anomalies is not used to try to hide any inconsistency. Even when there is not cancellation of anomalies, the renormalized theory is finite and unambiguous: in particular, the BRS anomaly is obtained without doing any tinkering. This is in stark contrast to the “naive” approach, where cancellation of anomalies is invoked to dangerously try to hide some inconsistencies.

We would like to stress the fact that it would not have been feasible to carry out the computations made in this paper without the help of the action principles and the BRS cohomology theory. This makes manifest the importance of these two theoretical tools in practical computations in chiral gauge theories.

Appendix A: Slavnov-Taylor Identities

This appendix is devoted to establishing eq. (19) by using the *Regularized Action Principle* as stated in eqs. (11) and (12).

Let $S_{\text{DReg}}^{(n)}$ be given by

$$S_{\text{free}}[\varphi, \Phi] + S_{\text{int}}[\varphi, \Phi; K_{\Phi}],$$

where

$$S_{\text{free}}[\varphi; \Phi] = S_{0\text{free}}[\varphi; \Phi],$$

$$S_{\text{int}}[\varphi, \Phi; K_{\Phi}; \lambda] = S_{0\text{int}}[\varphi, \Phi] + \int d^d x K_{\Phi}(x) s_d \Phi(x) + S_{\text{ct}}^{(n)}[\varphi, \Phi; K_{\Phi}].$$

The symbols φ and Φ stand, respectively, for sets of fields which transform linearly and non-linearly under arbitrary BRS transformations. The generalization to “ d -dimensions” of the BRS transformations are denoted by $s_d \phi(x)$ and $s_d \Phi(x)$. We define $s_d K_{\Phi}(x) = 0$

We next define

$$S_{\text{INT}}[\varphi, \Phi; J_{\varphi}, J_{\Phi}, K_{\Phi}] = S_{\text{int}}[\varphi, \Phi; K_{\Phi}] + \int d^d x (J_{\varphi}(x) \varphi(x) + J_{\Phi}(x) \Phi(x))$$

Following eq. (10), the generating functional Z_{DReg} is introduced next

$$Z_{\text{DReg}}[J_{\varphi}, J_{\Phi}, K_{\Phi}] = \langle \exp \left\{ \frac{i}{\hbar} S_{\text{INT}}[\varphi, \Phi; J_{\varphi}, J_{\Phi}, K_{\Phi}] \right\} \rangle_0,$$

where the symbol $\langle \cdots \rangle_0$ is defined as in the paragraph below eq. (10).

Let Δ and Δ_{ct} be defined as follows:

$$\Delta = s_d S_0, \quad \Delta_{\text{ct}} = s_d S_{\text{ct}}^{(n)},$$

where $S_0 = S_{\text{0free}}[\varphi; \Phi] + S_{\text{0int}}[\varphi, \Phi] + \int d^d x K_\Phi(x) s_d \Phi(x)$.

We now introduce $S_{\text{INT}}^{(\Upsilon)}$ for reasons that will become clear later:

$$S_{\text{INT}}^{(\Upsilon)} = S_{\text{INT}}[\varphi, \Phi; J_\varphi, J_\Phi, K_\Phi] + S_{\text{aux}},$$

where

$$S_{\text{aux}} = \int d^d x (\Upsilon_1(x) \Delta(x) + \Upsilon_2(x) \Delta_{\text{ct}}(x) + \Upsilon_3^\Phi(x) \frac{\delta S_{\text{ct}}^{(n)}}{\delta K_\Phi(x)}),$$

and $\Delta = \int d^d x \Delta(x)$ and $\Delta_{\text{ct}} = \int d^d x \Delta_{\text{ct}}(x)$. Υ_i , $i = 1, 2$ and 3 are external fields. We define $s_d \Upsilon_i(x)$ to be zero for any external field Υ_i . Then, we introduce the generating functional Z_{DReg}^Υ as follows

$$Z_{\text{DReg}}^\Upsilon[J_\varphi, J_\Phi, K_\Phi, \Upsilon_1, \Upsilon_2, \Upsilon_3^\Phi] = \langle \exp \left\{ \frac{i}{\hbar} S_{\text{INT}}^\Upsilon[\varphi, \Phi; J_\varphi, J_\Phi, K_\Phi, \Upsilon_i] \right\} \rangle_0,$$

where $\langle \cdots \rangle_0$ is defined as for Z_{DReg} .

It is obvious that the following equation holds in Dimensional Regularization:

$$Z_{\text{DReg}}[J_\varphi, J_\Phi, K_\Phi] = Z_{\text{DReg}}^\Upsilon[J_\varphi, J_\Phi, K_\Phi, \Upsilon_1 = 0, \Upsilon_2 = 0, \Upsilon_3^\Phi = 0]$$

The *Regularized Action Principle* (see eq. (11)) implies that the following equation holds in Dimensional Regularization:

$$\begin{aligned} s_d Z_{\text{DReg}}^\Upsilon &\equiv \int d^d x \left\langle \left(\Delta(x) + \Delta_{\text{ct}}(x) + (-1)^\varphi J_\varphi(x) s_\varphi(x) + (-1)^\Phi J_\Phi(x) s_\Phi(x) \right. \right. \\ &\quad \left. \left. + \text{terms proportional to } \Upsilon_i \right) \exp \left\{ \frac{i}{\hbar} S_{\text{INT}}^\Upsilon[\varphi, \Phi, J_\varphi, J_\Phi, K_\Phi, \Upsilon_i] \right\} \right\rangle_0 = 0. \end{aligned} \quad (\text{A.1})$$

Next, by using eq. (12), one easily obtains that

$$\left\langle J_\varphi(x) s_\varphi(x) \exp \left\{ \frac{i}{\hbar} S_{\text{INT}}^\Upsilon \right\} \right\rangle_0 = \frac{\hbar}{i} J_\varphi(x) \left[a_{\varphi\varphi} \frac{\delta Z_{\text{DReg}}^\Upsilon}{\delta J_\varphi(x)} + a_{\varphi\Phi} \frac{\delta Z_{\text{DReg}}^\Upsilon}{\delta J_\Phi(x)} \right] \quad (\text{A.2})$$

and

$$\begin{aligned} \left\langle J_\Phi(x) s_\Phi(x) \exp \left\{ \frac{i}{\hbar} S_{\text{INT}}^\Upsilon \right\} \right\rangle_0 &= J_\Phi(x) \left\langle \frac{\delta (S^\Upsilon - S_{\text{ct}} - S_{\text{aux}})}{\delta K_\Phi(x)} \right\rangle_0 = \\ &= \frac{\hbar}{i} J_\Phi(x) \frac{\delta Z_{\text{DReg}}^\Upsilon}{\delta K_\Phi(x)} - \frac{\hbar}{i} J_\Phi(x) \frac{\delta Z_{\text{DReg}}^\Upsilon}{\delta \Upsilon_3^\Phi(x)} + \text{terms proportional to } \Upsilon_i. \end{aligned} \quad (\text{A.3})$$

In eq. (A.2) the linear BRS transformations $s_d\varphi$ are given by $s_d\varphi = a_{\varphi\varphi}\varphi + a_{\varphi\Phi}\Phi$.

Now, by substituting eqs. (A.2) and (A.3) in eq. (A.1), one arrives at

$$\int d^d x \left[(-1)^\varphi \frac{\hbar}{i} J_\varphi(x) \left(a_{\varphi\varphi} \frac{\delta Z_{\text{DReg}}^\Upsilon}{\delta J_\varphi(x)} + a_{\varphi\Phi} \frac{\delta Z_{\text{DReg}}^\Upsilon}{\delta J_\Phi(x)} \right) + (-1)^\Phi \frac{\hbar}{i} J_\Phi(x) \left(\frac{\delta Z_{\text{DReg}}^\Upsilon}{\delta K_\Phi(x)} - \frac{\delta Z_{\text{DReg}}^\Upsilon}{\delta \Upsilon_3^\Phi(x)} \right) \right. \\ \left. + \frac{\hbar}{i} \frac{\delta Z_{\text{DReg}}^\Upsilon}{\delta \Upsilon_1(x)} + \frac{\hbar}{i} \frac{\delta Z_{\text{DReg}}^\Upsilon}{\delta \Upsilon_2(x)} + \text{terms proportional to } \Upsilon_i \right] = 0. \quad (\text{A.4})$$

Let us introduce as usual the generating functionals $Z_{c\text{DReg}}^\Upsilon[J_\varphi, J_\Phi; K_\Phi, \Upsilon_i]$ and $\Gamma_{\text{DReg}}^\Upsilon[\varphi, \Phi; K_\Phi, \Upsilon_i]$:

$$Z_{\text{DReg}}^\Upsilon[J_\varphi, J_\Phi; K_\Phi, \Upsilon_i] = \exp\left(\frac{i}{\hbar} Z_{c\text{DReg}}^\Upsilon[J_\varphi, J_\Phi; K_\Phi, \Upsilon_i]\right) \\ \Gamma_{\text{DReg}}^\Upsilon[\varphi, \Phi; K_\Phi, \Upsilon_i] = Z_{c\text{DReg}}^\Upsilon[J_\varphi, J_\Phi; K_\Phi, \Upsilon_i] - \int d^d x (J_\varphi(x)\varphi(x) + J_\Phi(x)\Phi(x)). \quad (\text{A.5})$$

where the functionals in the previous equations are to be understood as formal power series in $\hbar$ and the fields (both external and quantum). By taking advantage of eq. (A.5) one turns eq. (A.4) into an equation for the 1PI functional $\Gamma_{\text{DReg}}^\Upsilon[\varphi, \Phi; K_\Phi, \Upsilon_i]$:

$$\int d^d x \left[-\frac{\delta \Gamma_{\text{DReg}}^\Upsilon}{\delta \varphi(x)} (a_{\varphi\varphi}\varphi(x) + a_{\varphi\Phi}\Phi(x)) - \frac{\delta \Gamma_{\text{DReg}}^\Upsilon}{\delta \Phi(x)} \left(\frac{\delta \Gamma_{\text{DReg}}^\Upsilon}{\delta K_\Phi(x)} - \frac{\delta \Gamma_{\text{DReg}}^\Upsilon}{\delta \Upsilon_3^\Phi(x)} \right) \right. \\ \left. + \frac{\delta \Gamma_{\text{DReg}}^\Upsilon}{\delta \Upsilon_1(x)} + \frac{\delta \Gamma_{\text{DReg}}^\Upsilon}{\delta \Upsilon_3^\Phi(x)} + \text{term prop. to } \Upsilon_i \right] = 0 \quad (\text{A.6})$$

Finally, by setting $\Upsilon_i(x) = 0, \forall i$, in eq. (A.6) and taking into account that

$$\Gamma_{\text{DReg}}[\varphi, \Phi; K_\Phi] = \Gamma_{\text{DReg}}^\Upsilon[\varphi, \Phi; K_\Phi, \Upsilon_i = 0]$$

one obtains eq. (19) from eq. (A.6):

$$\int d^d x (s_d\varphi) \frac{\delta \Gamma_{\text{DReg}}}{\delta \varphi(x)} + \frac{\delta \Gamma_{\text{DReg}}}{\delta K_\Phi(x)} \frac{\delta \Gamma_{\text{DReg}}}{\delta \Phi(x)} = \\ \int d^d x \left[\Delta(x) \cdot \Gamma_{\text{DReg}} + \Delta_{\text{ct}}(x) \cdot \Gamma_{\text{DReg}} + \left(\frac{\delta S_{\text{ct}}}{\delta K_\Phi(x)} \cdot \Gamma_{\text{DReg}} \right) \frac{\delta \Gamma_{\text{DReg}}}{\delta \Phi(x)} \right]. \quad (\text{A.7})$$

Notice that

$$\left[\frac{\delta \Gamma_{\text{DReg}}^\Upsilon}{\delta K_\Phi(x)} - \frac{\delta \Gamma_{\text{DReg}}^\Upsilon}{\delta \Upsilon_3^\Phi(x)} \right]_{\Upsilon_i=0} = \frac{\delta \Gamma_{\text{DReg}}}{\delta K_\Phi(x)} - \frac{\delta S_{\text{ct}}}{\delta K_\Phi(x)} \cdot \Gamma_{\text{DReg}}, \\ \left[\frac{\delta \Gamma_{\text{DReg}}^\Upsilon}{\delta \Upsilon_1(x)} \right]_{\Upsilon_i=0} = \Delta(x) \cdot \Gamma_{\text{DReg}} \quad \text{and} \\ \left[\frac{\delta \Gamma_{\text{DReg}}^\Upsilon}{\delta \Upsilon_2(x)} \right]_{\Upsilon_i=0} = \Delta_{\text{ct}}(x) \cdot \Gamma_{\text{DReg}},$$

for eq. (12) holds.

Appendix B. Slavnov-Taylor Identities for ε -dependent BRS

This appendix is devoted to the study of the contributions to the right hand side of eq. (21) coming from BRS variations which depend explicitly on $\varepsilon = 4 - d$. We shall see that, unlike for the BRS transformations in eq. (31), the variation under these new BRS transformations of the singular counterterms yield a finite contribution to the right hand side of eq. (21).

To introduce an explicit ε dependence on the Dimensional Regularization BRS transformations, we shall replace the BRS transformation of A_μ given in eq. (31) with $s'_d A_\mu = \nabla_\mu \omega + C(d-4)\nabla_\mu \omega$. We shall assume, however, that s'_d acts on the rest of the fields as s_d does. Hence, s'_d defines BRS transformations with explicit ε dependence.

Although we have changed the Dimensional Regularization BRS transformations, we shall not take a new Dimensional Regularization classical action, which still will be S_0 in eq. (32). Thus, we will not modify the minimally renormalized 1PI functional Γ_{minren} , so that we will have the effect of the new Dimensional Regularization BRS transformations isolated.

The s'_d -variation of S_0 is given by:

$$s'_d S_0 = \hat{\Delta} + \Delta_{\text{f}}^\varepsilon + \Delta_{\text{gf}}^\varepsilon + \Delta_\rho^\varepsilon.$$

Here, $\hat{\Delta} = s_d S_0$ is the breaking given in eq. (34) and

$$\begin{aligned} \Delta_{\text{f}}^\varepsilon &= -\varepsilon C \int d^d x (\bar{\psi} \bar{\gamma}^\mu P_L T_L^a \psi + \bar{\psi}' \bar{\gamma}^\mu T_R^a \psi') (\nabla_\mu \omega)^a, \\ \Delta_{\text{gf}}^\varepsilon &= -\varepsilon C \int d^d x B^a \partial^\mu (\nabla_\mu \omega)^a - c^{abc} \partial^\mu \bar{\omega}^a (\nabla_\mu \omega)^b \omega^c, \\ \Delta_\rho^\varepsilon &= +\varepsilon C \int d^d x c^{abc} \rho^{a\mu} (\nabla_\mu \omega)^b \omega^c. \end{aligned}$$

are the new breakings (which are also evanescent operators). Notice that $s_d'^2 A_\mu^a \neq 0$, whereas $s_d^2 A_\mu^a = 0$.

Now, let Γ_0 denote dimensionally regularized 1PI functional obtained from S_0 , then, regularized Slavnov-Taylor equation for the s'_d -transformations reads:

$$\mathcal{S}'_d(\Gamma_0) = \left[\hat{\Delta} + \Delta_{\text{f}}^\varepsilon + \Delta_{\text{gf}}^\varepsilon + \Delta_\rho^\varepsilon \right] \cdot \Gamma_0, \quad (\text{B.1})$$

where $\mathcal{S}'_d$ is the Slavnov-Taylor operator for s'_d . Notice that

$$\begin{aligned} \mathcal{S}'_d(\Gamma_0) \equiv \int d^d x \left\{ (1 - C\varepsilon) \text{Tr} \frac{\delta \Gamma_0}{\delta \rho^\mu} \frac{\delta \Gamma_0}{\delta A_\mu} + \text{Tr} \frac{\delta \Gamma_0}{\delta \zeta} \frac{\delta \Gamma_0}{\delta \omega} + \text{Tr} B \frac{\delta \Gamma_0}{\delta \bar{\omega}} \right. \\ \left. + \frac{\delta \Gamma_0}{\delta \bar{L}} \frac{\delta \Gamma_0}{\delta \psi} + \frac{\delta \Gamma_0}{\delta \bar{R}} \frac{\delta \Gamma_0}{\delta \psi'} + \frac{\delta \Gamma_0}{\delta L} \frac{\delta \Gamma_0}{\delta \bar{\psi}} + \frac{\delta \Gamma_0}{\delta R} \frac{\delta \Gamma_0}{\delta \bar{\psi}'} \right\}. \end{aligned}$$

The next issue to address is the derivation of the (anomalous) Slavnov-Taylor identity satisfied by the minimal dimensional renormalization 1PI functional, Γ_{minren} . If we, naively,

just replace in eq. (B.1) every regularized object with their minimally renormalized counterpart, we will obtain the following equation

$$\mathcal{S}(\Gamma_{\text{minren}}) = \text{N} \left[\hat{\Delta} + \Delta_{\text{f}}^{\varepsilon} + \Delta_{\text{gf}}^{\varepsilon} + \Delta_{\rho}^{\varepsilon} \right] \cdot \Gamma_{\text{minren}}. \quad (\text{B.2})$$

This equation does not hold, though, since the coefficients of the regularized Slavnov-Taylor equation, eq. (B.1), do have an explicit ε dependence. Indeed, at the one-loop level the following identity holds

$$\mathcal{S}(\Gamma_{\text{minren}}) = \left[\text{N} \left[\hat{\Delta} \right] \cdot \Gamma_{\text{minren}} \right]^{(1)} + O(\hbar^2). \quad (\text{B.3})$$

This has been shown in subsection 3.4. Now, let us substitute the previous equation in eq. (B.2). We thus come to the conclusion that $\text{N} \left[\Delta_{\text{f}}^{\varepsilon} + \Delta_{\text{gf}}^{\varepsilon} + \Delta_{\rho}^{\varepsilon} \right] \cdot \Gamma_{\text{minren}}$ should vanish at the one-loop level. Which is nonsense, for a diagrammatic computation shows that

$$\begin{aligned} & \left[\text{N} \left[\Delta_{\text{f}}^{\varepsilon} + \Delta_{\text{gf}}^{\varepsilon} + \Delta_{\rho}^{\varepsilon} \right] \cdot \Gamma_{\text{minren}} \right]^{(1)} = \\ & \frac{-\hbar}{(4\pi)^2} C g^2 \left\{ (\alpha' C_{\text{A}} + 2 \alpha' C_{\text{L}}) \int d^4 x \frac{\delta S_{\text{cl}} \bar{\psi} \psi A}{\delta A_{\mu}^a} \partial_{\mu} \omega^a \right. \\ & + \left[\left(3 - \frac{3}{2}(1 - \alpha') \right) C_{\text{A}} + 2 \alpha' C_{\text{L}} \right] \int d^4 x \frac{\delta S_{\text{cl}} \bar{\psi} \psi A}{\delta A_{\mu}^a} c^{abc} A_{\mu}^b \omega^c \\ & + (\alpha' C_{\text{A}} + 2 \alpha' C_{\text{R}}) \int d^4 x \frac{\delta S_{\text{cl}} \bar{\psi}' \psi' A}{\delta A_{\mu}^a} \partial_{\mu} \omega^a \\ & + \left[\left(3 - \frac{3}{2}(1 - \alpha') \right) C_{\text{A}} + 2 \alpha' C_{\text{R}} \right] \int d^4 x \frac{\delta S_{\text{cl}} \bar{\psi}' \psi' A}{\delta A_{\mu}^a} c^{abc} A_{\mu}^b \omega^c \\ & - \frac{3}{2} (1 - \alpha') C_{\text{A}} \int d^4 x \left(\frac{\delta S_{\text{cl}} \bar{\omega} \omega A}{\delta A_{\mu}^a} + \frac{\delta S_{\text{cl}} \rho \omega A}{\delta A_{\mu}^a} \right) \partial_{\mu} \omega^a \\ & + 2 \alpha' C_{\text{A}} \int d^4 x \left(\frac{\delta S_{\text{cl}} \bar{\omega} \omega A}{\delta A_{\mu}^a} + \frac{\delta S_{\text{cl}} \rho \omega A}{\delta A_{\mu}^a} \right) c^{abc} A_{\mu}^b \omega^c \\ & + \left(1 + \frac{1}{2}(1 - \alpha') \right) C_{\text{A}} \int d^4 x (\partial_{\mu} B^a) (\partial^{\mu} \omega^a) \\ & \left. - \alpha' C_{\text{A}} \int d^4 x (\partial_{\mu} B^a) c^{abc} A_{\mu}^b \omega^c \right\}. \quad (\text{B.4}) \end{aligned}$$

Hence, a contribution to the right hand side of eq. (B.2) is missing. This missing contribution should apparently be furnished by the second and third terms on the right hand side of eq. (21), which is a consequence of eq. (A.7). But, it is essential to remember the assumption made in Appendix A in order to properly deduce eq. (A.7): the dependence in K_{Φ} in the starting dimensional regularized action should be $\int K_{\Phi} s'_d \Phi$, if s'_d is the variation under study.

Therefore, due to the fact that the dependence in ρ_μ of the chosen action is just $\int \rho_\mu s_d A^\mu$, an extra factor of $1 - C\varepsilon$ is needed in eq. (A.3), when studying s'_d variations:

$$\begin{aligned} \left\langle J_{A_a}^\mu(x) s'_d A_\mu^a(x) \exp\left\{\frac{i}{\hbar} S_{\text{INT}}^\Upsilon\right\} \right\rangle_0 &= (1 - C\varepsilon) \left\langle J_{A_a}^\mu(x) s'_d A_\mu^a(x) \exp\left\{\frac{i}{\hbar} S_{\text{INT}}^\Upsilon\right\} \right\rangle_0 = \\ &= (1 - C\varepsilon) \frac{\hbar}{i} J_{A_a}^\mu(x) \left(\frac{\delta Z_{\text{DReg}}^\Upsilon}{\delta \rho_\mu^a(x)} - \frac{\delta Z_{\text{DReg}}^\Upsilon}{\delta \Upsilon_3^{A^\mu}(x)} \right) + \text{terms proportional to } \Upsilon_i. \end{aligned}$$

Then, the equation (A.7) becomes

$$\begin{aligned} \mathcal{S}'_d(\Gamma_{\text{DReg}}) &= s'_d(S_0 + S_{\text{ct}}) \cdot \Gamma_{\text{DReg}} + \\ &+ \int d^d x \left[\sum_{\Phi \neq A} \left(\frac{\delta S_{\text{ct}}}{\delta K_\Phi(x)} \cdot \Gamma_{\text{DReg}} \right) \frac{\delta \Gamma_{\text{DReg}}}{\delta \Phi(x)} + (1 - C\varepsilon) \left(\frac{\delta S_{\text{ct}}}{\delta \rho_\mu^a(x)} \cdot \Gamma_{\text{DReg}} \right) \frac{\delta \Gamma_{\text{DReg}}}{\delta A_\mu^a(x)} \right], \end{aligned}$$

and the correct modification of eq. (B.2) is:

$$\begin{aligned} \mathcal{S}(\Gamma_{\text{minren}}) &= \left[N \left[\hat{\Delta} + \Delta_{\text{f}}^\varepsilon + \Delta_{\text{gf}}^\varepsilon + \Delta_\rho^\varepsilon \right] \cdot \Gamma_{\text{minren}} \right]^{(1)} + \\ &+ C \int d^4 x \left[(\nabla_\mu \omega)^a \frac{\delta \mathcal{F}}{\delta A_\mu^a} + \frac{\delta S_{\text{cl}}}{\delta A_\mu^a} \frac{\delta \mathcal{F}}{\delta \rho_a^\mu} \right] + O(\hbar^2). \end{aligned} \quad (\text{B.5})$$

Here, $\mathcal{F} = \text{LIM}_{d \rightarrow 4} \left\{ \varepsilon \Gamma_{0 \text{ sing}}^{(1)} \right\}$, with $\Gamma_{0 \text{ sing}}^{(1)}$ as given by eqs. (36) and (37). Notice that the second term on the right hand side of (B.5) cancels the contribution displayed in eq. (B.4). Eq. (B.3) is thus recovered.

Of course, if we want to use eqs. (19), (21) and (22)—with s'_d instead of s_d —, we have to choose the following new Dimensional Regularization classical action:

$$S'_0 = \left(S_0 - \int d^d x \rho_\mu^a s_d A^{a\mu} \right) + \int d^d x \rho_\mu^a s'_d A^{a\mu},$$

which cause the breaking

$$s'_d S'_0 = \hat{\Delta} + \Delta_{\text{f}}^\varepsilon + \Delta_{\text{gf}}^\varepsilon + (1 - C\varepsilon) \Delta_\rho^\varepsilon,$$

and a new perturbative renormalized 1PI functional Γ'_{minren} .

Therefore, in this case, from eq. (19), (21) and (22), we conclude that the (anomalous) one-loop renormalized Slavnov-Taylor identity reads is

$$\begin{aligned} \mathcal{S}(\Gamma'_{\text{minren}}) &= \left[N [s'_d S'_0] \cdot \Gamma'_{\text{minren}} \right]^{(1)} + C \int d^4 x \left[(\nabla_\mu \omega)^a \frac{\delta \mathcal{F}}{\delta A_\mu^a} + \frac{\delta S_{\text{cl}}}{\delta A_\mu^a} \frac{\delta \mathcal{F}}{\delta \rho_a^\mu} \right] + \\ &+ \int d^4 x \frac{\delta (S_{\text{cl} \rho \omega} + S_{\text{cl}})_{\rho \omega A}}{\delta \omega^a} \frac{\delta \mathcal{F}}{\delta \zeta^a} + O(\hbar^2). \end{aligned}$$

Appendix C. Comparison between “naive” and Breitenlohner and Maison minimal dimensional renormalization schemes

In this appendix we shall make a thorough comparison of the minimally renormalized one-loop 1PI functionals obtained within the “naive” dimensional renormalization prescription, on the one hand, and the Breitenlohner and Maison Dimensional Regularization scheme, on the other. The quantum field theory under scrutiny will be the theory already studied in section 3. Obviously, at the one-loop level, differences only arise in diagrams involving fermions. We will denote these contributions with the subscript $_F$. All along this appendix, the subscripts BM and “naive” will indicate that the corresponding quantity has been evaluated by using the Breitenlohner and Maison scheme and the “naive” prescription, respectively.

Let us begin with the self-energy of the gauge field. Here, only the fermionic loops may give rise to differences between the two calculational techniques. There are terms with one γ_5 and four γ s in the trace, but their contribution explicitly vanish, the reason being the contraction of two γ s of the antisymmetric trace with the same momentum q . The “naive” prescription does not yield ambiguous the results in this instance. These results read

$$\tilde{\Gamma}_{F AA \text{ naive}}^{(1) \mu\nu, ab}(p) = \frac{1}{4\pi^2}(T_L + T_R) \left[\frac{1}{3\varepsilon} - \frac{\gamma}{6} + \frac{5}{18} - \frac{1}{6} \ln \frac{-p^2 + i\epsilon}{4\pi\mu^2} + O(\varepsilon) \right] \delta^{ab} (p^\mu p^\nu - p^2 g^{\mu\nu}),$$

for the regularized theory and

$$\tilde{\Gamma}_{F AA \text{ naive}}^{R(1) \mu\nu, ab}(p) = \frac{1}{4\pi^2}(T_L + T_R) \left[-\frac{\gamma}{6} + \frac{5}{18} - \frac{1}{6} \ln \frac{-p^2 + i\epsilon}{4\pi\mu^2} \right] \delta^{ab} (p^\mu p^\nu - p^2 g^{\mu\nu}),$$

for the renormalized Green function. It should be understood a sum over the different fermionic multiplets: $\sum_{k=1}^{n_L} T_L^k + \sum_{r=1}^{n_R} T_R^r$. Notice that both results are transverse, so that the corresponding Slavnov-Taylor identity holds.

The calculation within the Breitenlohner and Maison framework is also straightforward and it yields the following regularized expression

$$\begin{aligned} \tilde{\Gamma}_{F AA \text{ BM}}^{(1) \mu\nu, ab}(p) = & \delta^{ab} \left\{ \frac{1}{4\pi^2}(T_L + T_R) \left[\frac{1}{3\varepsilon} - \frac{\gamma}{6} + \frac{5}{18} - \frac{1}{6} \ln \frac{-p^2 + i\epsilon}{4\pi\mu^2} \right] (\bar{p}^\mu \bar{p}^\nu - \bar{p}^2 \bar{g}^{\mu\nu}) \right. \\ & - \frac{1}{4\pi^2}(T_L + T_R) \frac{1}{12} \bar{g}^{\mu\nu} \bar{p}^2 \\ & \left. - \frac{1}{4\pi^2} \frac{(T_L + T_R)}{2} \left[\frac{1}{3\varepsilon} - \frac{\gamma}{6} + \frac{5}{18} - \frac{1}{6} \ln \frac{-p^2 + i\epsilon}{4\pi\mu^2} + \frac{1}{6} \right] \bar{g}^{\mu\nu} \hat{p}^2 \right\} + O(\varepsilon). \end{aligned} \quad (\text{C.1})$$

Notice that there is a singular contribution at $\varepsilon = 0$ whose residue is a an “evanescent” polynomial of the momentum. Minimal subtraction has to be applied also to this contribution; in the language of ultraviolet divergent counterterms: counterterms with hatted objects are also needed! This is important for the consistency of calculations in the next orders [32].

The renormalized 1PI fermionic contribution to the gauge field two-point function defined by minimal subtraction *à la* Breitenlohner and Maison reads

$$\tilde{\Gamma}_{F AA BM}^{R(1) \mu\nu,ab}(p) = \tilde{\Gamma}_{F AA \text{naive}}^{R(1) \mu\nu,ab}(p) - \delta^{ab} \frac{1}{4\pi^2} \frac{(T_L + T_R)}{12} p^2 g^{\mu\nu},$$

which differs from the “naive” result in a non-transversal quantity, so that it is clear that this result do not satisfy the Slavnov-Taylor identity.

But, we learned from eq. (40) that the Slavnov-Taylor identities for the regularized theory *à la* Breitenlohner and Maison have a breaking, so instead of transversality the following equation should hold

$$-ip_\mu \tilde{\Gamma}_{AA BM}^{(1) \mu\nu,ab}(p) = \tilde{\Gamma}_{A\omega; \hat{\Delta} BM}^{(1) \nu,ab}(p; 0) \quad (C.2)$$

(because the insertion of the breaking is integrated, it is an insertion at zero momentum).

The right hand side of eq. (C.2) is just the sum of the 1PI diagrams in fig. 5 with $\hat{\Delta}$ instead of Δ and p instead of p_1 . Their value is:

$$\begin{aligned} \tilde{\Gamma}_{A\omega; \hat{\Delta} BM}^{(1) \mu,ab}(p; q) = \delta^{ab} \left\{ \frac{i}{4\pi^2} \frac{(T_L + T_R)}{2} \left[\frac{1}{3\varepsilon} - \frac{\gamma}{6} + \frac{5}{18} - \frac{1}{6} \ln \frac{-p^2 + i\epsilon}{4\pi\mu^2} \right] \bar{p}^\mu \hat{p}^2 \right. \\ \left. + \frac{i}{4\pi^2} \frac{(T_L + T_R)}{12} \bar{p}^\mu p^2 \right\} + O(\varepsilon). \end{aligned} \quad (C.3)$$

As it should be. Indeed, eqs. (C.1) and (C.3) satisfy ($O(\varepsilon)$) eq. (C.2). The contribution from the sum of diagrams with bosonic and ghost loops is obviously transversal and thus it is of no bearing here.

From eq. (42) we get the renormalized counterpart to eq. (C.2):

$$-ip_\mu \tilde{\Gamma}_{AA BM}^{R(1) \mu\nu,ab}(p) = \tilde{\Gamma}_{A\omega; N[\hat{\Delta}] BM}^{R(1) \nu,ab}(p; 0),$$

which is also satisfied because minimal subtraction applied to eq. (C.3) yields

$$\tilde{\Gamma}_{A\omega; N[\hat{\Delta}] BM}^{R(1) \mu,ab}(p; q) = \delta^{ab} \frac{i}{4\pi^2} \frac{(T_L + T_R)}{12} p^\mu p^2.$$

We have checked thus eq. (42) by explicit computation of both its sides.

The following results will be needed as well

$$\begin{aligned} \tilde{\Gamma}_{\bar{\psi}\psi BM}^{R(1) \alpha i, \beta j}(p) - \tilde{\Gamma}_{\bar{\psi}\psi \text{naive}}^{R(1) \alpha i, \beta j}(p) = \\ = -\frac{1}{(4\pi)^2} g^2 \left[1 + (\alpha' - 1) \frac{2}{3} \right] C_L [\not{p} P_L]_{\alpha\beta} \delta_{ij}, \\ \tilde{\Gamma}_{A\bar{\psi}\psi BM}^{R(1) \nu, \alpha i, \beta j}(q, p) - \tilde{\Gamma}_{A\bar{\psi}\psi \text{naive}}^{R(1) \nu, \alpha i, \beta j}(q, p) = \\ = -\frac{1}{(4\pi)^2} g^2 (C_L - \frac{C_A}{2}) \left[2 + (\alpha' - 1) \frac{5}{6} \right] [\gamma^\mu P_L]_{\alpha\beta} [T_L^a]_{ij} + \\ - \frac{1}{(4\pi)^2} g^2 C_A \left[1 + (\alpha' - 1) \frac{5}{12} \right] [\gamma^\mu P_L]_{\alpha\beta} [T_L^a]_{ij} \\ = -\frac{1}{(4\pi)^2} g^2 C_L \left[2 + (\alpha' - 1) \frac{5}{6} \right] [\gamma^\mu P_L]_{\alpha\beta} [T_L^a]_{ij}. \end{aligned}$$

Now, we anticommute the γ_5 s of the projectors in the “naive” expression towards the right to get

$$N_{abc}^{\mu\nu\rho \text{ naive}} = \frac{T_R^{abc} + T_L^{abc}}{2} \text{Tr} [(\not{k} - \not{k}_1)\gamma^\mu \not{k}\gamma^\nu (\not{k} + \not{k}_2)\gamma^\rho] + \frac{T_R^{abc} - T_L^{abc}}{2} \text{Tr} [(\not{k} - \not{k}_1)\gamma^\mu \not{k}\gamma^\nu (\not{k} + \not{k}_2)\gamma^\rho \gamma_5]. \quad (\text{C.4})$$

Notice that when the integration over the k s is done a trace over six gammas, two of them contracted, and a γ_5 occurs. This is precisely the source of the ambiguity displayed by eq. (7)! Indeed, if we had decided to leave the matrix γ_5 in another place, the minimally renormalized results should differ in a local quantity: there is a divergent loop present, and the ambiguity of eq. (7) is proportional to $\varepsilon = 4 - d$. This is the reason why we have labeled the numerator with the subscript 1.

Therefore, we can write

$$iL_{abc}^{\mu\nu\rho}(k_1, k_2, k_3) + iR_{abc}^{\mu\nu\rho}(k_1, k_2, k_3) = \frac{T_R^{abc} + T_L^{abc}}{2} T_{VVV}^{\mu\nu\rho}(k_1, k_2, k_3) + \frac{T_R^{abc} - T_L^{abc}}{2} T_{VVA}^{\mu\nu\rho}(k_1, k_2, k_3), \quad (\text{C.5})$$

where T_{VVV} stands for the Abelian Green function (*i.e.* without gauge group matrices) and with three vector-like vertices and $T_{VVA}^{\mu\nu\rho}(k_1, k_2, k_3)$ denotes the Abelian diagram with an axial vertex $\gamma^\rho \gamma_5$ in the leg carrying momentum k_3 .

The Breitenlohner and Maison counterpart to eq. (C.4) is (we have used eq. (26)) the following

$$N_{abc}^{\mu\nu\rho \text{ BM}} = \frac{T_R^{abc} + T_L^{abc}}{2} \text{Tr} [(\bar{\not{k}} - \bar{\not{k}}_1)\bar{\gamma}^\mu \bar{\not{k}}\bar{\gamma}^\nu (\bar{\not{k}} + \bar{\not{k}}_2)\bar{\gamma}^\rho] + \frac{T_R^{abc} - T_L^{abc}}{2} \text{Tr} [(\bar{\not{k}} - \bar{\not{k}}_1)\bar{\gamma}^\mu \bar{\not{k}}\bar{\gamma}^\nu (\bar{\not{k}} + \bar{\not{k}}_2)\bar{\gamma}^\rho \gamma_5].$$

It is easy to calculate the difference between the renormalized result obtained with the help of eq. (C.4) (*do not move around the γ_5 “matrix”*) and the renormalized result obtained from eq. (C.5): the only contribution to this difference comes from the renormalization of the dimensionally regularized integral whose numerator is given by

$$\begin{aligned} & \frac{T_R^{abc} + T_L^{abc}}{2} \{ \text{Tr} [\hat{\not{k}}\bar{\gamma}^\mu \hat{\not{k}}\bar{\gamma}^\nu (\bar{\not{k}} + \bar{\not{k}}_2)\bar{\gamma}^\rho] + \text{Tr} [\hat{\not{k}}\bar{\gamma}^\mu \bar{\not{k}}\bar{\gamma}^\nu \hat{\not{k}}\bar{\gamma}^\rho] + \text{Tr} [(\bar{\not{k}} - \bar{\not{k}}_1)\bar{\gamma}^\mu \hat{\not{k}}\bar{\gamma}^\nu \hat{\not{k}}\bar{\gamma}^\rho] \} + \\ & \frac{T_R^{abc} - T_L^{abc}}{2} \{ \text{Tr} [\hat{\not{k}}\bar{\gamma}^\mu \hat{\not{k}}\bar{\gamma}^\nu (\bar{\not{k}} + \bar{\not{k}}_2)\bar{\gamma}^\rho \gamma_5] + \text{Tr} [\hat{\not{k}}\bar{\gamma}^\mu \bar{\not{k}}\bar{\gamma}^\nu \hat{\not{k}}\bar{\gamma}^\rho \gamma_5] + \\ & \quad \text{Tr} [(\bar{\not{k}} - \bar{\not{k}}_1)\bar{\gamma}^\mu \hat{\not{k}}\bar{\gamma}^\nu \hat{\not{k}}\bar{\gamma}^\rho \gamma_5] \}. \end{aligned}$$

The value of that difference is thus the following

$$\begin{aligned} & iL_{abc \text{ BM}}^{\mu\nu\rho R}(k_1, k_2, k_3) + iR_{abc \text{ BM}}^{\mu\nu\rho R}(k_1, k_2, k_3) \\ & - iL_{abc \text{ naive}_1}^{\mu\nu\rho R}(k_1, k_2, k_3) - iR_{abc \text{ naive}_1}^{\mu\nu\rho R}(k_1, k_2, k_3) = \\ & = \frac{i}{(4\pi)^2} \frac{T_R^{abc} + T_L^{abc}}{2} 4 \left[g^{\mu\nu} \frac{(k_1 - k_2)^\rho}{3} + g^{\nu\rho} \frac{(k_1 + 2k_2)^\mu}{3} - g^{\mu\rho} \frac{(2k_1 + k_2)^\nu}{3} \right] \\ & + \frac{i}{(4\pi)^2} \frac{T_R^{abc} - T_L^{abc}}{2} \text{Tr} \left[\frac{(\not{k}_1 - \not{k}_2)}{3} \gamma^\mu \gamma^\nu \gamma^\rho \gamma_5 \right]. \quad (\text{C.6}) \end{aligned}$$

Notice that the second term of the previous equation is not cyclic, *i.e.* if we make the change $(k_1, \mu, a) \rightarrow (k_2, \nu, b)$, $(k_2, \nu, b) \rightarrow (k_3, \rho, c)$, $(k_3, \rho, c) \rightarrow (k_1, \mu, a)$ the term changes. This seems to be incompatible with having a trace which is cyclic (or with the fact that we can start to read a closed fermion loop wherever we want). Of course, if a prescription for Feynman rules is consistent, the renormalized result should be also cyclic. This is the case for the Breitenlohner and Maison result, as can be checked explicitly. But in the “naive” prescription case, if we insist on an anticommuting γ_5 , the cyclicity of the trace is to be abandoned, lest the trace of four gammas and one γ_5 vanishes. The lack of cyclicity of eq. (C.6) is due, of course, to the inconsistency of the “naive” prescription with the cyclicity of the trace.

One could insist on keeping an anticommuting γ_5 and cyclicity of the trace. But, then, as we know from eq. (6), the trace of four gammas and a γ_5 should unavoidably vanish. This would not do since it would set to zero any dimensionally regularized Feynman integral that is finite by power-counting at $d = 4$ and whose Dirac algebra yields a contribution proportional to the trace of four gammas and one γ_5 .

The cyclicity of the diagram is also needed for obtaining a Bose symmetric 1PI function after summing the crossed diagram. Taking into account this crossed diagram the difference between the complete 1PI functions in both schemes is:

$$\begin{aligned} & \tilde{\Gamma}_{abc}^{(1)\mu\nu\rho R}{}_{\text{BM}}(k_1, k_2, k_3) - \tilde{\Gamma}_{abc}^{(1)\mu\nu\rho R}{}_{\text{naive}_1}(k_1, k_2, k_3) = \\ &= \frac{1}{(4\pi)^2} \frac{2}{3} i c^{abc} (T_L + T_R) [g^{\mu\nu}(k_1 - k_2)^\rho + g^{\nu\rho}(k_2 - k_3)^\mu - g^{\mu\rho}(k_3 - k_1)^\nu] \\ &+ \frac{1}{(4\pi)^2} \frac{1}{6} [\text{Tr}\{T_R^a, T_R^b\}T_R^c - \text{Tr}\{T_L^a, T_L^b\}T_L^c] \text{Tr}[(\not{k}_2 - \not{k}_1)\gamma^\mu\gamma^\nu\gamma^\rho\gamma_5]. \quad (\text{C.7}) \end{aligned}$$

The second term on the right hand side of the previous equation is not Bose symmetric unless it is zero.

Notice that the problem of the inconsistency of the “naive” prescription can be swept under the carpet by choosing the matter content so that $\text{Tr}\{T_R^a, T_R^b\}T_R^c - \text{Tr}\{T_L^a, T_L^b\}T_L^c$ ($\equiv d_R^{abc} - d_L^{abc} \equiv d_{R-L}$) vanish, *i.e.* if there is cancellation of anomalies. Moreover, we have compared the unambiguous Breitenlohner and Maison computation with the “naive” result obtained by putting the γ_5 after the γ^ρ (see eq. (C.4)). If we had put it after, say, the γ^μ , the new difference with the result *à la* Breitenlohner and Maison has exactly the same first term as the difference in eq. (C.7) but its second term is the second contribution in the right hand side of eq. (C.7), upon having changed cyclically the indices and momenta of the latter. That is, *the one-loop difference between two possible “naive” calculations is a term involving a product of an epsilon tensor and the coefficient $d_{R-L}^{bca} = d_{R-L}^{abc} = d_{R-L}^{abc} \dagger$* . This justifies the statement that *if there is cancellation of anomalies there is no ambiguity in the one-loop “naive” computation of the three boson vertex*. In ref. [29], it was correctly claimed that the one-loop essential anomaly could be also obtained with the “naive” prescription for the Abelian case from a VVA triangle diagram by simply not moving around the *unique* γ_5 which appear in the traces, computing explicitly the

$\dagger \{ \dots \}$ enclosing indices means symmetrization and $[\dots]$ antisymmetrization

dimensional integrals and contracting, at the end of the day, the final expression with the momentum in the axial vertex. In fact, with our notation

$$k_3^\rho T_{\text{VVA}}^{\mu\nu\rho}(k_1, k_2, k_3) = -\frac{i}{(4\pi)^2} \text{Tr}[\not{k}_1 \not{k}_2 \gamma^\mu \gamma^\nu \gamma_5], \quad (\text{C.8})$$

which, with the crossed diagram, gives correctly the Abelian chiral anomaly. Of course, also is checked explicitly that

$$k_1^\mu T_{\text{VVA}}^{\mu\nu\rho}(k_1, k_2, k_3) = k_2^\nu T_{\text{VVA}}^{\mu\nu\rho}(k_1, k_2, k_3) = 0.$$

It is widely known that a zero (so, incorrect) result can be obtained by performing shifts in the integration variables and using the cyclicity of the trace and the *anticommutativity* of γ_5 . Therefore, the correctness of the results above seems to be due to the fact that in the explicit calculations no use of the anticommutativity of γ_5 have been made. Unfortunately, for this restricted -the γ_5 is not moved around- “naive” prescription action principles cannot be usefully applied (the action is not invariant if the γ_5 is not anticommutated) and, more importantly, *in local gauge chiral theories*, such as the model studied here, *triangle diagrams VAA and AAA also appear*. If one decides not to anticommute the γ_5 s before the subtraction is made, one cannot tell before hand whether the result will satisfy the Slavnov-Taylor identities with the correct essential breaking, unless explicit check is made order by order. And if one naively anticommutes the γ_5 s, then one will obtain ambiguous results (which turns out to be safe in one-loop calculations if there is cancellation of anomalies, but it is not clear at all what happens at higher orders)

Of course, the Breitenlohner and Maison scheme is a consistent method and has none of these problems. From eq. (42) we get the explicit identity which involves $\tilde{\Gamma}_{AAA}^{\text{R}(1)}$:

$$\begin{aligned} & -i(k+p)^\mu \tilde{\Gamma}_{AAA}^{\text{R}(1) bca}{}_{\nu\rho\mu}(k, p) \\ & -i c^{bce} [(2k+p)_\rho g_{\mu\nu} - (k+2p)_\nu g_{\rho\mu} + (p-k)_\mu g_{\nu\rho}] \tilde{\Gamma}_{w;N[sA]}^{\text{R}(1) ae}{}_\mu(; k+p) \\ & + c^{eba} \tilde{\Gamma}_{AA}^{\text{R}(1) ce}{}_{\rho\nu}(p) + c^{eca} \tilde{\Gamma}_{AA}^{\text{R}(1) be}{}_{\nu\rho}(k) - 1/g^2 p^2 P_T^{\rho\mu}(p) \tilde{\Gamma}_{Aw;N[sA]}^{\text{R}(1) bac}{}_{\nu\mu}(k; p) \\ & - 1/g^2 k^2 P_T^{\rho\mu}(k) \tilde{\Gamma}_{Aw;N[sA]}^{\text{R}(1) cab}{}_{\rho\mu}(p; k) = \tilde{\Gamma}_{AAw;N[\hat{\Delta}]}^{\text{R}(1) bca}{}_{\nu\rho}(k, p; 0), \end{aligned}$$

where $P_T^{\mu\nu}(k)$ is the transversal Lorentz projector $g^{\mu\nu} - \frac{k^\mu k^\nu}{k^2}$.

The diagrams without fermions satisfy the Slavnov-Taylor identity without breaking; therefore,

$$\begin{aligned} & -i(k+p)_\mu \tilde{\Gamma}_{FAAA}^{\text{R}(1) bca}{}_{\nu\rho\mu}(k, p) + c^{eba} \tilde{\Gamma}_{FAA}^{\text{R}(1) ce}{}_{\rho\nu}(p) + c^{eca} \tilde{\Gamma}_{FAA}^{\text{R}(1) be}{}_{\nu\rho}(k) = \\ & \tilde{\Gamma}_{AAw;N[\hat{\Delta}]}^{\text{R}(1) bca}{}_{\nu\rho}(k, p; 0). \quad (\text{C.9}) \end{aligned}$$

The explicit calculation of the right hand side of the last equation gives a result which contains spurious anomalies, *i.e.* which can be canceled by adding finite counterterms to

the action, this we have already seen in subsection 3.6., and a part which is the correct essential anomaly:

$$\begin{aligned}\tilde{\Gamma}_{AAw;N[\hat{\Delta}]}^{R(1)}{}_{\nu\rho}{}^{bca\text{ BM}}(k, p; 0) &= -\frac{1}{(4\pi)^2} \frac{1}{3} (p^2 - k^2) c^{bca} g_{\nu\rho} (T_R + T_L) \\ &\quad - \frac{1}{(4\pi)^2} \frac{2}{3} (k_\nu k_\rho - p_\nu p_\rho) c^{bca} (T_R + T_L) \\ &\quad + \frac{1}{(4\pi)^2} \frac{1}{3} \frac{d_{R-L}^{bca}}{2} \text{Tr} [\not{k} \not{p} \gamma_\nu \gamma_\rho].\end{aligned}$$

Notice that even if there $d_{R-L} = 0$ there are spurious anomalies in the calculation.

For the “naive” prescription, by using eq. (C.5) and *anticommuting the γ_5* , it is possible to write

$$\begin{aligned}\tilde{\Gamma}_{FAAA}^{R(1)}{}_{\nu\rho\mu}{}^{bca\text{ naive}}(k, p) &= i c^{bca} \frac{T_R + T_L}{2} T_{VVV}^{\nu\rho\mu}(k, p, -k - p) \\ &\quad + \frac{d_{R-L}^{bca}}{2} T_{VVA}^{\nu\rho\mu}(k, p, -k - p).\end{aligned}$$

Therefore, the left hand side of eq. (C.9) in the “naive” prescription (anticommuting γ_5) reads:

$$\begin{aligned}&c^{bca} \frac{T_R + T_L}{2} \{ (k + p)_\mu T_{VVV}^{\nu\rho\mu}(k, p, -k - p) - T_{VV}^{\nu\rho}(p) + T_{VV}^{\nu\rho}(k) \} \\ &- \frac{d_{R-L}^{bca}}{2} i (k + p)_\mu T_{VVA}^{\nu\rho\mu}(k, p, -k - p).\end{aligned}$$

But the first term between brackets vanish in the “naive” prescription without incurring into any inconsistency -no γ_5 is involved. Hence, we arrive at the conclusion that

$$\text{l.h.s. of eq. (C.9)} = \frac{d_{R-L}^{bca}}{2} i (-k - p)^\mu T_{VVA}^{\nu\rho\mu}(k, p, -k - p), \quad (\text{C.10})$$

in the “naive” dimensional regularization prescription (anticommuting γ_5). Next, one could think of applying eq. (C.8) to obtain an “anomaly” $\frac{1}{(4\pi)^2} \frac{d_{R-L}^{bca}}{2} \text{Tr} [\not{k} \not{p} \gamma_\nu \gamma_\rho]$, which is different from the correct Breitenlohner and Maison result. Moreover, eq. (C.10) was obtained by putting the one γ_5 that is left after the anticommutation process is done in the position μ . If we had put it in the position ν , we would not have obtained any anomaly at all. (The “anomaly” obtained with the first choice for a place for the γ_5 is three times the correct Breitenlohner and Maison result and three is the number of choices). This shows clearly that the “naive” prescription cannot give consistently the one-loop anomaly in chiral local gauge theories. Therefore, the claim that “we do not need discuss the graphs containing the AAA triangle since these behave in the same manner as the AVV triangle” of [30] seems to be not well justified.




Figure 14: one-loop 1PI Feynman diagrams with four bosonic legs

Let us move on to computation of the one-loop four gauge field vertex. This vertex also contributes to the anomaly. We have already shown in subsection 3.5. that the Breitenlohner and Maison scheme gives the correct relative factors between the anomaly from the three gauge field vertex and from four gauge field vertex.

The one-loop 1PI four gauge field function is given by the sum of the diagram depicted in fig. 14 and all the diagrams which are obtained from the first by performing all the possible permutations of three legs.

Proceeding as for the three boson vertex case, the difference between the renormalized minimal Breitenloner and Maison result for the diagrams of fig. 14 and the “naive” result defined with the by putting the γ_5 after the γ^λ -the result depends on the position of the γ_5 -

$$\begin{aligned}
& -\frac{i}{(4\pi)^2} \frac{2}{3} \left\{ g^{\mu\nu} g^{\rho\lambda} \left[3 \frac{T_{L+R}^{abcd}}{2} + 3 \frac{T_{L+R}^{abdc}}{2} + 3 \frac{T_{L+R}^{acdb}}{2} + 3 \frac{T_{L+R}^{adcb}}{2} - 5 \frac{T_{L+R}^{acbd}}{2} - 5 \frac{T_{L+R}^{adbc}}{2} \right] + \right. \\
& \quad g^{\mu\lambda} g^{\nu\rho} \left[3 \frac{T_{L+R}^{abcd}}{2} + 3 \frac{T_{L+R}^{acbd}}{2} + 3 \frac{T_{L+R}^{adbc}}{2} + 3 \frac{T_{L+R}^{adcb}}{2} - 5 \frac{T_{L+R}^{abdc}}{2} - 5 \frac{T_{L+R}^{adcb}}{2} \right] + \\
& \quad \left. g^{\mu\rho} g^{\nu\lambda} \left[3 \frac{T_{L+R}^{acbd}}{2} + 3 \frac{T_{L+R}^{acdb}}{2} + 3 \frac{T_{L+R}^{adbc}}{2} + 3 \frac{T_{L+R}^{abdc}}{2} - 5 \frac{T_{L+R}^{abcd}}{2} - 5 \frac{T_{L+R}^{adcb}}{2} \right] \right\} \\
& -\frac{i}{(4\pi)^2} \frac{1}{2} \text{Tr} [\gamma^\mu \gamma^\nu \gamma^\rho \gamma^\lambda \gamma_5] \left\{ \frac{T_{R-L}^{abcd}}{2} + \frac{T_{R-L}^{acdb}}{2} + \frac{T_{R-L}^{adbc}}{2} - \frac{T_{R-L}^{acbd}}{2} - \frac{T_{R-L}^{abdc}}{2} - \frac{T_{R-L}^{adcb}}{2} \right\}.
\end{aligned}$$

Notice that the first term between brackets is cyclic but that the second one is, in fact, $3 T_{R-L}^{a[bcd]}$, *i.e.* antisymmetric in the last three indices but not in the four indices, so, again, is not cyclic.

Then, the complete difference, taking into account all diagrams, is:

$$\begin{aligned}
& -\frac{i}{(4\pi)^2} \frac{2}{3} \left\{ g^{\mu\nu} g^{\rho\lambda} \left[3 \frac{T_{L+R}^{\{a,b\}\{c,d\}}}{2} - 5 \frac{T_{L+R}^{acbd}}{2} - 5 \frac{T_{L+R}^{adbc}}{2} \right] + \right. \\
& \quad g^{\mu\lambda} g^{\nu\rho} \left[3 \frac{T_{L+R}^{\{a,d\}\{c,b\}}}{2} - 5 \frac{T_{L+R}^{abdc}}{2} - 5 \frac{T_{L+R}^{adcb}}{2} \right] + \\
& \quad \left. g^{\mu\rho} g^{\nu\lambda} \left[3 \frac{T_{L+R}^{\{a,c\}\{b,d\}}}{2} - 5 \frac{T_{L+R}^{abcd}}{2} - 5 \frac{T_{L+R}^{adcb}}{2} \right] \right\} \\
& + \frac{i}{(4\pi)^2} \frac{1}{2} \text{Tr} [\gamma^\mu \gamma^\nu \gamma^\rho \gamma^\lambda \gamma_5] \left\{ \mathcal{D}_R^{abcd} - \mathcal{D}_L^{abcd} \right\},
\end{aligned}$$

where we have defined

$$T_{L+R}^{\{a,b\}\{c,d\}} = \text{Tr} \{T_L^a, T_L^b\} \{T_L^c, T_L^d\} + \text{Tr} \{T_R^a, T_R^b\} \{T_R^c, T_R^d\},$$

$$\mathcal{D}_L^{abcd} = -i 3! T_L^a T_L^{[b} T_L^c T_L^{d]} = \frac{1}{2} (d_L^{abe} c^{ecd} + d_L^{ace} c^{edb} + d_L^{ade} c^{ebc}) = \mathcal{D}_L^{a[bcd]},$$

and similarly $\mathcal{D}_R^{abcd}$.

Again, the first term between brackets is a local term which can be checked to be just proportional to the Feynman rules of $\text{Tr}_{R+L} A_\mu A^\mu A_\nu A^\nu$ and $\text{Tr}_{R+L} A_\mu A_\nu A^\mu A^\nu$; whereas the second term is not Bose symmetric and reflects the inconsistency of the “naive” prescription. Also, we can use these result to show that the *ambiguities in the “naive” calculation of the one-loop four boson vertex* is proportional to the epsilon tensor and to the coefficient $\mathcal{D}_R^{abcd} - \mathcal{D}_L^{abcd}$, which vanish if there is cancellation of anomalies. For example, the one-loop renormalized difference between the “naive” calculation with the γ_5 after $\gamma^\lambda P_{L(R)}$ and the “naive” calculation with the γ_5 after $\gamma^\mu P_{L(R)}$ is $i/8\pi^2 \varepsilon_{\mu\nu\rho\lambda} (\mathcal{D}_R^{abcd} - \mathcal{D}_L^{abcd} + \mathcal{D}_R^{bcda} - \mathcal{D}_L^{bcda})$.

We have exhibited by explicit computation the many inconsistencies that plague the computations leading to a renormalized one-loop chiral gauge theory if the “naive” Dimensional Regularization prescription is used. The classical action of the theory was given in subsection 3.1. To close this appendix we shall show that in spite of these inconsistencies, and if the chiral theory is anomaly free- a compulsory constraint-, the renormalized BRS invariant chiral theories obtained by means of the “naive” dimensional regularization prescription, on the one hand, and Breitenlohner and Maison Dimensional Regularization procedure, on the other, are equivalent at the one-loop level, in the sense that they are related by finite BRS symmetric renormalizations of the fields and couplings of the theory.

As we have previously compared the Breitenlohner and Maison renormalized theory with the “naive” renormalized theory when no finite BRS-asymmetric counterterms have been added to the former to restore the BRS symmetry, the comparison between the two renormalized theories, when the BRS-asymmetric counterterms have been added to the minimal Breitenlohner and Maison renormalized theory, is very easy.

Due to the fact that, in the situation of cancellation of anomalies, the “naive” minimal dimensionally renormalized theory satisfy the Slavnov-Taylor identities at order $\hbar^1$, the finite counterterms relating the two renormalized theories should appear in b -invariant combinations. And this is what nicely happen, upon inclusion of finite BRS asymmetric counterterms, the Breitenlohner and Maison renormalized theory equals the “naive” renormalized theory if the coefficients $-l_g^{(1)}$, $l_\psi^{(1)}$, $l_{\psi'}^{(1)}$, $l_A^{(1)}$ and $l_\omega^{(1)}$ - of the BRS-symmetric counterterms in eq. (69) are chosen to be given by

$$l_g^{(1)} = -\frac{1}{(4\pi)^2} \frac{5}{24} g^2, \quad l_{\psi^{(\prime)}}^{(1)} = \frac{1}{(4\pi)^2} g^2 C_{L(R)} [1 + (\alpha' - 1) \frac{5}{12}], \quad l_A^{(1)} = l_\omega^{(1)} = 0.$$

This corresponds, of course, to a mass-independent multiplicative finite renormalization of the fields and parameters of the effective action. Hence, the beta function and anomalous dimensions of both renormalized theories are the same at the one-loop level. The finite renormalizations we have just mentioned read

$$\begin{aligned}
A &= A^*, & B &= B^*, & \omega &= \omega^*, & \bar{\omega} &= \bar{\omega}^*, \\
\rho &= \rho^*, & \zeta &= \zeta^*, & \alpha &= \alpha^*, \\
g &= (1 + \frac{1}{(4\pi)^2} \frac{5}{48} g^{*2} \hbar^1) g^*, \\
\psi &= (1 + \frac{1}{(4\pi)^2} g^{*2} C_L [1 + (\alpha' - 1) \frac{5}{12}] \hbar^1 P_L) \psi^*, \\
\bar{\psi} &= \bar{\psi}^* (1 + \frac{1}{(4\pi)^2} g^{*2} C_L [1 + (\alpha' - 1) \frac{5}{12}] \hbar^1 P_R), \\
\psi' &= (1 + \frac{1}{(4\pi)^2} g^{*2} C_R [1 + (\alpha' - 1) \frac{5}{12}] \hbar^1 P_R) \psi'^*, \\
\bar{\psi}' &= \bar{\psi}'^* (1 + \frac{1}{(4\pi)^2} g^{*2} C_R [1 + (\alpha' - 1) \frac{5}{12}] \hbar^1 P_L), \\
\bar{L} &= \bar{L}^* (1 + \frac{1}{(4\pi)^2} g^{*2} C_L [1 + (\alpha' - 1) \frac{5}{12}] \hbar^1 P_L), \\
L &= (1 + \frac{1}{(4\pi)^2} g^{*2} C_L [1 + (\alpha' - 1) \frac{5}{12}] \hbar^1 P_R) L^*, \\
\bar{R} &= \bar{R}^* (1 + \frac{1}{(4\pi)^2} g^{*2} C_R [1 + (\alpha' - 1) \frac{5}{12}] \hbar^1 P_R), \\
R &= (1 + \frac{1}{(4\pi)^2} g^{*2} C_R [1 + (\alpha' - 1) \frac{5}{12}] \hbar^1 P_L) R^*.
\end{aligned}$$

Where the fields and constants with a star are the renormalized fields and constants in the “naive” minimal Dimensional Regularization prescription and the fields and constant without stars stand for the corresponding renormalized quantities in renormalized theory obtained by means of the Breitenlohner and Maison scheme (minimal subtraction plus addition of the finite counterterms in eq. (69), with $l_g^{(1)}$, $l_\psi^{(1)}$, $l_{\psi'}^{(1)}$, $l_A^{(1)}$ and $l_\omega^{(1)}$ set to zero).

Acknowledgments

The financial support from the Universidad Complutense de Madrid under grant PR156/97-7164 and the Consejería de Educación y Cultura of the Junta de Comunidades de Castilla-La Mancha are acknowledged.

References

- [1] J. Ashmore, Lett. Nuovo Cimento 4 (1972) 289; C.G.Bollini and J.J. Giambiagi, Phys. Lett. B40 (1972) 566.
- [2] G. t’Hooft and M. Veltman, Nucl. Phys. B44 (1972) 189-213.
- [3] P. Breitenlohner and D. Maison, Comm. Math. Phys 52 (1977) 11-38.
- [4] P. Breitenlohner and D. Maison, Comm. Math. Phys 52 (1977) 39-54.
- [5] P. Breitenlohner and D. Maison, Comm. Math. Phys 52 (1977) 55-75.
- [6] G. Bonneau, Nucl. Phys. B167 (1980) 261-284.
- [7] G. Bonneau, Nucl. Phys. B180 (1980) 477-508.
- [8] G. Bonneau, Nucl. Phys. B177 (1980) 523-527.
- [9] G. Leibbrandt, Rev. Mod. Phys. 47 4 (1975) 849.
- [10] J. Collins, Renormalization (Cambridge University Press, Cambridge, 1984).
- [11] G. ‘t Hooft, Nucl. Phys. B61 (1973) 455.

- [12] G. 't Hooft, Nucl. Phys. B62 (1973) 444; O.V. Tarasov, A.A. Vladimirov and A.N. Zharkov, Phys. Lett. B93 (1980) 65; F.V. Tachkov, Phys. Lett. B100 (1981) 65; D.I. Kazakov, Phys. Lett. B133 6 (1983) 406; J.M. Campbell, E.W.N. Glover and D.J. Miller, Nucl. Phys. B498 (1997) 397; L. Brucher, J. Franzkowski and D. Kreimer, *Oneloop 2.0 hep-th 9709209*.
- [13] M.E. Machacek and M.T. Vaughn, Nucl. Phys. B222 (1983) 83; *ibid.* B236 (1984) 221; I. Jack and D.R.T. Jones, Nucl. Phys. B249 (1985) 472; C. Ford, I. Jack and D.R.T. Jones, Nucl. Phys. B387 (1992) 373; C. Ford, D.R.T. Jones, P.W. Stephenson and M.B. Einhorn, Nucl. Phys. B395 (1993) 17; T. Van Ritbergen and J.A.M. Vermaseren, Phys. Lett. B400 (1997) 379; B.A. Kniehl, Int. J. Mod. Phys. A10 4 (1995) 443 *and references therein*.
- [14] G. Bonneau, Phys. Lett. B96 (1980) 148-150.
- [15] D.A. Akyeampong and R. Delbourgo, Nuovo Cimento 17A (1973) 578; *ibid.* 18A (1973) 94; *ibid.* 19A (1974) 219.
- [16] D. Kreimer, Phys. Lett. B237 (1990) 59, "The role of γ_5 in Dimensional Regularization", *hep-ph/9401354*.
- [17] J.C. Collins, Nucl. Phys. B92 (1975) 477.
- [18] O. Piguet and S. Sorella, Algebraic renormalization (Springer-Verlag, Berlin, 1995) *and references therein*.
- [19] F. Brandt, Commun. Math. Phys. 190 (1997) 459; E. Krauss, Renormalization of the Electroweak Standard Model to all orders, *hep-th/9709154*, *to appear in Ann. Phys.*
- [20] K. Hepp, Renormalization theory in statistical mechanics and quantum field theory, in "Les Houches XX 1970" (Gordon and Breach, New York, 1971).
- [21] G. Bonneau, Int. Journ. Mod. Phys. A5 (1990) 3831-3859.
- [22] W.A. Bardeen, R. Gastmans and B. Lautrup, Nucl. Phys. B46 (1972) 319; M. Chanowitz, M. Furman and I. Hinchliffe, Nucl. Phys. B159 (1979) 225.
- [23] L. Alvarez-Gaumé and E. Witten, Nucl. Phys. B234 (1984) 269.
- [24] K. Wilson, Phys. Rev. D7 (1973), 2924-2926.
- [25] O. Piguet and A. Rouet, Phys. Rep. 76 (1981) 1-77.
- [26] J.G. Körner, N. Nasrallah and K. Schilcher, Phys. Rev. D41 (1990) 888-890.
- [27] R. Ferrari, A. Le Yaouanc, L. Oliver and J.C. Raynal, Phys. Rev. D52 (1995) 3036-3047.
- [28] A. Barroso, M.A. Doncheski, H. Grotch, J.G. Körner and K. Schilcher, Phys. Lett. B261 (1991) 123-126.
- [29] S. Gottlieb and J.T. Donohue, Phys. Rev. D20 (1979) 3378-3389.
- [30] B.A. Ovrut, Nuc. Phys. B213 (1983) 241-265.
- [31] C. Itzikson and J.B. Zuber, Quantum Field Theory (McGraw-Hill, 1980)
- [32] M. Bos, Ann. Phys. 181 (1988) 197; H. Osborn, Ann. Phys. 200 (1990) 1; I. Jack, D.R.T. Jones and K.L. Roberts, Z. Phys. C 63 (1994) 151.